

1 Use of simulation approaches to evaluate the consequences of catch-and-release angling on the
2 migration behaviour of adult Atlantic salmon (*Salmo salar*)

3
4
5
6
7
8

9 Robert J. Lennox^{1,2,§}, Steven J. Cooke¹, Ola H. Diserud², Torgeir B. Havn², Martin R. Johansen²,
10 Eva B. Thorstad², Frederick G. Whoriskey³, and Ingebrigt Uglem²

11
12
13
14
15
16

¹Fish Ecology and Conservation Physiology Laboratory, Department of Biology, Carleton
University, Ottawa, Ontario, Canada K1S 5B6 ²Norwegian Institute for Nature Research, P. O.
Box 5685, Sluppen, N-7485 Trondheim, Norway ³Ocean Tracking Network, c/o Dalhousie
University, Halifax, NS B3H 4J1, Canada

17 [§]Corresponding Author: Email: robert.lennox@carleton.ca; telephone: 1-613-408-3474

18 Lennox, Robert J.; Cooke, Steven J.; Diserud, Ola Håvard; Havn, Torgeir Børresen; Johansen, Martin R.; Thorstad, Eva Bonsak; Whoriskey,
Frederick G.; Uglem, Ingebrigt.
19 Use of simulation approaches to evaluate the consequences of catch-and-release angling on the migration behaviour of adult Atlantic salmon (*Salmo salar*). *Ecological Modelling* 2016 ;Volum 333. s. 43-50 DOI 10.1016/j.ecolmodel.2016.04.010 (CC BY-NC-ND 4.0)

20 Abstract

21

22 Given that most salmon released by anglers survive (97% in this study), economically and
23 culturally important recreational Atlantic salmon fisheries are increasingly incorporating catch-
24 and-release. Sublethal alterations to behaviour with potential individual fitness costs are a potential
25 consequence of catch-and-release but are difficult to measure empirically relative to uncaptured
26 fish. To test for sublethal effects of angling on migratory movements, 39 salmon were captured by
27 recreational anglers, externally tagged with radio transmitters, and released. Data from the annual
28 visual drift count of spawning salmon were used to calculate the probability of spawning in each
29 pool of the river and input into simulation models. Simulation models were implemented to test
30 the hypothesis that catch-and-release did not affect the upriver movement of salmon. Ten thousand
31 simulation steps selected a spawning pool for each of the tagged salmon, permitting a calculation
32 of the average expected movement by salmon for comparison to the average observed movement.
33 The average observed movement by the released salmon was significantly less than the average
34 expected movement generated by all three models, indicating a sublethal effect of catch-and-
35 release on the migration of Atlantic salmon.

36

37 Keywords: telemetry, recreational fisheries, sublethal effects, angling, simulation

38

39 1.0 Introduction

40 Atlantic salmon migration in freshwater incorporates multiple phases of activity
41 including active upriver movement, holding, and searching with upstream and downstream
42 movements before staging near the eventual spawning destination weeks or months in advance
43 of spawning (Økland et al. 2001). Atlantic salmon are philopatric with most individuals able to
44 locate their natal rivers (Fleming 1996) and even specific tributaries within a system (Heggberget
45 et al. 1988; Verspoor et al. 1991). The timing and speed of migration by Atlantic salmon through
46 freshwater depends on a variety of factors, including sex (Lucas et al. 1993), size (Kristinsson et
47 al. 2015), and experience (Niemelä et al. 2006). However, anthropogenic challenges including
48 pollution (Thorstad et al. 2005), artificial barriers (Croze 2008), and climate change (Baisez et al.
49 2011) alter migratory patterns exhibited by salmon. In addition, recreational fishery practices
50 such as catch-and-release have the potential to influence the migratory behaviour of salmon in
51 rivers.

52 Recreational fisheries are popular worldwide and can be important components of the
53 economy for many communities (Arlinghaus and Cooke 2009). The sustainability of recreational
54 fisheries, however, depends on the ability of the targeted fish population to persist in spite of
55 harvest and non-harvest mortality imposed by angling activities (Coggins et al. 2007; Cooke and
56 Schramm 2007). Traditionally, many recreational anglers harvested their catch; however, catch-
57 and-release is now increasing in many fisheries. From a regulatory perspective, catch-and-release
58 focuses on maintaining the socio-economic benefits of fisheries while sustaining fish populations
59 that are being exploited. As a result, catch-and-release practices assume that fish released by
60 anglers have high survival and experience limited sublethal consequences to their lifetime
61 reproductive success (Arlinghaus et al. 2007; Wilson et al. 2014). Catch-and-release is

62 increasingly practiced in recreational salmon fisheries but scientific evaluations of catch-and-
63 release for salmon have focused on demonstrating that mortality for caught and released fish is
64 infrequent (< 0.10 ; Thorstad et al. 2003; Gargan et al. 2015). However, mortality studies alone
65 probably underestimate the impacts of catch-and-release because they do not consider sublethal
66 effects (Cooke et al. 2002). Sublethal effects occur as a consequence of aerobic debt (Kieffer
67 2000; Lee et al. 2003), metabolic disturbance (Wood et al. 1983), physiological stress (Pankhurst
68 2011), and exhaustion induced by angling. The resulting prolonged recovery can result in
69 behavioural impairment, causing significant indirect and direct impairments to potential fitness,
70 such as reduced growth or fecundity (Cooke et al. 2002; Wilson et al. 2014). Sublethal effects of
71 catch-and-release can be difficult to measure because equating a capture event to fitness is
72 challenging. However, migrating salmonids provide a useful model for identifying sublethal
73 effects of angling because the upriver migration towards spawning grounds might be a reflection
74 of fitness (Dingle 1980).

75 There is correlative evidence that angling alters migration patterns of Atlantic salmon.
76 Two documented alterations to migratory patterns that have been observed for Atlantic salmon
77 released by anglers are downriver movement from the release site (Mäkinen et al. 2000; Thorstad
78 et al. 2003; Havn et al. 2015) and shortened migration distance (Tufts et al. 2000; Lennox et al.
79 2015a). However, the extent to which catch-and-release actually causes significant changes to an
80 individual's migration is unclear. Determining whether migration is negatively affected by
81 angling requires an estimation of where salmon would spawn if they were not captured by
82 anglers. It is difficult to know where salmon are destined to spawn in the river prior to the
83 spawning period itself, necessitating the development of a novel tool using an estimate of the
84 spawning distribution of non-angled fish within the river as a proxy for the ultimate distribution

85 of released fish at spawning time. This information provides a natural baseline against which
86 hypotheses regarding the hypothesized impacts of catch-and-release can be tested. To do so, we
87 tested the observed movement of salmon against model-predicted movement given no effect of
88 catch-and-release. Model predictions were generated from the distribution of salmon at spawning
89 time based on the results of a passive drift count. These were compared to the upriver progress
90 and spawning locations used by Atlantic salmon after catch-and-release as determined by radio
91 telemetry.

92 2.0 Methods

93

94 2.1 Study Area

95

96 River Lakselva is a 45 km long river that drains into the Porsangerfjord in Porsanger,
97 Finnmark, Norway. The confluence of Lakselva with the fjord is at 70.078757 N, 24.926302 E.
98 Lakselva is a large, unregulated river with one major tributary (Vuolajohka) and two large lakes
99 (Figure 1). Atlantic salmon enter Lakselva during the spring and summer and spawn in Lakselva
100 and Vuolajohka in October. The recreational fishery is regulated by the Lakselva Landowner's
101 Association, which limits access to most of the fishery via a licensing system. There are also
102 stretches of river where angling is regulated by single landowners or local lodges. The annual
103 salmon fishing season in Lakselva begins June 1 and continues through August 31. Average
104 annual catch in Lakselva (2007-2015) is 1464 ± 229 (SD) Atlantic salmon (www.scanatura.no).

105

106 2.2 Tagging

107

108 Historical catch records indicate that few salmon enter this river in June; therefore, we
109 focused our tagging efforts between July 13 and August 28, 2014. Salmon selected for tagging
110 (N = 39) were those that were typical of caught-and-released fish, and not moribund (see Lennox
111 et al. 2015a). After being landed by an angler, salmon were transferred to a water-filled tube
112 where they were placed in a prone position. The individual was measured and a radio transmitter
113 in the frequency range 142.114 – 142.213 (Advanced Telemetry Systems [ATS], Minnesota,
114 USA) was attached externally below the dorsal fin. The tagging methods followed Lennox et al.
115 (2015a), using sterile hypodermic needles and stainless steel wire to secure the radio tag through
116 the dorsal musculature. Anglers that captured salmon handled them naturally and we did not
117 attempt to interfere with their fish handling (e.g. by telling them to use a net, not to air expose the
118 fish too long, etc.). However, we declined to tag two angled salmon; one salmon was critically
119 injured (hooked in gills) and the other was too small to support the tag comfortably. In total, 39
120 Atlantic salmon (89 ± 16 cm TL, range: 62 – 121 cm) captured by anglers were radio tagged and
121 released. Many (N = 18) of the salmon were caught and released in pools relatively close to the
122 head of the tide and most (N = 26) were considered to be fresh fish based on their silver colour
123 and/ or the presence of salmon lice. Mean water temperature at capture was 14 ± 1 °C whereas
124 temperature stress begins to become an important issue in Atlantic salmon angling at > 20 °C
125 (Dempson et al. 2002, Havn et al. 2015). All handling and tagging were conducted according to
126 Norwegian regulations for treatment and welfare of animals.

127

128 2.3 Tracking

129

130 To ensure adequate coverage of the watershed, four stationary data logging stations were
131 set up at key points in the river to monitor passage of salmon. Data logging receivers (Advanced
132 Telemetry Systems [ATS], Minnesota, USA; R4520CD Coded Receiver-Datalogger) were set up
133 with paired Lotek (Newmarket, Canada; 6 element Yagi tuned to 142 MHz) antennas (one
134 pointing upriver and one pointing downriver) to establish directionality of movement by salmon
135 past the receiver. The stations were set up above and below each of the lakes and also near the
136 mouth of the tributary Vuolajohka (Figure 1). The listening stations were checked biweekly and
137 were active throughout the summer and into the autumn. In addition to the stationary logging
138 stations, mobile tracking was conducted along the river using a vehicle mounted receiver and a
139 magnetic whip antenna (Magnetic Roof-Mount Dipole, Laird Technologies, Missouri, USA).
140 Salmon positions were determined on alternating days starting on July 14 and continuing through
141 the end of the angling season on August 31, 2014. During the autumn, positioning occurred on
142 September 2, September 15, September 24, and October 24. We used the salmon's position on
143 September 24 as an estimate of the spawning position in the river. On September 24-25, a
144 snorkel survey was conducted in conjunction with radio tracking to visually confirm survival of
145 some salmon with nominal movement after release.

146

147 2.4 Drift Count

148

149 Each year in Lakselva the Landowner's Association conducts a visual count to estimate
150 the total number of salmon in the river. The count is conducted by two experienced persons who
151 drift passively downriver while snorkeling. For each section of the river (typically delineated by
152 pools), the number of spawning salmon is estimated based on these visual observations.

153 Although drift counts are considered underestimates of the total number of salmon, Orell and
154 Erkinaro (2007) found that they provided accurate estimates of spawning biomass during the
155 salmon spawning season. In 2014, the drift count in Lakselva was conducted on September 13-
156 14, and spawning was observed to have commenced (E. Liberg, personal communication). Staff
157 were aware of and noted the presence of tagged salmon based on visual identification of the
158 external radio tags. We collected drift count data from Lakselva for 2011, 2013, 2014, and 2015.

159

160 2.5 Data Analysis

161

162 Each pool in the drift count was assigned a number with the pool closest to the fjord
163 being Pool 1 and the pool farthest upriver being number 57 (Figure 1). Pools that could not be
164 enumerated by divers due to poor visibility were assigned 0 salmon for the purposes of analysis.
165 The release and spawning pools (the latter being inferred from the position of the salmon in the
166 river on September 24) were compared to assess the movement of salmon released by anglers.
167 The analyses could be conducted on 30 of the 39 tagged salmon, because one died, one exited
168 the river, and seven were recaptured and killed by anglers prior to spawning season. We used a
169 Pearson correlation to quantify the relationship between the salmon's release and spawning
170 pools. To test whether catch-and-release affected the movement of salmon within the river, a
171 series of simulations was conducted to create a distribution of the most probable average
172 movement of salmon from the release site under the null hypothesis of no effect of catch-and-
173 release.

174 The simulation tests were implemented as follows: each pool was assigned a probability
175 that a salmon would spawn there based on the proportion of salmon observed spawning near

176 there by the 2014 drift count. These pool probabilities were calculated and applied to each of the
177 30 radio tagged salmon. A single simulation step was implemented using the *sample* function in
178 R (R Core Team 2014), which selected a spawning pool for each salmon based on the assigned
179 probabilities, permitting a calculation of expected movement by subtracting the number of the
180 release pool from the number of the simulated spawning pool. For example, a fish captured and
181 released in Pool 1 could be assigned Pool 10 as a spawning pool in a simulation step, equating to
182 an expected movement of nine pools. Averaging the expected movement among the 30 salmon
183 and repeating the simulation 10,000 times, a probability distribution was generated that described
184 the average expected movement of salmon from the site of their release to spawn. The average
185 expected movement was then compared to the average observed movement of the 30 radio
186 tracked salmon. The total number of simulated movement values greater or equal to the observed
187 mean movement value was divided by the number of simulations (10,000), yielding a probability
188 (p-value) that the average observed movement differed from the average expected movement.

189 We ran three simulations each using different assumptions (described below) and
190 generating different null models. All null models assumed that there was no impact of being
191 caught and released on a salmon's movement.

192 Finally, we present data from the drift count in Lakselva for 2011, 2013, 2014, and 2015
193 to assess temporal stability in the distribution of spawning salmon within the river. We used
194 violin plots as implemented by *ggplot2* (Wickham 2009), which show the density of spawners
195 along the longitudinal axis of the river. To test for differences in the average spawning position
196 across years we used a Kruskal-Wallis non-parametric analysis of variance.

197

198 2.5.1 Free distribution

199

200 In the first simulation, radiotagged salmon were assumed in the null model to distribute
201 anywhere in the river to spawn, independently of where they were caught and released. The
202 probability of choosing a given spawning pool was estimated as the proportion of the total
203 number of spawners in the river observed in this particular pool during the drift count. This
204 corresponds to assuming that salmon will freely distribute in a river and concentrate in some
205 areas, presumably of high spawning substrate. Although it is well known that salmon are
206 positively rheotactic and migrate primarily upriver to spawning sites, this simulation assumed
207 that no matter where salmon were captured, they could in theory move up or down independent
208 of the release location by maintaining equal spawning pool probabilities for all salmon.

209

210 2.5.2 Salmon only move upriver

211

212 In the second simulation, spawning pool probabilities were adjusted based on the release
213 pool for each radiotagged salmon such that any pools downriver of the release pool had zero
214 probability of salmon spawning there and upriver pool spawning probabilities were adjusted
215 accordingly for each fish.

216

217 2.5.3 Most salmon move upriver

218

219 The third simulation was identical to the second, with the exception that it excluded
220 salmon that spawned at or below the release site. This restricted the simulation to 15 salmon that
221 spawned at least one pool upriver from the release location. Fifteen salmon that spawned at or

222 below the release pool were excluded under the assumption that these fish were captured after
223 completing their migration whereas the other 15 were captured during their upriver migration.

224

225 3.0 Results

226

227 3.1 Catch-and-release

228

229 Only one of the 39 tagged salmon is known to have died. This occurred soon after catch-
230 and-release, and its drifting carcass was observed by an angler downriver of the release site just
231 hours later (E. Liberg, pers.comm.). Therefore, survival from catch-and-release was high (0.97 of
232 released fish). Total mortality (N = 2) from angling was 0.95 (total N = 40) after including one
233 moribund salmon that was not released because of bleeding. One tagged salmon left from the
234 river in August, which was a grilse (i.e. one-sea-winter salmon) that had exhibited erratic
235 behaviour after release, first moving upriver within hours of release and eventually moving
236 downriver two kilometres below the initial release site before exiting in August (last tracked
237 August 24), several weeks prior to the spawning period. Given the movement trajectory of that
238 salmon, it was determined that it had survived catch-and-release but we were unable to test
239 whether its river exit was associated with catch-and-release or whether it left the river to spawn
240 in another, adjacent river (Havn et al. 2015). Nine salmon (0.23) were reported as having been
241 recaptured by anglers later in the angling season, with seven of them being harvested and two re-
242 released. One of the seven harvested salmon was recaptured twice before being killed. Two
243 tagged salmon that were captured multiple times remained in the river through the spawning

244 season. One of the recaptured salmon was angled as a kelt the year after tagging on June 20,
245 2015.

246

247 3.2 Spawning distribution of catch-and-release salmon

248

249 There was a strong positive correlation between the catch-and-release location and the
250 final spawning position, indicating that there was limited upriver movement ($R^2 = 0.74$ Figure 2).
251 During the spawning period, all of the salmon that were still present in the river were located in
252 regions of the river known to be spawning locations for salmon. In addition, 20 (0.71) of the
253 tagged salmon were visually identified in spawning aggregations during the drift count. The
254 Lakselva Landowners' Association counted 1341 salmon spawning in the main stem of Lakselva
255 during the autumn spawning count in 2014. The drift count was conducted in 72 pools in the
256 river, which we reduced to 57 pools for analysis based on the locations of pools in the river and
257 counts from previous seasons. According to the drift count, the majority of salmon spawned
258 below the lakes, with only ten salmon counted above Øvrevatnet. However, there were some
259 areas in the river that were too turbid for the counting staff to conduct the count, making some
260 areas of the river appear depauperate in the count. Most notably, sections of the river between
261 Øvrevatnet and Nedrevatnet were not counted to poor visibility, nor was the tributary
262 Vuolajohka. However, given that these regions were upstream of where all the tagged salmon
263 spawned we suggest that this would not affect our results.

264

265 3.3 Simulation tests

266

267 3.3.1 Free distribution

268

269 When the simulation permitted salmon to distribute themselves anywhere within the river
270 to spawn, salmon were predicted to move on average 7.04 pools upriver from the catch-and-
271 release site (Figure 3A). In other words, a theoretical 30 salmon released in the given pools
272 (Table 1) would move on average 7.04 pools each toward spawning grounds if they were
273 assumed to freely distribute themselves as the wild fish in the river did. This was mostly because
274 the majority of radio tagged fish were captured in lower reaches of the river and would therefore
275 be most likely to move upriver where the majority of the salmon were counted during the drift
276 count. Based on fish positions from tracking data from September, the tagged salmon moved on
277 average only 2.33 pools upriver from the release site, significantly less than expected based on
278 the free distribution hypothesis ($p = 0.03$).

279

280 3.3.2 Salmon only move upriver

281

282 When salmon in this null model were restricted from backtracking to downriver
283 spawning grounds, the simulation indicated that salmon should move on average 20.01 pools
284 upriver from the release location. This makes sense because many fish were captured in the
285 lower parts of the river and would therefore be highly likely to migrate to middle or upper
286 reaches for spawning where the highest numbers of wild fish were found during the visual
287 counts. However, as noted above the radiotracked fish showed limited movement. In this
288 simulation, where downstream movements were discounted and assigned 0 values, the average

289 movement was 2.33 pools per individual, again a highly significant difference from the model's
290 prediction ($p = 0.00$; Figure 3B).

291

292 3.3.3 Most salmon move upriver

293

294 When the second simulation was repeated excluding all salmon that showed any
295 downriver movements, we found that the simulation reduced the predicted movement per fish to
296 only 9.95 pools upriver per individual. For the radio tracked sample, after removing the salmon
297 that moved downriver, the observed movement was 6.07 pools per individual, still a highly
298 significant difference compared to the model's expected movement ($p = 0.01$; Figure 3C).

299

300 3.4 Seasonal differences in drift count observations

301

302 Average spawning pools were calculated from historic drift counts and it was determined
303 that the average spawning pools in Laksevla were 30 in 2011 ($N = 849$), 25 in 2013 ($N = 1254$),
304 21 in 2014 ($N = 1337$), and 26 in 2015 ($N = 832$). We observed some temporal inconsistency in
305 the distribution of spawning salmon within Lakselva (Figure 4). Indeed, there was a significant
306 difference in the distribution of spawners across years ($\chi^2 = 250.22$, $df = 3$, $p < 0.01$). However,
307 visual analysis (Figure 4) demonstrated consistent shapes in the distribution of spawning salmon
308 and indicated that the majority of spawning salmon are consistently below Pool 49, which was
309 the last pool prior to the first lake, Nedrevatnet (Figure 1). Moreover, most salmon in the river
310 spawned in pools in the middle of the anadromous stretch of the river.

311

312 4.0 Discussion

313

314 Similar to other studies on the effects of catch-and-release angling on Atlantic salmon, we
315 identified high survival of the fish released by anglers. One mortality among 39 salmon
316 represents a high probability of survival for salmon given good angling practices. Interestingly,
317 we calculated an exceptionally high recapture rate of salmon in Lakselva. Generally, instances of
318 recapture are infrequent in salmon fisheries and Lennox et al. (2015b) calculated a recapture
319 frequency of about 0.18 from multiple Norwegian rivers (including Lakselva) in 2012-2013. In
320 2014, 0.23 salmon were recaptured in Lakselva including one individual that was recaptured
321 twice (but counted in the proportion only once) and excluding one individual that was recaptured
322 as a kelt the following summer. This frequent recapture is interesting because there have been no
323 studies on the effects of multiple capture on salmon during their spawning migration, perhaps
324 because it is considered to be an infrequent occurrence. Some individuals tend to have higher
325 vulnerability to angling than others and would be captured more frequently than expected by
326 chance (Cox and Walters 2002; Tsuboi and Morita 2004). However, Lennox et al. (2015a) found
327 that salmon in a control group (captured prior to river entry by bag net) were not captured less
328 frequently than salmon that had already been captured by anglers. That a relatively high
329 proportion of salmon released by anglers goes on to be recaptured begs questions about how
330 effective catch-and-release can be in some fisheries with high exploitation rates (e.g. Gudjonsson
331 et al. 1996; Downton et al. 2001). High recapture of salmon suggests that further research is
332 necessary to evaluate the physiological and behavioural effects of recapture for salmon during a
333 potentially physiologically sensitive life stage. Indeed, encounters with recreational anglers are
334 stressful for fish in the short-term. Burst exercise during angling increases the concentration of

335 circulating stress hormones and results in osmoregulatory disruptions (Wood 1991; Kieffer 2000;
336 Barton 2002). After release, there is an energetic burden associated with repayment of oxygen
337 debt (Scarabello et al. 1991) and restoration of intramuscular fuels (Kieffer 2000).

338 Our simulation models indicated that the caught and released salmon in Lakselva had
339 shorter migrations than expected from model inputs. Two other studies have identified reduced
340 migratory distances traveled by salmon as a sublethal consequence of catch-and-release (Lennox
341 et al. 2015a; Tufts et al. 2000). However, these studies used a reference group of radio tagged
342 fish that had been captured using means other than angling (traps or nets), and these capture
343 methods could also have stressed the fish potentially confounding their utility as controls. The
344 novel approach of this study used uncaptured fish from a passive count rather than a control
345 group that had potentially been subjected to stress to generate a more robust estimate of expected
346 movement by released salmon.

347 Although we identified a sublethal effect of angling on Atlantic salmon, it is not clear
348 what the impacts of movement reductions would have on individual fitness and salmon
349 population dynamics. For Atlantic salmon released by anglers, reduced upriver migration
350 resulting from catch-and-release has the potential to decrease fitness via density-dependent egg
351 or fry mortality (Einum and Nislow 2005) or via outbreeding effects when salmon do not
352 successfully reach their natal spawning destination (Heggberget et al. 1986). However, the extent
353 of genetic substructuring by Atlantic salmon within rivers is probably low in general (Garant et
354 al. 2000) particularly within smaller rivers such as Lakselva without major tributaries (Jordan et
355 al. 1992; Vähä et al. 2011). It could be suggested that short migrations are symptomatic of larger
356 disturbances associated with stress or exhaustion given that breeding success is influenced by
357 physiological condition on spawning grounds (de Gaudemar and Beall 2004; Hendry and Beall

2004). However, other studies of released salmon have found that parr densities increased in years following catch-and-release (Whoriskey et al. 2000; Thorstad et al. 2003), that late season catch-and-release does not affect gamete or fry quality (Davidson et al. 1994; Booth et al. 1995), and that wild salmon released by anglers are able to successfully reproduce (Richard et al. 2013). Nonetheless, if reduced migration following catch-and-release corresponds to reduced activity overall, there could be fewer reproductive encounters by released salmon corresponding with decreased fitness. Even though salmon in this study did not travel as far as was expected based on the simulation, every salmon (except the one that exited the river prematurely and the one that died) was tracked at suitable spawning territory and many were also visually observed in aggregations of spawning conspecifics during drift counting.

An alternative explanation for our findings is that the salmon captured by anglers never intended to continue migrating because they were in the holding phase of migration (Økland et al. 2001). This implies that salmon are more likely to be captured by anglers at the end of migration than during the upriver migration phase. Vulnerability to recreational angling is a complex function of the biotic and abiotic environment (Stoner 2004), individual-level characteristics (Cooke et al. 2007), and the fisheries environment (i.e. gear types used; Lennox et al. 2015b). However, changing vulnerability to angling at different stages of fish migration has not previously been explored; however, behaviour does change at different stages of the migration, which has the potential to influence angling vulnerability. For example, dominant males become aggressive on spawning grounds (Hendry and Beall 2004), a behavioural change that could influence vulnerability to angling. Therefore, behavioural vulnerability could increase when salmon arrive at spawning grounds and indeed many fish remain in holding pools near spawning grounds for long periods of time prior to spawning (Økland et al. 2001) meaning that

381 salmon spend most of their time in freshwater at or near their spawning sites. This suggests that
382 angling vulnerability – and capture probability – should be higher on spawning grounds than
383 during the migration and that the “shortened migration” we observed was actually a function of
384 this change in capture probability.

385 Combining a visual survey with the radio telemetry in this study proved important for
386 estimating survival of salmon after catch-and-release. We had several salmon exhibit limited
387 post-release movement, including some that would have been categorized as dead using
388 established protocols for the interpretation of electronic tagging data (Lennox et al. 2015a) based
389 on their lack of movement, that were confirmed to be alive via visual observation. Indeed,
390 telemetry studies can also underestimate the movement of animals (Ovidio et al. 2000),
391 particularly without fine-scale positioning systems (Hanson et al. 2007). Although we are
392 confident that our periodic tracking allowed us to accurately identify the movements of salmon at
393 a coarse scale (i.e. among pools), it is possible for salmon to make forays up or downriver in
394 short periods of time that could have been missed (i.e. searching behaviour; Økland et al. 2001).
395 For example, one salmon tagged in Pool 2 was tracked once in Pool 5; however, it returned to
396 Pool 2 before the next tracking and remained there until spawning. Such transient movements
397 can only be detected by chance when tracking is periodic. Moreover, Taggart et al. (2001) noted
398 that salmon may move up to 5 km between redds during the spawning season. Although we
399 accept that our methods may not have captured all movements caught and released salmon made,
400 the overall trend observed among salmon was striking because upriver movement was largely
401 restricted throughout the remainder of the summer and into the spawning season.

402 Using simulation methods to test hypotheses about salmon movement was a novel
403 approach for answering our research question. Salmon are dynamic animals and although well

404 studied, their behaviour remains somewhat cryptic. Simulation provided an analytical tool for
405 exploring different but equally rational hypotheses to develop models of expected movement by
406 the released salmon. Although we found that there was some inconsistency in the spawning
407 distribution of salmon in Lakselva across years, it was interesting and important to our study to
408 note that general trends were similar. Ultimately, the results of all three simulations were
409 concordant allowing us to make inferences about the population that we studied. Annual visual
410 spawning counts of fish similar to those that we used to generate spawning pool probabilities are
411 available for many rivers making this method a valuable tool for work over and beyond stock
412 assessment in the future.

413

414 5.0 Conclusion

415

416 Consistent with other studies, high survivorship of salmon released by anglers in
417 Lakselva is promising for salmon conservation efforts and demonstrates the utility of catch-and-
418 release for management of the salmon fishery. However, our model predicted longer migrations
419 after catch-and-release than we observed, suggesting that the upriver migration could have been
420 hindered by angling, which could be a relevant sublethal effect of catch-and-release. Future
421 research into the behavioural vulnerability of salmon at different stages of migration are
422 necessary to develop a mechanistic understanding of these observations. Moreover, studies that
423 monitor the fitness-related endpoints of released salmon could provide important information
424 about the effects of catch-and-release on reproduction including gamete development prior to
425 spawning and intraspecific competition for mating opportunities or fertilization success.

426

427 Acknowledgements

428 We thank Egil Liberg and the Lakselva River Owner's Association for their support of this study.

429 R JL was funded by a Natural Sciences and Engineering Research Council (NSERC) graduate

430 scholarship. This research was financed by the Research Council of Norway, contract 216416/O10

431 and by the Norwegian Environmental Agency. Cooke was supported by the Canada Research

432 Chairs Program and NSERC. Thanks to Colin Davis for helping with simulation coding.

433

434 6.0 References

435

436 Arlinghaus, R. and Cooke, S.J. 2008. Recreational fishing: socio-economic importance,
437 conservation issues and management challenges. *In* Recreational Hunting, Conservation
438 and Rural Livelihoods: Science and Practice. *Edited by* B. Dickson, J. Hutton, and B.
439 Adams. Blackwell Publishing, Oxford, pp 39-58.

440 Arlinghaus, R., Cooke, S.J., Lyman, J., Policansky, D., Schwab, A., Suski, C., Sutton, S.G., and
441 Thorstad, E.B. 2007. Understanding the complexity of catch-and-release in recreational
442 fishing: an integrative synthesis of global knowledge from historical, ethical, social, and
443 biological perspectives. *Rev. Fish. Sci.* **15**:75-167.

444 Baisez, A., Bach, J.M., Leon, C., Parouty, T., Terrade, T., Hoffmann, M., and Laffaille, P. 2011.
445 Migration delays and mortality of adult Atlantic salmon *Salmo salar* en route to
446 spawning grounds on the River Allier, France. *Endanger. Spec. Res.* **15**:265-270.

447 Barton, B.A. 2002. Stress in fishes: a diversity of responses with particular reference to changes
448 in circulating corticosteroids. *Integr. Comp. Biol.* **42**:517-525.

449 Booth, R.K., Kieffer, J.D., Tufts, B.L., Davidson, K., and Bielak, A.T. 1995. Effects of late-
450 season catch-and-release angling on anaerobic metabolism, acid-base status, survival, and
451 gamete viability in wild Atlantic salmon (*Salmo salar*). *Can. J. Fish. Aquat. Sci.* **52**:283-
452 290.

453 Coggins, L.G., Catalano, M.J., Allen, M.S., Pine, W.E., and Walters, C.J. 2007. Effects of
454 cryptic mortality and the hidden costs of using length limits in fishery management. *Fish.*
455 *Fisheries* **8**:196-210.

456 Cooke, S.J. and Schramm, H.L. 2007. Catch-and-release science and its application to
457 conservation and management of recreational fisheries. *Fish. Manage. Ecol.* **14**:73-79.

458 Cooke, S.J., Messmer, V., Tobin, A.J., Pratchett, M.S., and Clark, T.D. 2014. Refuge-Seeking
459 Impairments Mirror Metabolic Recovery Following Fisheries-Related Stressors in the
460 Spanish Flag Snapper (*Lutjanus carponotatus*) on the Great Barrier Reef. *Physiol.*
461 *Biochem. Zool.* **87**(1):136-147.

462 Cooke, S.J., Schreer, J.F., Dunmall, K.M., and Philipp, D.P. 2002. Strategies for quantifying sub-
463 lethal effects of marine catch-and-release angling: insights from novel freshwater
464 applications. *Am. Fish. Soc. Symp.* **30**:121-134.

465 Cooke, S.J., Suski, C.D., Ostrand, K.G., Wahl, D.H., and Philipp, D.P. 2007. Physiological and
466 Behavioral Consequences of Long-Term Artificial Selection for Vulnerability to
467 Recreational Angling in a Teleost Fish. *Physiol. Biochem. Zool.* **80**:480-490.

468 Cox, S.P. and Walters, C. 2002. Modeling exploitation in recreational fisheries and implications
469 for effort management on British Columbia rainbow trout lakes. *N. Am. J. Fish. Manage.*
470 **22**:21-34.

471 Croze, O. 2008. The impact of the channeled part of the Auline River (France) on the upstream
472 migration of returning adult Atlantic salmon as determined by radio-tracking. *Am. Fish.*
473 *Soc. Symp.* **65**:23-37.

474 Davidson, K., Hayward, J., Hambrook, M., Bielak, A.T., and Sheasgreen, J. 1994. The effects of
475 late-season angling on gamete viability and early fry survival in Atlantic salmon.
476 Canadian Technical Report of Fisheries and Aquatic Sciences **1982**:1-12.

477 de Gaudemar, B.D. and Beall, E. 1998. Effects of overripening on spawning behaviour and
478 reproductive success of Atlantic salmon females spawning in a controlled flow
479 channel. *J. Fish Biol.* **53**:434-446.

480 Dempson, J.B., Furey, G. and Bloom, M. 2002. Effects of catch and release angling on Atlantic
481 salmon, *Salmo salar* L., of the Conne River, Newfoundland. *Fish. Manage. Ecol.* **9**:139-
482 147.

483 Dingle, H. 1980. Ecology and evolution of migration. *In* Animal migration, orientation, and
484 navigation. *Edited by* A. Cauthreaux. Academic Press, New York, pp. 1-101.

485 Donaldson, M.R., Arlinghaus, R., Hanson, K.C., and Cooke, S.J. 2008. Enhancing catch-and-
486 release science with biotelemetry. *Fish. Fisheries* **9**:79-105.

487 Downton, P.R., Reddin, D. G., and Johnson, R. W. 2001. Status of Atlantic salmon (*Salmo salar*
488 L.) in Campbellton River, Notre Dame Bay (SFA 4), Newfoundland in 2000. Department
489 of Fisheries and Oceans Canadian Science Advisory Secretariat Research Document
490 2001/031. 73 pp.

491 Einum, S., Nislow, K.H. 2005. Local-scale density-dependent survival of mobile organisms in
492 continuous habitats: an experimental test using Atlantic salmon. *Oecologia* **143**:203-210.

493 Fleming, I.A. 1996. Reproductive strategies of Atlantic salmon: ecology and evolution. *Rev. Fish*
494 *Biol. Fisheries* **6**:379-416.

495 Garant, D., Dodson, J.J., and Bernatchez, L. 2000. Ecological determinants and temporal
496 stability of the within-river population structure in Atlantic salmon (*Salmo salar* L.). *Mol.*
497 *Ecol.* **9**:615-628.

498 Gargan, P.G., Stafford, T., Økland, F., and Thorstad, E.B. 2015. Survival of wild Atlantic salmon
499 (*Salmo salar*) after catch and release angling in three Irish rivers. *Fish. Res.* **161**:252-260.

500 Gudjonsson, S., Antonsson, T., and T. Tomasson. 1996. Exploitation ratio of Atlantic salmon in
501 relation to Atlantic Salmon run in three Icelandic rivers. International Council for the
502 Exploration of the Sea Statutory Meeting, ANACAT Committee, M:8.

503 Hanson, K.C., Cooke, S.J., Suski, C.D., Niezgodá, G., Phelan, F.J.S., Tinline, R., and Philipp,
504 D.P. 2007. Assessment of largemouth bass (*Micropterus salmoides*) behavior and activity
505 at multiple spatial and temporal scales utilizing a 3-D whole-lake ecological telemetry
506 observatory. *Hydrobiologia* **582**:243–256.

507 Havn, T.B., Uglem, I., Solem, Ø., Cooke, S.J., Whoriskey, F.G., and Thorstad, E.B. 2015. The
508 effect of catch-and-release angling at high water temperatures on behavior and survival of
509 Atlantic salmon during spawning migration. *J. Fish Biol.* **87**:342-359.

510 Heggberget, T.G., Hansen, L.P., and Næsje, T.F. 1988. Within-river spawning migration of
511 Atlantic salmon (*Salmo salar*). *Can. J. Fish. Aquat. Sci.* **45**:1691-1698.

512 Heggberget, T.G., Lund, R.A., Ryman, N., and Ståhl, G. 1986. Growth and genetic variation of
513 Atlantic salmon (*Salmo salar*) from different sections of the River Alta, North
514 Norway. *Can. J. Fish. Aquat. Sci.* **43**:1828-1835.

515 Hendry, A.P. and Beall, E. 2004. Energy use in spawning Atlantic salmon. *Ecol. Freshw. Fish.*
516 **13**:185-196.

517 ICES. 2013. Report of the Working Group on North Atlantic Salmon, Copenhagen, Denmark,
518 pp. 1-380.

519 Jordan, W.C., Youngson, A.F., Day, D.W., and Ferguson, A. 1992. Genetic protein variation in
520 natural populations of Atlantic salmon (*Salmo salar*) in Scotland: temporal and spatial
521 variation. *Can. J. Fish. Aquat. Sci.* **49**:1863-1872.

522 Kieffer, J.D. 2000. Limits to exhaustive exercise in fish. *Comp. Biochem. Physiol. A* **126**:161-
523 179.

524 Kristinsson, K.O., Gudbergsson, G., and Gislason, G.M. 2015. Variable migration and delay in
525 two stock components of an Atlantic salmon population. *Env. Biol. Fish.* **98**:1513-1523.

526 L'Abée-Lund, J.H. and Aspås, H. 1999. Threshold values of river discharge and temperature for
527 anglers' catch of Atlantic salmon *Salmo salar* L. *Fish. Manage. Ecol.* **6**:323-333.

528 Lee, C.G., Farrell, A.P., Lotto, A., Hinch, S.G., and Healey, M.C. 2003. Excess post-exercise
529 oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*)
530 salmon following critical speed swimming. *J. Exp. Biol.* **206**:3253-3260.

531 Lennox, R.J., Diserud, O.H., Cooke, S.J., Thorstad, E.B., Whoriskey, F.G., Solem, Ø., Havn,
532 T.B., and Uglem, I. In Press. Influence of gear switching on recapture of Atlantic salmon
533 (*Salmo salar*) in catch-and-release fisheries. *Ecol. Freshw. Fish.* DOI: 10.1111/eff.12223

534 Lennox, R.J., Uglem, I., Cooke, S.J., Næsje, T.F., Whoriskey, F.G., Havn, T.B., Ulvan, E.,
535 Solem, Ø., and Thorstad, E.B. 2015. Does Catch-and-Release Angling Alter the Behavior
536 and Fate of Adult Atlantic Salmon During Upriver Migration? *Trans. Am. Fish. Soc.*
537 **144**:400-409.

538 Lucas, M.C. 1994. Heart rate as an indicator of metabolic rate and activity in adult Atlantic salmon,
539 *Salmo salar*. *J. Fish Biol.* **44**:889-903.

540 Mäkinen, T.S., Niemelä, E., Moen, K., Lindström, R. 2000. Behaviour of gill-net and rod-
541 captured Atlantic salmon (*Salmo salar* L.) during upstream migration and following radio
542 tagging. *Fish. Res.* **45**:117-127.

543 Niemelä, E., Orell, P., Erkinaro, J., Dempson, J.B., Brørs, S., Svenning, M.A., Hassinen, E.
544 2006. Previously spawned Atlantic salmon ascend a large subarctic river earlier than their
545 maiden counterparts. *J. Fish Biol.* **69**:1151-1163.

546 Økland, F., Erkinaro, J., Moen, K., Niemelä, E., Fiske, P., McKinley, R.S., and Thorstad, E.B.
547 2001. Return migration of Atlantic salmon in the River Tana: phases of migratory
548 behaviour. *J. Fish Biol.* **59**:862-874.

549 Orell, P. and Erkinaro, J. 2007. Snorkelling as a method for assessing spawning stock of Atlantic
550 salmon, *Salmo salar*. *Fish. Manage. Ecol.* **14**:199-208.

551 Ovidio, M., Philippart, J.C., Baras, E. 2000. Methodological bias in home range and mobility
552 estimates when locating radio-tagged trout, *Salmo trutta*, at different time intervals.
553 *Aquat. Liv. Res.* **13**:449–454

554 Pankhurst, N.W. 2011. The endocrinology of stress in fish: an environmental perspective. *Gen.*
555 *Comp. Endocr.* **170**:265-275.

556 Parrish, D.L., Behnke, R.J., Gephard, S.R., McCormick, S.D., and Reeves, G.H. 1998. Why
557 aren't there more Atlantic salmon (*Salmo salar*)? *Can. J. Fish. Aquat. Sci.* **55**:281-287.

558 Persson, P., Sundell, K., Björnsson, B.T., and Lundqvist, H. 1998. Calcium metabolism and
559 osmoregulation during sexual maturation of river running Atlantic salmon. *J. Fish Biol.*
560 **52**:334-349.

561 Pollock, K.H. and Pine, W.E. 2007. The design and analysis of field studies to estimate catch-
562 and-release mortality. *Fish. Manage. Ecol.* **14**:123-130.

563 R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for
564 Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL [http://www.R-](http://www.R-project.org)
565 [project.org](http://www.R-project.org).

566 Raby, G.D., Donaldson, M.R., Hinch, S.G., Patterson, D.A., Lotto, A.G., Robichaud, D.,
567 English, K.K., Willmore, W.G., Farrell, A.P., Davis, M.W., and Cooke, S.J. 2012.
568 Validation of reflex indicators for measuring vitality and predicting the delayed mortality
569 of wild coho salmon bycatch released from fishing gears. *J. Appl. Ecol.* **49**:90-98.

570 Richard, A., Dionne, M., Wang, J., and Bernatchez, L. 2013. Does catch and release affect the
571 mating system and individual reproductive success of wild Atlantic salmon (*Salmo salar*
572 L.)? *Mol. Ecol.* **22**:187-200.

573 Scarabello, M., Heigenhauser, G.J.F., and Wood, C.M. 1991. The oxygen debt hypothesis in
574 juvenile rainbow trout after exhaustive exercise. *Resp. Physiol.* **84**:245-259.

575 Taggart, J.B., McLaren, S.M., Hay, D.W., Webb, J.H., and Youngson, A.F. 2001. Spawning
576 success in Atlantic salmon (*Salmo salar* L.): A long-term DNA profiling-based study
577 conducted in a natural stream. *Mol. Ecol.* **10**:1047-1060.

578 Thorstad, E.B., Forseth, T., Aasestad, I., Økland, F., and Johnsen, B.O. 2005. In situ avoidance
579 response of adult Atlantic salmon to waste from the wood pulp industry. *Water Air Soil*
580 *Poll.* **165**:187-194.

581 Thorstad, E.B., Næsje, T.F., Fiske, P., and Finstad, B. 2003. Effects of hook and release on
582 Atlantic salmon in the River Alta, northern Norway. *Fish. Res.* **60**:293-307.

583 Tsuboi, J.I. and Morita, K. 2004. Selectivity effects on wild white-spotted charr (*Salvelinus*
584 *leucomaenis*) during a catch and release fishery. *Fish. Res.* **69**:229-238.

585 Tufts, B.L., Davidson, K., and Bielak, A.T. 2000. Biological implications of "catch-and-release"
586 angling of Atlantic salmon. *In Managing Wild Atlantic Salmon – New Challenges, New*
587 *Techniques. Edited by F.G. Whoriskey and K.B. Whelan. Atlantic Salmon Federation, St.*
588 *Andrews, New Brunswick, pp. 100–138*

589 Vähä, J.P., Erkinaro, J., Niemelä, E., Primmer, C.R., Saloniemi, I., Johansen, M., Svenning, M.,
590 and Brørs, S. 2011. Temporally stable population-specific differences in run timing of
591 one-sea-winter Atlantic salmon returning to a large river system. *Evol. Appl.* **4**:39-53.

592 Verspoor, E., Fraser, N.H.C., and Youngson, A.F. 1991. Protein polymorphism in Atlantic
593 salmon within a Scottish river: evidence for selection and estimates of gene flow between
594 tributaries. *Aquaculture* **98**:217-230.

595 Whoriskey, F.G., Prusov, S., and Crabbe, S. 2000. Evaluation of the effects of catch-and-release
596 angling on the Atlantic salmon (*Salmo salar*) of the Ponoï River, Kola Peninsula, Russian
597 Federation. *Ecol. Freshw. Fish* **9**:118-125.

598 Wickham, H. 2009. *ggplot2: Elegant graphics for data analysis*. Springer, New York.

599 Wilson, S.M., Raby, G.D., Burnett, N.J., Hinch, S.G., and Cooke, S.J. 2014. Looking beyond the
600 mortality of bycatch: sublethal effects of incidental capture on marine animals. *Biol.*
601 *Cons.* **171**:61-72.

602 Wood, C.M. 1991. Acid-base and ion balance, metabolism, and their interactions, after
603 exhaustive exercise in fish. *J. Exp. Biol.* **160**:285-308.

604 Wood, C.M., Turner, J.D., and Graham, M.S. 1983 Why do fish die after severe exercise? *J. Fish*
605 *Biol.* **22**:189-201.

606
607
608
609

610 Tables

611 Table 1. Individual data on the radio tagged salmon in the Lakselva River, Norway. Thirty-nine
 612 salmon were captured between July 13 and August 28 2014, eight of which were recaptured later
 613 in the migration, one of which disappeared, and one of which died. One of the recaptured salmon
 614 was re-released and remained in the river for spawning. The spawning pool was determined by
 615 radio tracking in the fall during the spawning season and the net movement is the number of pools.


Capture Date	Total Length (cm)	Fate	Release Pool	Spawning Pool	Net Movement (number of pools)
July 13	73	Survived to Spawn	7	18	11
July 14	97	Recaptured			
July 15	98	Recaptured			
July 16	91	Recaptured			
July 16	90	Survived to Spawn	21	27	6
July 17	95	Survived to Spawn	1	1	0
July 17	80	Recaptured			
July 19	95	Survived to Spawn	17	27	10
July 19	62	Disappeared			
July 24	66	Survived to Spawn	1	1	0
July 26	63	Survived to Spawn	8	10	2
July 27	121	Survived to Spawn	2	2	0
July 30	111	Survived to Spawn	18	24	6

July 30	103	Recaptured	18	34	16
July 30	81	Survived to Spawn	18	14	-4
July 31	102	Survived to Spawn	2	1	-1
August 1	111	Survived to Spawn	18	18	0
August 2	109	Survived to Spawn	18	18	0
August 2	93	Survived to Spawn	18	14	-4
August 2	112	Survived to Spawn	18	14	-4
August 5	112	Survived to Spawn	21	18	-3
August 9	67	Recaptured			
August 9	90	Died			
August 10	64	Survived to Spawn	1	3	2
August 10	94	Survived to Spawn	1	2	1
August 12	94	Survived to Spawn	2	3	1
August 13	99	Survived to Spawn	2	10	8
August 13	69	Survived to Spawn	14	20	6
August 14	69	Survived to Spawn	1	2	1
August 14	84	Recaptured	1	13	12
August 14	91	Survived to Spawn	27	21	-6
August 15	76	Recaptured			
August 15	89	Survived to Spawn	27	27	0
August 16	101	Survived to Spawn	21	24	3
August 17	102	Recaptured			
August 17	112	Survived to Spawn	2	2	0

August 20	83	Survived to Spawn	21	27	6
August 24	77	Survived to Spawn	21	21	0
August 28	66	Survived to Spawn	1	2	1

616


617



620 Figure 1. Lakselva watershed in Porsanger, Finnmark, Norway. The watershed incorporates two
621 major lakes, Øvrevatnet and Nedrevatnet. Atlantic salmon return to the river from the ocean
622 through the Porsangerfjord throughout the summer and migrate upriver to spawning grounds. For
623 this study, salmon released by anglers were tagged at various points in the river although mostly
624 in the lower reaches. Some pool numbers are provided for reference. Note that the river flows
625 south to north.

626

627




628

629 Figure 2. Relationship between the release location and spawning position of 30 salmon released
630 by anglers. Discrete spawning pools are assigned based on locations where spawning counts
631 occurred in September 2015. The dashed line indicates a 1:1 relationship between release pool
632 and spawning pool (i.e. no upriver movement). Points are jittered to reduce overlap. $R^2 = 0.74$.

633

634


A



635

636

B



637


638

639

640

641


C



642

643 Figure 3. Simulated test statistic distributions for the mean number of pools moved for Atlantic
644 salmon under alternative null hypothesis of no effect of catch-and-release on post release
645 movements and final choice of spawning pool. The black diamonds indicate the observed mean
646 number of pools moved from the release location to the spawning location among the tagged
647 salmon. Panel A gives the simulated distribution for the free distribution of salmon, B shows the
648 distribution for the upriver movement only simulation, and C the distribution for the upriver
649 movement simulation that excludes all salmon that spawned at or below the release site. Observed
650 movement (black diamond) in Panels A and B are based on 30 salmon whereas panel C includes
651 15 salmon after removing individuals that spawned downriver of the release site (see Table 1 for
652 list of salmon with negative movement that were excluded).

653



654

655 Figure 4. Violin plots of annual drift counts in Lakselva. The width of violins indicate the spawning
656 densities at corresponding pools of the river based on observations by drift counters. Note that
657 across years there was some inconsistency in visibility, excluding some pools from the count; for
658 example, the lower pools in the 2011 count. Only data from the 2014 count were used for the
659 simulation models.