


2057

NINA Report

Environmental impacts of floating bridges

Arne Follestad, Johanna Järnegren, Evert Johannes Mul, Carolyn Rosten & Frode Thomassen Singaas



NINA Publications

NINA Report (NINA Rapport)

This is NINA's ordinary form of reporting completed research, monitoring or review work to clients. In addition, the series will include much of the institute's other reporting, for example from seminars and conferences, results of internal research and review work and literature studies, etc. NINA

NINA Special Report (NINA Temahefte)

Special reports are produced as required and the series ranges widely: from systematic identification keys to information on important problem areas in society. Usually given a popular scientific form with weight on illustrations.

NINA Factsheet (NINA Fakta)

Factsheets have as their goal to make NINA's research results quickly and easily accessible to the general public. Fact sheets give a short presentation of some of our most important research themes.

Other publishing.

In addition to reporting in NINA's own series, the institute's employees publish a large proportion of their research results in international scientific journals and in popular academic books and journals.

Environmental impacts of floating bridges

Arne Follestad
Johanna Järnegren
Evert Johannes Mul
Carolyn Rosten
Frode Thomassen Singsaas

Follestad, A., Järnegren, J., Mul, E.J., Rosten, C.M. & Singasaas, F.T. 2022. Environmental impacts of floating bridges. NINA Report 2057. Norwegian Institute for Nature Research

Trondheim, January 2022

ISSN: 1504-3312

ISBN: 978-82-426-4840-2

COPYRIGHT

© Norwegian Institute for Nature Research

The publication may be freely cited where the source is acknowledged

AVAILABILITY

Open

PUBLICATION TYPE

Digital document (pdf)

QUALITY CONTROLLED BY

Jørn Thomassen

SIGNATURE OF RESPONSIBLE PERSON

Research director Svein-Håkon Lorentsen (sign.)

CLIENT(S)/SUBSCRIBER(S)

Statens Vegvesen, Utbygging

CLIENTS/SUBSCRIBER CONTACT PERSON(S)

Arianna Minoretti

COVER PICTURE

Submerged bridge © Statens Vegvesen

KEY WORDS

Ferry-free E39, floating bridge, submerged floating tube bridge, sustainable transport, environmental impact assessment, birds, marine mammals, fish, marine invertebrates, sediments, anthropogenic disturbance, literature review.

NØKKELOD

Fergefri E39, flytende bro, nedsenkbare tunneller, bærekraftig transport, miljøkonsekvensutredning, fugler, sjøpattedyr, fisk, marine evertebrater, sedimenter, menneskeskapte forstyrrelser, litteraturstudium.

CONTACT DETAILS

NINA head office

P.O.Box 5685 Torgarden
NO-7485 Trondheim
Norway
P: +47 73 80 14 00

NINA Oslo

Sognsveien 68
0855 Oslo
Norway
P: +47 73 80 14 00

NINA Tromsø

P.O.Box 6606 Langnes
NO-9296 Tromsø
Norway
P: +47 77 75 04 00

NINA Lillehammer

Vormstuguvegen 40
NO-2624 Lillehammer
Norway
P: +47 73 80 14 00

NINA Bergen:

Thormøhlens gate 55
NO-5006 Bergen.
Norway
P: +47 73 80 14 00

www.nina.no

Abstract

Follestad, A., Järnegren, J., Mul, E.J., Rosten, C. & Singasaas, F.T. 2022. Environmental impacts of floating bridges. NINA Report 2057. Norwegian Institute for Nature Research.

The Norwegian government has an ambition of better connecting the west coast of Norway between Kristiansand and Trondheim. Improvement of the fjord crossings will be the primary means of improving connections and reducing travel time. This project has been termed the Ferry Free E39 project and is the responsibility of the Norwegian Public Roads Administration (NPRA). Many of the fjord crossings on the west coast present difficulties for existing bridge and tunnel technologies due to the spans that need to be crossed, or the depths of the fjords. The NPRA is therefore scoping new solutions in the form of floating bridges and submerged, floating tube bridges (SFTB). Floating bridges and SFTB use new approaches such as suspension of bridges, or submerged tunnels, from pontoons. This enables deeper (over 400 m), or wider (over 2 km) stretches to be crossed.

When evaluating new technological solutions, it is important to consider their potential environmental footprints. For this reason, the NPRA has commissioned the Norwegian Institute for Nature Research (NINA) to evaluate the potential environmental effects of floating bridges and SFTB through literature study and expert evaluation. We applied a semi-systematic literature review approach and identified 193 potentially relevant publications. Since floating bridge technology is new and has, as of yet, rarely been applied in practice, little literature that directly assesses environmental effects of floating bridges is available. We addressed this by including literature from other anthropogenic structures in marine environments (e.g., offshore wind farms) which have met some of the same challenges during their construction or operation.

The literature review identified four main classes of environmental impact: habitat alterations, noise pollution, light pollution, and gateways to new ecosystems (e.g. islands) for predators. These impacts were discussed in depth with consideration for both aquatic (benthic invertebrates, zooplankton, fishes and marine mammals) and terrestrial (birds, mammals) organisms and communities. Since fjord crossings already exist in the form of ferries at all locations being considered for floating bridges, potential environmental impacts were compared to ferries, and not a null-crossing situation.

We then present our expert evaluation based upon findings from the literature review and the authors expertise. Two phases were considered during the expert evaluation, the construction phase and the operation phase, since these present different environment challenges. We identified the following groups of effects: (i) habitat changes in the form of reduction in habitat quality through avoidance or acoustic masking, and new habitat availability through additional physical structures for benthic organism growth or removal of barriers to island habitats for predators and other mammals; (ii) physical injury, specifically from noise produced during the construction phase; and (iii) barrier effects in the form of either physical barriers (e.g. SFTBs) or barriers to migration caused by noise, light, or hydrodynamics. Finally, we present mitigation possibilities to reduce the negative environmental impacts of floating bridges and SFTBs, and highlight some key future research needs in this novel and largely unexplored field.

Arne Follestad, NINA, P.O.box 5685 Torgarden, 7485 Trondheim, arne.follestad@nina.no.
Johanna Järnegren, NINA, P.O.box 5685 Torgarden, 7485 Trondheim, johanna.jarnegren@nina.no
Evert Johannes Mul, NINA, Framsenderet, P.O.box 6606 Langnes, 9296 Tromsø, evert.mul@nina.no
Carolyn Rosten, NINA, P.O.box 5685 Torgarden, 7485 Trondheim, carolyn.rosten@nina.no
Frode Thomassen Singasaas, NINA, P.O.box 5685 Torgarden, 7485 Trondheim, frode.singasaas@nina.no

Sammendrag

Follestad, A., Järnegren, J., Mul, E.J., Rosten, C. & Singsaas, F.T. 2022. Miljøeffekter av flytende bruer. NINA Rapport 2057. Norsk institutt for naturforskning

Norske myndigheter har en ambisjon om en bedre veiforbindelse langs kysten mellom Kristiansand og Trondheim. Nye og bedre måter å krysse fjordene på vil være det beste virkemidlet for å redusere reisetiden. Prosjektet har blitt kalt ferjefri E39, der Statens vegvesen (SVV) er ansvarlig for prosjektet. Mange av fjordene på Vestlandet er for brede eller for dype til å bygge tradisjonelle broer eller tunneler. Prosjektet undersøker derfor konsekvensene av alternative løsninger i form av flytende broer. De kan ha forskjellig utforming, fra nedsenkbare tuneller som kan festes til bunn eller henge i pongtonger på overflaten, til flytebroer montert på pongtonger. Flytende broer kan være et alternativ for å krysse dype fjorder, der største dyp er fra 400 meter eller mer, og der fjorden er over to kilometer bred.

Ved vurdering av nye teknologiske løsninger er det viktig å ta hensyn til mulige miljøeffekter. Statens Vegvesen ga derfor NINA oppdraget med å vurdere mulige miljøeffekter av slike broløsninger gjennom et litteraturstudium og en ekspertvurdering. Vi utførte en semi-systematisk litteraturstudie og identifiserte 193 potensielt relevante publikasjoner. Siden teknologien bak flytende broer er ny, og foreløpig i liten grad tatt i bruk, finnes det lite litteratur som direkte tar for seg miljøeffekter av flytende broer. Vi løste problemet ved å inkludere litteratur fra liknende menneskeskaptede strukturer i marine miljøer, som offshore vindmøller, som har hatt flere av de samme utfordringene knyttet til konstruksjon og drift.

Litteraturstudien indentifiserte fire hovedtyper av miljøpåvirkning: habitatendring, lydforurensing, lysforurensing, og inngangsporter til nye økosystemer (for eksempel øyer) for predatorer. Vi diskuterte disse påvirkningsfaktorene med hensyn til både vannlevende (bunndyr, dyreplankton, fisk og sjøpattedyr) og landlevende (fugler og pattedyr) arter og samfunn. Siden ferger allerede krysser fjorden på alle lokalitetene hvor flytende broer vurderes, har vi sammenliknet mulige miljøeffekter med ferger, og ikke uberørte områder uten trafikk over fjorden.

Vi presenterer vår ekspertvurdering basert på funn i litteraturstudien og artikkelforfatternes ekspertise. Vurderingen ble gjennomført i to faser, konstruksjons- og driftsfase, siden disse medfører ulike miljømessige utfordringer. Vi delte miljøeffektene inn i følgende grupper: (i) habitatendringer i form av redusert kvalitet på eksisterende habitat, nye fysiske strukturer danner nye habitater eller fjerning av barrierer, (ii) fysiske skader, spesielt fra støy i konstruksjonsfasen og (iii) barriereeffekter i form av enten fysiske barrierer (for eksempel nedsenkbare tuneller) eller lyd- og lysbarrierer. Vi presenterer til slutt mulige avbøtende tiltak som kan redusere negative miljøpåvirkninger av flytende broer og nedsenkbare tuneller, og fremhever framtidige forskningsbehov i dette nye, og i stor grad utforskede fagfeltet.

Arne Follestad, NINA, Postboks 5685 Torgarden, 7485 Trondheim, arne.follestad@nina.no.
Johanna Järnegren, Postboks 5685 Torgarden, 7485 Trondheim, johanna.jarnegren@nina.no
Evert Johannes Mul, Framsenteret, Postboks 6606 Langnes, 9296 Tromsø, evert.mul@nina.no
Carolyn Rosten, Postboks 5685 Torgarden, 7485 Trondheim, carolyn.rosten@nina.no
Frode Thomassen Singsaas, Postboks 5685 Torgarden, 7485 Trondheim, frode.singsaas@nina.no

Contents

Abstract	3
Sammendrag	4
Contents	5
Foreword	6
1 Introduction	7
1.1 Background	7
1.2 Plans for a ferry free E39	7
1.3 Floating bridges	7
1.3.1 Floating oversea bridge	7
1.3.2 Submerged floating tube bridge (SFTB)	8
1.3.3 Suspension bridge	10
1.4 Offshore wind turbines	10
2 Methods	12
3 Results	13
3.1 Existing literature on floating bridges	13
3.2 Potential effects of floating bridges	13
3.2.1 Habitat change	13
3.2.2 Noise	15
3.2.3 Light	19
3.2.4 Gateways	20
3.2.5 Other	21
3.3 Ecological effects of ferries	22
4 Expert evaluation	24
5 Recommendations	28
5.1 Potential mitigation measures that may be built into bridge design	28
5.2 Research areas on the environmental impacts and mitigation measures of floating bridges	29
6 References	31
7 Appendix: Included papers from WoS literature search	41

Foreword

The Norwegian Public Roads Administration (NPRA) commissioned NINA to carry out a literature study and expert evaluation on the potential environmental effects of floating bridges, since they show much promise for providing fixed links particularly in deep and large fjord crossings. This report provides the basis needed to include environmental impact assessment early in the design process and may be used as a decision tool during the design phase.

The assignment for this project includes:

- Mapping of existing knowledge on how floating constructions may impact marine and terrestrial organisms and seabirds with focus on noise, vibrations (movements in the structure), presence of the construction and how it may change the surrounding environment.
- Perform an assessment of existing sources of noise connected to floating bridges related to traffic to make it possible to compare impacts of floating bridges and ferries.
- Obtain experience with other sources of noise, as sonars and seismic, and how these may impact biodiversity in the marine environment.
- Establish a basis for which measures and considerations should be made during the planning phase to facilitate appropriate mitigating measures and solutions to minimize environmental footprint of the structures.

From the literature search, a total of 193 relevant publications were identified as relevant for the present study. This report includes a broad range of references related to the potential effects that are associated with floating bridges. Although these references do not discuss floating bridges, they may be relevant in the evaluation of potential ecological effects.

12.01.2022 Arne Follestad

1 Introduction

1.1 Background

Examples of permanent floating bridges are relatively rare. Although the concept of floating bridges is not new, most designs have been used in non-permanent settings, often in the context of military activities. As a result, the body of scientific literature that specifically addresses ecological effects of floating bridge designs is almost non-existent. Despite the lack of literature, it is evident that floating bridges can alter habitats and affect species (see literature review in Ghanim et al. 2021, Moore et al. 2013). One strategy to evaluate how floating bridges may affect local ecosystems is to aggregate and review studies that addressed other anthropogenic structures in marine environments and use expert evaluation to extrapolate to floating bridges. For example, numerous studies have addressed the ecological effects of windfarms, pile-driving and other construction activities, aquaculture facilities, fisheries shipping, and pipelines (e.g. Lindeboom et al. 2011, Kuşku et al. 2018, Cox et al. 2018, Bray et al. 2016, Nabe-Nielsen et al. 2014, Fliessbach et al. 2019). Many of these ecosystem effects are likely to overlap with those that may be caused by floating bridges, either during the construction phase or during the operational phase. This report therefore consists of an extensive literature review and expert evaluation to identify potential environmental impacts from the construction and use of different floating bridge designs.

1.2 Plans for a ferry free E39

The plan for a ferry free E39 project aims to eliminate all ferries along the coastal highway E39 in Norway, from Stavanger to Trondheim. This route crosses several fjords which are characterized by great widths (up to 5 km) and depths (up to 1 km). Recent development of offshore technology has now made it possible to plan for new, non-conventional engineering solutions for crossing these fjords without ferries. Floating bridges, submerged floating tube bridges (SFTB) with pontoons or vertical tethers, and suspension bridges on tethers (TLP) have now become a possibility for such crossings. An SFTB will, when built, be the first of its kind worldwide (Source: NPRA).

The Norwegian Public Roads Administration (NPRA) has completed several scoping studies comparing different solutions for crossing the fjords along the E39. These solutions all have advantages and disadvantages, also related to environmental impacts. Up until now no specific literature has been available in their planning of bridges.

Better knowledge of the impacts of floating bridges is important to compare the different solutions and their potential impacts on species and ecosystems. Some of these evaluations will be specific to a geographic area, while others are of a more general nature and connected to the type of construction. General features can be used in the planning process of all fjord crossings, also for comparison to other types of infrastructures, like ferries and fixed bridges.

The NPRA wishes to understand how existing and new technology may impact the biological environment in the vicinity of a bridge, both above and below water surface, so that environmental considerations can be included in the planning process. This report outlines environmental perspectives that ought to be considered during the planning of floating, and traditional, bridges, to ensure environmental impacts are minimize and subsequent mitigation costs are reduced.

1.3 Floating bridges

1.3.1 Floating oversea bridge

A floating oversea bridge floats on pontoons. The structure has an open deck, connected with pillars to the pontoons. If the crossing must allow the boats to pass, a part of the deck is



Figure 1. Floating bridge crossing Bjørnafjorden (from Statens Vegvesen).

elevated, with the help of towers and cables (cable stayed bridge, **Figure 1**). Fillings near the shores or along the structure are usually evaluated as alternatives for the tower location or to shorten the length of the crossing.

Floating bridges are typically considered where undersea tunnels, or suspension bridges are not possible due to the distance of the crossing, or depth of the fjord. In these cases, fjord crossings with ferries are already established. It is important to highlight, that when a floating overseas bridge is a part of a solution to permit a physical fjord crossing, the environmental effects of the crossing solution should be evaluated as an alternative, and in comparison, to ferries.

1.3.2 Submerged floating tube bridge (SFTB).

The submerged floating tube bridge consists of submerged tubes floating under the water surface (**Figure 2**). These structures may have different cross-sections (one or several tubes, with connecting elements, of circular/rectangular/elliptical shape) and the bridge may have intermediate elements (pontoons, tethers, pillars), depending on the depth of the water and the length of the bridge. The tethers are the same elements as those used for TLPs.

Along the E39 road, there are some crossings that are very exposed to environmental loads. These crossings can also be too deep and wide to consider a traditional bridge, such as a suspension bridge, that can reach lengths of up to 2 km. In this case, a SFTB would be a competitive solution, because, submerged from the sea-surface, the structure naturally reduces the main loads with the depth.

When the SFTB is hanging in green pontoons (**Figure 3**), they may be attractive nesting places for some seabirds, such as gulls, terns, and oystercatchers, as these man-made islands are safe for mammalian predators.




Figure 2. Submerged floating twin-tube bridge with floating pontoons (above) and tension legs to the bottom (below) (from Statens Vegvesen).




Figure 3. SFTB with green floating pontoons (from Statens Vegvesen).

1.3.3 Suspension bridge

The suspension bridge is a deck suspended on cables that are attached to towers (**Figure 4**). The structure is an overseas bridge, but for lengths more than approximately 2 km, intermediate towers must be provided. These towers can be ground-based, but for deep waters a solution of TLP (tethers) towers is an alternative (**Figure 5**).



Figure 4. Suspension bridge on TLP for Bjørnafjord (from Statens Vegvesen).



Figure 5. Suspension bridge on TLP for Bjørnafjord – underwater view (from Statens Vegvesen).

1.4 Offshore wind turbines

Since so few floating bridges or SFTB have yet been built, or planned, evaluating their environmental footprint is challenging. One way around this problem is to use the experiences from other anthropogenic structures in marine environments. Offshore wind turbines utilise some of the same approaches as floating bridges, so evaluations of potential environmental impacts of these can provide valuable information and are therefore discussed below.

Expected Effects of Offshore Wind Farms

Biological effects resulting from the construction, and operation, of offshore wind farms (OWFs) were identified in a review of studies in Northern European Seas (Bray et al. 2016). Wind farms affect resident and migrating birds, through avoidance behavior, habitat displacement, and collision mortality. Parameters that are heavily weighted on the risks of collision mortality includes flight altitude, flight maneuverability, percentage of flight time, nocturnal flight altitude, disturbance by wind farm structures, ship and helicopter traffic, and habitat specialization. The risk of collision with OWFs is also related to the movement of the upper part of the installation and its dimensions. In the case of 'static', 'not-moving' bridge elements, this event is regarded as rarer.

When planning for OWFs several countries in Northern Europe and in the Mediterranean have identified potential OWF “hotspots”. In the Mediterranean, using lessons learned in Northern Europe (see Bradbury et al. 2014, Bray et al. 2016, Furness et al. 2013, Garthe & Huppopp 2004) they have identified sensitive species and habitats that will likely be influenced by OWFs in both these hotspot areas and at a basin level. This information will be valuable to guide policy governing OWF development and will inform the industry when environmental impact assessments are required for the Mediterranean Sea.

In the breeding season, many seabird species can fly long distances (> 100 km) to find food for their chicks. If birds from colonies far from the planned sites for floating bridges find their food close to these sites, the possible occurrences of birds from colonies or breeding areas far from the site, must also be evaluated.

For fish, population- and community-level effects are poorly known, as available studies are typically short term or relate to individual fish species. Fish are expected to be affected by OWF in both positive and negative ways. A positive effect may occur due to the increased habitat complexity provided by the foundations and any additional scour protection structures. The introduced structures provide increased shelter and colonization substrates for many marine organisms, which in turn may also attract foraging species. This effect is also well known from other anthropogenic structures in the sea, such as oil platforms, piers, wrecks etc. Concerns have also been raised that fish may be repelled from OWF areas because of noise disturbance or disturbance from electromagnetic fields created around cables on the seafloor (references are given in Bergström et al. 2014, Bergström et al. 2013).

Surveillance studies at the Lillgrund wind farm in Sweden on benthic fish communities, revealed no large-scale effects on fish diversity and abundance after establishment of the wind farm when compared to the development in two reference areas. However, changes at smaller spatial scales were evident. Increased densities of all studied piscivores (cod, eel, shorthorn sculpin) were observed close to the foundations in the first years of operation. The increase was probably attributed mainly to local changes in distribution. These results give some indication that OWF might provide long-term benefits by enhancing local ecosystem services (Bergström et al. 2013). It is, however, difficult to say if bridges will have the same positive effect. The offshore windfarm foundations provide hard-substrate habitat for benthic species (etc.), which ultimately leads to increased fish density, but this is in areas where there is perhaps not much hard substrate habitat to begin with. The fjords along the E39 have plenty of hard-substrate habitat, so they will likely not have the same effect in such an environment. This speculation ought to be tested before drawing conclusions.

2 Methods

Traditional, qualitative literature reviews can sometimes represent subjective views, they can lack transparency, or they can be liable to biases. To avoid this, we based our literature search on some of the more systematic techniques known from search building in evidence reviews (Collins et al. 2015).

Some test searches were conducted, and then relevant search terms and synonyms were identified and sorted into three different categories:

- Terms related to floating bridges and other relevant constructions/vessels
- Terms related to impacts
- Terms related to animals and environment

The search terms were then put together using Boolean operators: OR between terms within the same category, and AND between different categories. In some cases the Boolean operator NEAR/* was used for further narrowing of certain terms.

The following search was conducted in Web of Science Core Collection on October 26th 2021:

(TS=(float NEAR/3 bridge*) OR TS=(discrete-pontoon) OR TS=(continuous-pontoon) OR TS=(pontoon-separated) OR TS=(subsea NEAR/3 tunnel*) OR TS=(underwater NEAR/3 tunnel*) OR TS=(submerged NEAR/3 tunnel) OR TS=(submerged NEAR/3 bridge) OR TS=(Suspend* NEAR/3 tunnel) OR TS=(Archimedes NEAR/3 bridge) OR TS=(Float* NEAR/3 structure*) OR TS=("wind turbine" AND (sea* OR offshore OR water)) OR TS=("wind power" AND (sea* OR offshore OR water)) OR TS=("wind farm" AND (sea* OR offshore OR water)) OR TS=((cable* OR cord* OR Rope*) AND (bottom OR seabed OR seafloor OR "ocean floor")) OR TS=(ferry OR ferries))*

AND

(TS=(pollution) OR TS=(nois*) OR TS=(light* AND (pollut* OR disturb*)) OR TS=(vibrat*) OR TS=(barrier*) OR TS=(current* AND (ocean OR sea OR river OR water)) OR TS=(collision*) OR TS=(temperat*) OR TS=(entangle*))*

AND

(TS=(Plant) OR TS=(animal*) OR TS=(seabird*) OR TS=(invertebrat*) OR TS=(whale*) OR TS=(fish*) OR TS=(seal*) OR TS=(mammal*) OR TS=(Bentho*) OR TS=(environment*))*

The search resulted in 1441 publications in Web of Science Core collection. As many of these references were not relevant for this review study, all 1441 references were subsequently evaluated using the reviewing tool "Rayyan" (www.rayyan.ai). The result list was split into four equal parts for screening, one part for each researcher in our team. All references were then screened, based on title and abstract. 1154 references were excluded from the study, while 161 references were deemed relevant by one evaluator in the first evaluation round. In addition, 126 references were initially marked as potentially relevant – from which 32 were eventually included based on the assessment of a second evaluator. The total list of publications that were deemed relevant thus consists of 193 publications.

Based on this list of relevant publications, a number of potential consequences of floating bridges were identified. Most of these consequences have been addressed in relation to other anthropogenic activities, even though these studies might not have been related directly to floating bridges. Whenever relevant, such publications have been used in this report to address the potential consequences of floating bridges in more detail. In addition, "grey literature" was searched for in Google, and some papers referred to in the examined publications were also included in the report.

3 Results

3.1 Existing literature on floating bridges

Few scientific studies (Ghanim et al. 2021, Moore et al. 2013) have directly addressed the impact of floating bridges on the surrounding ecosystem. However, our extensive automated literature search yielded 1441 references, from which 193 were deemed relevant after manual evaluation. Most of these studies did not directly evaluate ecological consequences of floating bridges, but they address elements that overlap with floating bridge designs and discuss associated ecological consequences.

3.2 Potential effects of floating bridges

Based on the assembled literature, the ecological consequences of floating bridges and other anthropogenic activities in the marine environment can be divided into different categories. These can be broadly summarised as habitat changes, noise, light, gateways and other considerations. These categories are addressed in the sections below.

3.2.1 Habitat change

Hydrodynamics

The physical presence of an artificial structure will alter local hydrodynamics around the structure, effecting currents, wave energy and flow. Depending on depth, this may influence sediment distribution, sediment properties and grain size which typically leads to changes in environmental diversity, abundance and species composition (Martin et al. 2005, Moschella et al. 2005, Walker et al. 2008). These changes are most pronounced in the immediate vicinity of the structure. Recent research suggests that a floating bridge may alter hydrodynamic patterns in a fjord (Khangaonkar et al. 2018a). Circulation in a classic fjord is characterised by a shallow brackish layer at the surface over a deep saltwater column that is vulnerable to disruptions. The presence of floating structures could constrict the mixing and transport in the upper layers (<20 m) of the water column (Khangaonkar et al. 2018b). In addition to currents, this may effect also salinity and temperature downstream of the construction (Khangaonkar et al. 2018a).

Artificial reefs

Research has shown that if a structure remains intact long enough to allow ecological succession, the dominating species on floating structures are filter feeders, such as tunicates, sponges, hydrozoans and polychaetes (Connell 2000, Perkol-Finkel et al. 2008, Vandendriessche et al. 2015, Whomersley & Picken 2003). However, artificial structures differ physically from natural habitats with respect to substratum composition, complexity, surface area, age, orientation, movement and disturbance regimes. It is shown to support different ecological communities compared to natural habitats, often characterised by greater abundances of opportunistic and non-native species. Surveys have shown that artificial structures favour non-indigenous species over native hard-bottom species (reviewed by Bishop et al. 2017, Momota & Hoskawa 2021). Opportunistic colonists of hard substrates are often tolerant to a wide range of environmental conditions and have larvae present in the water column for much of the year and are therefore able to rapidly recruit to new substrates. Artificial structures have played a role in establishment and spread of invasive algae (Bulleri & Airoldi 2005) and mussels (Baker et al. 2007, Spinuzzi et al. 2013) but also indigenous species (Rooker et al. 1997) and it may play a role in facilitating climate migrants (reviewed by Bishop et al. 2017). It is important to note that artificial floating structures do not have a natural counterpart, meaning that these structures create novel habitats for both benthic and pelagic organisms (Holloway & Connell 2002, Perkol-Finkel et al. 2006).

The biomass on a submerged artificial structure may reach up to 500-fold the biomass found in the surrounding soft sediment and mainly consists of filter-feeders (Picken et al. 2000). Through this it may act as a bio-filter, affecting the water chemistry and particle composition, and depleting primary organic matter in the close vicinity (<100 m) (Maar et al. 2009).

Floating structures offer a shaded habitat with overhang, something that is rare in natural marine habitats. These are habitats where invasive species often thrive, but here ecological engineering can play an important part by adding elements to lessen shading effects and include 3D complexity offering shelter for juvenile fish etc. (Hadary et al. 2022).

Another effect that has received little attention is the fact that organisms attached to floating structures live at a fixed depth and do not experience tidal cycles. This may effect behavioural patterns of benthic intertidal communities (Hadary et al. 2022). If this has any importance to these systems remains to be seen.

The anchor and connecting system of any artificial structure will destroy habitats on the ocean floor. It is important that this system is designed for minimal contact with the seabed to prevent erosion and destruction of valuable habitat. Concrete gravity based anchor or sinker have greater eco-engineering potential than mooring solutions, as they can be designed to provide habitat or shelter to the affected area (Hadary et al. 2022).

Migration barrier

Migratory species are more likely to encounter and be affected by anthropogenic barriers than non-migrating species. Studies indicate that a floating bridge may function as a barrier to salmonid fish passage. Slower migration times, higher mortality rates in the vicinity of the bridge relative to other areas on the migration route, and unique behaviour and mortality patterns at the bridge suggests they impede migration and increasing predation of salmonids (Celedonia et al. 2008, Moore et al. 2013). During early migration Atlantic salmon (*Salmo salar*) post-smolts usually swim close to the surface (<3 m), but make irregular dives down to 7 m depth (Thorstad et al. 2012). Adult salmon and adult sea trout (*Salmo trutta*) also occur mainly in the upper 5 m in the water column both when returning to rivers and migrating out again after spawning (Davidsen et al. 2013, Thorstad et al. 2016). Although diving intensity increases with time and distance from the targeted estuary, adult Atlantic salmon show large individual variation in how deep they dive, from >300 m during outward migration and around 100 m during inward migration (Kjellman 2015). The few studies of depth use of both Atlantic salmon and sea trout in their marine migration have been conducted during the summer months, but more knowledge is needed during other times of the year (Thorstad et al. 2016). From the perspective of a migrating salmonid, a suspension bridge using TLP or SFTB, affecting the upper water levels (<20 m) would be preferable to having a pontoon-supported structure.

Little is known about the effects of habitat alteration due to bridge or tunnel designs on marine mammals. The distribution of marine mammals in Norwegian fjords is linked to the distribution of prey, which means that any effects of bridges on prey species might indirectly affect the distribution of marine mammals. Direct effects, such as a barrier effect have not been addressed, but the diving behaviour of species that are common along the Norwegian coast could help identify potential conflict-depths. Harbour porpoises (*Phocoena phocoena*) can dive to more than 400 meters, and foraging depths may be related to the water depth (Nielsen et al. 2018, Westgate et al. 1995). Dives that are related to transiting behaviour are typically V-shaped, rather than U-shaped foraging dives, and occur primarily within the top 20 m (Bjørge 2003). Harbour seals (*Phoca vitulina*) can dive to 500 m, but studies performed in different areas showed conflicting results for average diving depths. One study, performed in the St. Lawrence estuary (Canada) indicated that more than half of the recorded dives occurred within the top 4 meters (Lesage et al. 1999), while a study performed in Norway indicated that seals often dive to the bottom to forage (Bjørge et al. 1995). Depending on the design of the bridge, entanglement may be considered a pressure. Although whale entanglements in underwater cables, such as telecommunication cables, have been reported in the past, such events are considered rare (Wood & Carter 2008). However, cables that may be used in the design or anchoring of floating bridges may differ from telecommunication cables. There are currently no methods to deter marine mammals from the vicinity of cables. Deterrent devices that are used on fishing gear or near construction activities are developed for short-term use only.

Connectivity

Artificial structures may increase connectivity of species by acting as steppingstones. Steppingstones across unfavourable stretches of habitat such as extensive soft sediment areas may aid dispersal of organisms or resources that previously only rarely transgressed a barrier. For hard-bottom organisms, the large areas of ocean or even coastal areas that lack or have low densities of hard substrates can be a natural barrier to dispersal. Artificial structures provide new hard substrate, which may be used by some species in addition to their natural substrates and may provide new dispersal pathways by serving as destinations and sources of larvae. Also, coastal structures may provide habitat for species dispersed by shipping and other vectors. Dispersal of species with short pelagic larval durations may particularly benefit from artificial structures serving as 'stepping stones', while species with longer pelagic larval durations may be less affected (Bishop et al. 2017).

3.2.2 Noise

Sound is one of the key pressures associated with anthropogenic activities in the marine environment. Airborne sounds of certain activities can have a relatively large range (Van Renterghem et al. 2014), and may disturb behaviour of seals or seabirds in some situations (Acevedo-Gutierrez & Cendejas-Zarelli 2011). However, ecological consequences of underwater noise can be far more severe. Since water has a much higher density than air, sound travels much faster and further underwater (Jelle et al. 1988). The perceived intensity of sound is also much higher underwater. Many aquatic organisms rely on sound production, transmission and reception for key aspects of their life-histories. Sound is used for orientation, migration, habitat selection, communication, mating behaviour and the detection of prey or predators (Dolman & Jasny 2015, Kuşku et al. 2018). Anthropogenic sound can therefore influence many species throughout the food web, including fish, marine mammals and plankton (Culloch et al. 2016, Kuşku et al. 2018, McCauley et al. 2017). However, this important source of information is under increasing threat due to the increased prevalence of underwater human-produced sounds. A wide range of human activities produce sound in the hearing and sound production ranges of a wide range of organisms (**Figure 6**). However, since floating and submerged bridges are so cutting edge, little to no information exists on their contribution to sound pollution. Conclusions must therefore be drawn from studies of other sound pollution sources and extrapolation to floating and submerged bridges.

Focus of marine sound pollution literature is currently biased towards marine mammals (Erbe 2012), though the studies that do exist also find anthropogenic sound impacts on fishes, invertebrates, marine birds and reptiles (Gentry 2002, Murchy et al. 2019, Popper & Hawkins 2019, Sørensen et al. 2020). Effects of noise on marine animals range from death to physical injury, displacement, and behavioural changes. Physical injury may come in the form of physical damage to the ear or swimbladder (Smith & Monroe 2016), or physiological changes reflecting, for example, stress (Celi et al. 2016). Masking by anthropogenic noise can cause a decrease in detectability of biologically relevant sounds such as those of predators or prey, sounds of conspecifics or acoustic cues used for orientation (Kaplan et al. 2016, Pine et al. 2016). Behavioural responses are varied and can reflect both startle responses and short- or long-term avoidance of sound sources (Popper & Hawkins 2019). While reactions such as startle response may only be transient, avoidance can result in changes to migration routes or abandonment of feeding or breeding grounds (de Jong et al. 2020). It is the population scale effects resulting from all these effects of anthropogenic noise that guide the assessment of severity. If noise impacts on populations in a way that causes population declines, then regulation and mitigation will need to be more restrictive than if the populations can tolerate the impacts.




Figure 6. Sources and animal receivers of sound in the ocean soundscape. Approximate sound production and hearing ranges of marine taxa and frequency ranges of selected anthropogenic sound sources. These ranges represent the acoustic energy over the dominant frequency range of the sound source, and colour shading roughly corresponds to the dominant energy band of each source. Dashed lines represent sonars to depict the multifrequency nature of these sounds. Reproduced from Duarte et al. 2021.

Marine mammals

The majority of studies that address the ecological effects of anthropogenic underwater noise are focused on marine mammals (Cominelli et al. 2018, Gervaise et al. 2012, Koschinski et al. 2003, Nabe-Nielsen et al. 2014), which rely heavily on sound for communication, prey location, orientation and predator avoidance (Kastelein et al. 2013). Based on sound production and perception (hearing), marine mammals can be divided into different groups (see also **Figure 6**). Baleen whales (mysticeti) use low-frequency sound for communication, while toothed whales (Odontoceti) primarily rely on high-frequency sound for finding prey (echolocation). Seals (including all seals, sealions and walrus within the families Phocidae, Otariidae & Odobenidae)

use a wide frequency range that falls between the two whale groups. All three groups are represented in Norwegian waters, and members of each of these groups can be found near the coast. The most common baleen whale in Norwegian waters is the Minke whale (*Balaenoptera acutorostrata*) (Hammond et al. 2017) but several other species are also frequently observed along the Norwegian coast. While baleen whales can be found in Norwegian fjords, this is more common in northern Norway. However, baleen whales communicate using low-frequency sounds, and are therefore sensitive to low-frequency anthropogenic sounds (Terhune & Killorn 2021), which can travel over long distances. This means that potential consequences of activities that produce low-frequency sounds should be considered, even if the activity is located in the fjords. Some marine mammals are often found in the fjords along the coast of southern Norway. Harbor porpoises are abundant along the entire Norwegian coast and are often found deep in fjords. Like other toothed whales (e.g. dolphins, sperm whales and pilot whales), they use high-frequency sound to locate prey (echolocation) (Carstensen et al. 2006). Two species of seals are also common along the Norwegian coast and are often found deep inside fjords: harbor seals and grey seals (*Halichoerus grypus*) (Olsen et al. 2010, Øigård et al. 2012). Their hearing overlaps with that of both baleen whales and toothed whales, and sound plays an important role, for example in the reproduction process (mating calls) and acoustic detection of predators, such as killer whales (Hastie et al. 2015).

Seabirds

Studies on the effects of underwater sounds on seabirds are lacking. One effect that has recently come to light is the masking of the sounds from seabird's prey. Recent studies into the hearing of great cormorants shows that they have a better sense of hearing in water than they do above the surface (Baker 2020, Mooney et al. 2019). Most seabirds have a poor sense of sight underwater, thus cormorants are more likely to hear the fish they are trying to catch, instead of seeing them before they are close to them. Most seabirds will first locate their prey by smell, following chemical trails like dimethyl sulphide produced by plankton to find areas of high productivity (Baker 2020). Thus, predation by seabirds may be impacted by underwater noise.

Bridges as a source of noise

Anthropogenic sound can be classified into two groups; *impulsive sound* most commonly produced during construction phases, and *continuous sound* most commonly produced during operational phases.

Impulsive noise – construction phase

The construction phase of floating bridges is short-term, however, sound production during this phase can be of higher intensity and therefore warrants consideration. Sound types and intensities will vary with construction type. Sound sources from the construction phase are piling, drilling, groundwork and increased road or boat traffic (Thomsen et al. 2006). Other than traffic sounds, these are typically transient, brief (<1s), broadband and show high peak sound pressure with a rapid rise time and rapid decay and are termed impulsive sounds. Piling is the most commonly studied of these. Passage of sound and vibration into the substrate, which can be caused by sources such as pile driving, may result in waves propagating through the substrate, both as compression waves and interface waves (Popper & Hawkins 2019). They may generate evanescent sound pressure and particle motion waves that propagate through the water. This effect can occur both during underwater piling, but also when piling occurs near to the marine environment, as is the case when building landfall arrangements for bridges.

While few studies exist, those existing studies find both considerable effects of impulsive sounds, but also considerable variation between species and settings (Popper & Hawkins 2019). Physical injury, such as tissue damage or temporary hearing loss may occur when organisms pass a given cumulative level of sound energy (Halvorsen et al. 2012a, Halvorsen et al. 2012b, Smith & Monroe 2016). Behavioural changes such as altered feeding behaviour may also ensue (Roberts & Laidre 2019). In many cases however, those organisms that are able to do so will avoid the affected area during construction works (Popper & Hawkins 2019).

Construction of various bridge designs (floating or traditional) might require pile driving, which can have a devastating effect on many marine mammals (Brandt et al. 2018, Lindeboom et al. 2011). Pile-driving leads to a dramatic reduction in sound production by porpoises, which can either indicate a reduction in foraging and socializing behaviour (best case scenario), or it can indicate that porpoises leave the area altogether (Brandt et al. 2011). This effect is noticeable within 20 km from the construction area, but it is possible that porpoises are affected at distances up to 50 km (Dähne et al. 2013). Area avoidance behaviour in response to pile-driving events have also been recorded in seals, and both seals and porpoises within 10 km of the sound source are exposed to sound levels that are sufficient to cause temporary or long-term hearing damage (Hastie et al. 2015). One possible mitigation measure to reduce the exposure of marine mammals to impulsive (temporary) noise, is the deployment of bubble-net curtains during the construction activity (Wursig et al. 2000; Verfuß, 2014). These bubble-nets create an air barrier which reduce the radiation of noise.

Continuous sound - operational phase

Noise from the operation phase is typically long-duration signals that can increase the overall sound level in the environment for extended periods of time. In the cases of floating or SFTB this will likely arise from transfer of vibrations resulting from traffic over or through the bridge, or its natural resonance which varies with wind or current. This will be transferred to the water column through the submerged structure, or in the case of submerged tunnels, through the tunnel itself. Activities onshore, including the passage of vehicles, may increase noise levels in the sea, lakes and rivers, especially if they generate substrate vibration. A recent study by Evergreen Point floating bridge across Lake Washington, Abadi et al. (2018) found that noise levels measured in the lake near to the bridge followed public data on both traffic load and wind speed over the bridge. Likewise, another study showed that passing vehicles in a tunnel (within a riverbed), radiated low-frequency (12-25 Hz) at approximately 14 dB greater levels, compared to the background noise (Song et al. 2020). However, while this demonstrates that sound is transferred from the bridge to the surrounding aquatic environment, the spatial extent of the elevated sound levels requires further research.

Though the sound intensity during the operational phase may often be lower than for construction noise, their continuous character can cause long term impacts, particularly through behavioural changes. Anthropogenic sounds may interfere with foraging behaviour either by masking the relevant sounds or by resembling the sounds that the prey may generate (Purser & Radford 2011). Likewise, avoidance behaviour by prey may depend on listening for the sounds that the predators produce, either deliberately or inadvertently (Luczkovich & Keusenkothen 2008, Ramage-Healey & Bass 2006). If these impacts are biased to the advantage of either the predator or the prey species, they may cause population scale changes. Another potential impact regards movement and migration. Many fishes and marine mammals migrate between feeding or spawning areas. During migrations, they may use a variety of cues to orientate and navigate, including natural soundscapes. If these are altered by noise at key frequencies, or high intensity, ability to navigate may be reduced and/or altered. High level sounds may also cause direct avoidance responses of a given area (Montgomery et al. 2006, Stanley et al. 2012).

Sound is also used directly in communication between individuals of the same species. Communication during spawning is a common example. Any interference with detection of spawning sounds can have a significant effect on reproductive success of a population (de Jong et al. 2020, Popper & Hawkins 2019).

Mitigation measures that were developed to deter marine mammals or other animals from a certain location, such as fishing nets or pile-driving locations, are all only used during short periods. It is therefore important to consider the continuous noise levels during the design phase of the floating bridge, and to adopt a design that is aimed to minimise the radiation of continuous noise. The soundscape that is produced by a floating bridge in operation should be estimated, in order to evaluate potential effects on local fauna.

Anthropogenic noise generated by today's solutions

Current crossing solutions are by means of conventional bridges, tunnels, and ferries. These solutions can generate sound pollution during both construction and operation phases, and it is therefore important to consider these during evaluation of potential new solutions. Current state-of-the-art bridges (e.g. cable-stayed bridges or suspension bridges) require multiple support towers with bottom mounted fundaments, depending on the total required span. Each contact point with the water and seabed provides a point for transmission of soundwaves and vibrations. In addition, noise generated during the construction phase will increase relative to the number of structures being built. Very few studies exist that quantify sound pollution from bridge structures. However, in a recent study of an international bridge crossing between Norway and Sweden, De Clippele & Risch (2021) found significant differences in sound levels on the nearby Tisler cold-water coral reef pre- and post-border closure due to COVID-19. This demonstrates the existence of noise transfer from bridges to the marine environment, at least at busy bridge crossings.

A significant proportion of anthropogenic noise in the ocean is created by motorised vessels, including large ships, ferries, fishing and pleasure boats (Erbe 2013, Erbe et al. 2012). Indeed, during the initial COVID-19 lockdown, a reduction in shipping traffic was reflected in noise levels in the shipping band of frequencies (Thomson & Barclay 2020). Most vessels produce predominately low frequency sound (i.e., <1 kHz) from onboard machinery and hydrodynamic flow around the hull. Cavitation at propeller blade tips is also a significant source of noise across all frequencies (Ross 1993). Low frequency sounds from ships can travel hundreds of kilometres and can increase ambient noise levels over large areas of the ocean (Ellison et al. 2012). Sound production will be to some extent mitigated by new generation boats with electric boat motors and boat design to reduce sound from, for example, machinery (Parsons et al, 2020).

However, sound production from boats will not be entirely eliminated. It ought to be noted that when considering boat traffic from much of the literature in comparison to applications in Norway, many of the studies have been conducted in busy international ports or with the combination of all boat traffic (e.g. shipping, ferries, fishing and pleasure) in focus. Ferries, which are the focus of the current study, make up only a proportion of the boat generated noise. It is difficult to compare the sound-related impact of a (floating) bridge with ferries in the same place, that transport the same number of vehicles. Both solutions will generate different soundscapes, as bridges produce a relatively continuous sound level, while the sound from ferries is more localized and temporary. To our knowledge, no studies have been performed to compare the sound production of (floating) bridges with that of ferries, and their impact on the local ecosystem.

Sound exposure criteria and guidelines

Sound exposure criteria define the levels of sound that are likely to affect aquatic animals negatively, in order to enable regulation of noise generation in aquatic environments. Principal focus has been placed on marine mammals, with little effort placed on developing criteria and guidelines for fishes or invertebrates (Popper & Hawkins 2019). Underwater noise is included in the European Union's Marine Strategy Framework Directive (EU 2008), with the purpose of maintaining good environmental status in the marine environment. The Swedish Environmental Protection Agency published a review solely focusing on pile driving sounds (Andersson et al. 2016b). In this review they propose maximum exposure values based on interim values proposed by the USA (Buehler et al. 2015). These values have a very limited scientific basis and will be adapted in time, but they at least set a precedent for the use of sound exposure criteria and guidelines and an impression of guidelines that will be implemented in the future.

3.2.3 Light

Light pollution is increasing, and artificial light sources may have great impacts on many animals (see literature review in Follestad 2014). For migrating birds, collisions caused by artificial light pollution are a significant source of mortality. Laboratory studies have demonstrated that birds have different visual sensitivities to different colors of light. In a study of the impact of wavelength on phototaxis at two gathering sites of nocturnally migrating birds in Southwest China, short-

wavelength blue light caused the strongest phototactic response (Zhao et al. 2020). In contrast, birds were rarely attracted to long-wavelength red light. The attractive effect of blue light was greatest during nights with fog and headwinds. Zhao et al. (2020) thus suggest that switching to longer wavelength lights is a convenient and economically effective way to reduce bird collisions.

Artificial light sources can, especially under some weather conditions, cause migratory birds to fly toward a light source. Some of these effects are summarized in Follestad (2014). Artificial light sources have increased dramatically in recent decades, with 23 % of the world's land surface between 75°N and 60°S experiencing light-polluted nights (Falchi et al. 2016). This has seriously affected bird migration (La Sorte et al. 2017, Cabrera-Cruz et al. 2018, Horton et al. 2019) and their choices of stopover habitat (McLaren et al. 2018). Migratory birds may aggregate around light sources (Van Doren et al. 2017), and they may thus collide with lighthouses, ships, oil rigs, airports, communication towers, and tall buildings that are illuminated at night (Avery & Cassel 1976, Crawford & Engstrom 2001, Jones & Francis 2003, Gehring 2009, Kerlinger 2010, Merkel & Johansen 2011, Longcore et al. 2008).

A number of factors influence how birds respond to artificial light, including the time of year, weather conditions and properties of the light (Zhao et al. 2014). Poot et al. (2008) found that red light tends to attract migratory birds and suggested that the failure of geomagnetic orientation abilities under red light stimulation may cause collisions with artificial light sources. However, the notion that red light attracts birds has been questioned (Evans 2010). With the current advances in development of new lamps, such as LEDs, and studies on effects of different wavelengths and temperature of the light, an update on this should be made before deciding lightening of components of floating bridges. Design of lights on deck (as illustrated in **Figure 7**) may help reducing the amount of light spread to the surroundings.



Figure 7. The suspension bridge crossing Bjørnarfjord, where lights on deck do not spread light to the surrounding sea surface (from Statens Vegvesen).

3.2.4 Gateways

Bridges and tunnels may introduce mammalian predators to new sites/islands. All types of floating bridges may act as a spreading route for carnivores (such as red fox *Vulpes vulpes*, European badger *Meles meles*, European pine marten *Martes martes*) from the mainland to islands where they are currently not present. This may increase predation on many birds, mammals, amphibians etc. on these islands that are not adapted to such predation. Bergan & Giæver (2002)

describe environmental consequences from Nordøyvegen based on the prerequisite that these predators will NOT spread to the island connected by a number of bridges and tunnels as barriers to stop them, are effective. The American mink *Neogale vison* is already present along most of the Norwegian coast, and there is little reason to believe that floating bridges may expand its distribution.

If barriers built to prevent spread of mammalian predators are not 100 % effective, introduction of new predators may have severe effects on populations of birds, wildlife and other animals (like amphibians). When islands become connected to the mainland by tunnels or bridges, there is a real threat that animals on the mainland, or a nearby island, may spread to new islands. This has been well documented on the island Tautra in the Trondheimsfjord, where red foxes, badgers and pine martens crossed Svaet from the mainland to Tautra by the stone causeway (Thingstad 1994). Thingstad (1994) summarized what happened (sitat): *Tautra, including Svaet, was established as a Ramsar area in July 1985, thus earning international status as an important wetland area for birds. However, after Tautra was connected to the mainland via an approximately 2,5 km long stone causeway across Svaet in 1976 it has become inhabited by predators (particularly foxes and pine martens) that were previously not present there. This has led to heavy predation pressure on the seabird colonies on the island which, among other things, has strongly decimated the nesting populations of eider ducks and common gulls.*

To mitigate the spread of predators, barriers must be planned from the beginning and must be specific for the local needs. A replacement of a stone causeway by a bridge and physical barriers, as a mitigating measure, may have a positive effect of the wildlife.

3.2.5 Other

Effects of road fills

Where floating bridges may include road fills or stone causeways at one or both ends of the link, several effects need to be discussed. In an environmental impact assessment of a planned approximately 1.5 km road link including a 400 m bridge from the island of Kråkvåg to the neighboring island of Storfosna across Kråkvågsvaet, Thingstad & Hokstad (1997) discuss the very local effects of changed currents close to a small road fill in one end of the road. The shallows and rocky/muddy shores at Kråkvågsvaet have achieved international conservation status as a Ramsar area because of their great ornithological qualities as a wetland. Marine biological investigations have shown a demersal fauna rich in species and individuals associated with the shallows and its currents. One alternative for the bridge, if chosen, would cause serious changes to the pattern of currents. This could give a significant reduction in the production of several groups of demersal marine organisms especially important as a source of food for several aquatic bird species.

A road fill can also be a new site for breeding otters or minks, if there is a lack of safe breeding sites nearby. Increase in otter numbers could be considered as positive as it may give locals better opportunities to experience this native species. The mink may, however, be a negative supplement to the local biodiversity, as a ferocious predator of several species. Otters may displace mink close to its dens, with a possible effect of reducing the predation pressure by the mink (Follestad et al. 2005).

Floating or submerged bridges may have an advantage over bridges with towers and wires, which might cause collisions with migrating birds. This may be the case both for seabirds during foraging and during ordinary seasonal migrations.

When bridges cross deep fjords, there will probably be small or no conflicts with feeding areas for seabirds. But in areas near to known, or suspected, important feeding areas, mapping how seabirds utilize the area close to the bridge should be performed in an early stage of the planning phase. This should also include alternative crossing alternatives like "normal" bridges or ferries.

Bridges as nesting place for birds

Birds may use bridges for resting or nesting, and their acid droppings may badly damage concrete. For floating bridges, kittiwakes may start breeding on edges under the bridge, as they have done at the Gjermundnessund bridge. Some years ago, nests were washed away to prevent such damage, but now a foil has been added on the concrete in order to prevent damage and enable the nests stay (A. Follestad own information). Problems with kittiwakes breeding on oil platforms have been discussed by Christensen-Dalsgaard et al. (2020). If kittiwakes and other birds could potentially start breeding on a floating bridge, concern should be taken to minimize problems (see also [Bird Control under Bridges](#)).

3.3 Ecological effects of ferries

A ferry crossing is a less permanent alternative to bridges or tunnels. Ferries do not form a constant ecological pressure, but each crossing can be linked to multiple environmental impacts. Although few studies have addressed the effects of ferries in isolation, many effects related to shipping also apply to ferries. A comprehensive overview of ecological effects related to coastal shipping is presented in Jägerbrand et al. (2019).

Birds

Most studies investigating the effects of ship traffic on seabirds in offshore waters have focused on behavioral responses of scavenging seabird species towards fishing vessels (Votier et al., 2010; Tew Kai et al., 2013; Bodey et al., 2014; Sommerfeld et al., 2016; Le Bot et al., 2018, see references in Burger et al. 2019). With respect to bird disturbance by ships, only few studies exist, mainly due to the difficulty of studying these interactions. Using research vessels as observation platforms, high response distances from approaching vessels were recorded for common scoters (*Melanitta nigra*) and divers (Bellebaum et al., 2006; Schwemmer et al., 2011; Fliessbach et al., 2019). A recent study by Mendel et al. (2019) found the strongest impact of ships on red-throated divers (*Gavia stellata*) for a radius of 5 km and within 5 min of the passage of a ship. However, long-term effects over several hours could not be investigated.

Literature on environmental effects of ferries on seabirds is primarily focussed on high-speed passenger ferries crossing open sea like in the North Sea, Skagerrak, Kattegat or in the Baltic Sea, with partly shallow waters. We have found no studies on disturbance effects on birds from the typical slower Norwegian ferries.

Noise

Vessel noise is perhaps the most prominent pressure associated with ferries, and with shipping in general (Jägerbrand et al. 2019; Cox et al, 2018). Sound levels are related to gross tonnage, where small or medium size vessels (such as ferries) typically produce sound levels up to 175 dB re 1 μ Pa (Kuşku et al. 2018). A car ferry line across the Saguenay Fjord (Canada) added 30-35 dB to ambient noise levels during crossings, which was substantially higher than the noise contribution of smaller whale watching vessels in the same area (Gervaise et al. 2012). Main sources of shipping noise include engines and other machinery, water displacement from the hull and propeller cavitation. The most intensive shipping noise occurs within 10 to 1000 Hz (Celi et al. 2016, Jagerbrand et al. 2019), although smaller vessels may produce intense high-frequency noise as well (see **Figure 6**). Shipping noise may lead to area avoidance behaviour (Fliessbach et al. 2019), masked communication (Terhune & Killorn 2021), elevated stress (Kuşku et al. 2018) and behaviour disturbance (Dehnhard et al. 2019). However, due to the intermittent character of ferry-induced noise, the long-term impacts of sound pollution from ferries may be less severe, compared to continuous sound sources of similar intensity and frequency ranges (Blom et al. 2019).

Other impacts from ferries

Collisions between moving vessels and marine organisms have been reported throughout the world, particularly when large whales are involved. However, collisions with other marine mammals, birds and fish also occur, although it is difficult to evaluate the potential risk of collisions, due to a lack of data (Schoeman et al. 2020). In addition, shipping may contribute to a number

of other environmental pressures, including pollution, either in air or in water (Andersson et al. 2016a, Corbett & Farrell 2002), marine litter (Grøsvik et al. 2018), and light pollution (Jägerbrand et al. 2019).

Electric ferries

The installation of electric engines in ferries may help reduce certain ecological pressures, such as pollution and noise. The radiated noise from a vessel is comprised of various elements, including cavitation noise of the rotating propeller, sound from machinery, which is transported through the vessel hull, and the sound of water displacement by the vessel. The use of electric engines in vessels should not affect cavitation noise or sounds from water displacement, but it is likely to reduce the sound of the engine itself. As a result, electric engines should reduce the overall radiated noise level of vessels. Although reports comparing the ecological impact of conventional ferries and electric ferries are scarce, one publication found a significant reduction in noise, as a conventional ferry was 12 dB louder at distance of 55 m than an electric ferry. Low frequencies (<500 Hz) were specifically reduced, up to 25 dB at short distances from the source (Parsons et al. 2020).

4 Expert evaluation

The literature review has gathered the knowledge basis currently available for assessing potential effects of floating bridges on the environment. In this section the authors process the available information to provide expert evaluation of the potential positive or negative effects of floating bridges. This list is not exhaustive and can be refined and improved as more information becomes available. Potential mitigation measures are suggested in chapter 5 of this report.

It is important to highlight that the effects listed in the sections below are not independent and may result in cumulative effects on animal populations and the environment through combination with other impacts (e.g. light, physical barriers or other noise effects).

Expert evaluation of potential impacts from anthropogenic noise from floating or SFTB bridges

The following potential impacts of noise from construction of floating or SFTB bridges have been identified:

- *Habitat quality reduction as a result of avoidance* - anthropogenic noise may cause marine animals to avoid areas in the vicinity of the construction site. This is relevant for marine invertebrates (zooplankton, bottom fauna), fish and mammals. The extent of the impact will depend on the spatial extent of the area avoided and eventual periodicity. Assuming the construction phase is not long lasting, this will likely not result in population scale effects. However, if construction takes place during important feeding or reproduction periods it may result in population scale effects. Therefore, such periods must be identified prior to the construction, and construction phases that have consistent noise effects should be avoided during these events.
- *Physical injury* – anthropogenic noise may cause physical injury or death to those animals unable to avoid it. This is relevant for marine invertebrates (zooplankton, bottom fauna), fish and mammals. The extent of the impact will depend on the spatial extent of the area in which physical injuries occur and the duration of the impact. If the extent is widespread, then population scale effects may occur. Construction phases that are likely to produce high levels of noise (e.g. pile driving) require mitigation measures that are specifically tailored for the local fauna should be investigated. Potential mitigation measures are suggested in chapter 5 of this report.

The following potential impacts of noise from operation of floating or SFTB bridges have been identified:

- *Migration barrier* – anthropogenic noise may create a barrier to movement through, or into, an area. This is particularly relevant for salmonid migration in and out of freshwater and migrations of marine fish and mammal species between different habitats. The extent of the impact will be dependent on frequency range, intensity and periodicity. Variability in sound intensity or presence with, for example, traffic or weather conditions (wind), will drive the degree of influence.
- *Habitat quality reduction as a result of masking* – anthropogenic noise may mask signals marine animals use for activities such as foraging, predator avoidance and reproduction. This is relevant for marine invertebrates (zooplankton, bottom fauna), fish and mammals. The extent of the impact will depend on sensitivity of each species to the frequencies produced, sound intensity and periodicity (i.e. whether sound pollution is present at high intensities during key feeding or reproduction seasons for each species)
- *Habitat quality reduction as a result of avoidance* – anthropogenic noise may cause marine animals to avoid areas in the vicinity of the bridge structure. This is relevant for marine invertebrates (zooplankton, bottom fauna), fish and mammals. The extent of the impact will depend on the spatial extent of the area avoided and eventual periodicity.

Knowledge about local biodiversity, migratory species and migration routes is essential to optimize the structure design to minimize environmental effects.

Expert evaluation on the comparison of anthropogenic noise from floating or SFTB bridges

It has been documented during the literature review and expert evaluation that all methods of waterway crossing can result in elevated levels of anthropogenic noise in the surrounding marine environment. The effect of noise at a given location is dependent on many things. The geological structure of the surrounding (underwater) area, the species that are present in, or migrate to or through, that area and which ecosystem functions it provides. Therefore, environmental impact assessment is needed on a case-by-case basis to identify problems, needs and solutions prior to the bridge planning phase. There are, however, some general conclusions that can be drawn:

- Floating bridges have less contact structures with the seabed. This is anticipated to reduce the scale of some construction-based noise, such as that from piling. It may also reduce the potential for transfer of noise during the operation phase through contact with the seabed. However, it is currently unknown whether there is any difference (positive or negative) in transfer of sound through fixed contacts with the seabed or floating structures. Further investigations is therefore required to document the full advantages or disadvantages of each approach.
- Boat traffic has been documented in the literature review as being a source of underwater noise. The majority of studies included all boat traffic, not only ferries. Therefore, it is the expert opinion that in most cases, reduction of ferry traffic through replacement with a bridge will not cause major reductions in boat-related underwater noise. Likewise, although ferry traffic will likely increase over time (for example resulting from increased traffic on the better connected E39), it is anticipated that an increase in new, modern ferries will not result in noise increases to significant levels. This must, however, be considered case by case taking into account site-specific conditions and information.

SFTBs are a novel approach which are largely untested. Testing is required to investigate the sound transfer properties of SFTBs to the surrounding aquatic environment. Questions that need to be answered include: To what extent does sound from the SFTB structures transfer to the aquatic environment? Is sound from traffic vibrations transferred to the aquatic environment? Do vibrations or movement of the SFTB in the water column cause sound that is transferred to the aquatic environment? What are the common frequencies and sound intensity of the sounds produced? To what extent can this be mitigated through material choice or design? Studying the existing floating bridges in Norway and existing ferry-connection would make it possible to answer these questions.

Expert evaluation on the comparison of effects on the benthic community from floating or SFTB bridges

The following potential impacts from construction on marine benthic organisms have been identified:

- Destruction of habitat – during construction of bridge types using anchoring, the sea floor will unavoidably be affected by their attachment and existing habitat will be destroyed.
- Turbidity in the water column – installation of anchor systems will result in sediment particles in the water mass. This may affect benthic organisms such as filter feeders and the increased turbidity provides difficulties for visual predators. Depending on length and timing of this phase, it may influence larval production and spreading.
- Migration barrier – depending on length and timing of this phase, the activity may impede both inwards and outwards migration of salmonid fishes, resulting in increased mortality.

The following potential impacts from operation on marine benthic organisms have been identified:

- **Habitat quality reduction** – altered hydrodynamics can change sediment distribution, temperature and salinity which may locally reduce species diversity and change species composition.
- **Providing new habitat** – providing hard substrates as new habitats may have both positive and negative implications. On the positive side it may construct an artificial reef that provide shelter and food for many species, thereby increasing biodiversity and biological biomass. On the negative side the population on the structure differs from natural habitats due to its physical properties, it may act as steppingstones for both invasive and native species and act as a bio-filter, depleting primary organic matter in the near vicinity.
- **Physical barrier** – creating a physical barrier in the ocean will affect the hydrodynamics on this water system on many levels as well as migrating species as well as introduce shading.

Knowledge about site specific biodiversity, oceanography and hydrology is essential to optimize the structure design to minimize environmental effects.

Expert evaluation on the comparison of benthic community changes between floating or SFTB bridges

It has been documented in this review that means of crossing a waterway will have effects on the benthic communities. The degree of the effects will vary depending on a range of factors that has to be assessed for each potential development site. But some general conclusions can be drawn:

- A submerged structure will likely have the most profound effect on the benthic communities. A large, submerged structure will have a significant impact on the hydrodynamics of the area. It will also act as a physical barrier for marine animals on both local and regional scale. It will also provide a prominent area of hard substrate for colonisation of a range of organisms, which, depending on the design, may have both negative and positive effects. If supported by pontoons, it will not have any contact with the sea floor which avoids the destruction of habitat in the construction phase, which is positive, but the pontoons also add to the area of new substrate. If supported by anchors, the benefit of avoiding bottom destruction is lost but the surface area is reduced.
- A suspended bridge is likely the solution that will impact the benthic communities the least. Even if supported by anchors and pontoons, the effect on hydrodynamics is minimized and the introduction of new substrate will be considerably less. It will also have a considerably less effect on migratory species.
- To understand the potential effects a construction may have and to best validate mitigating procedures it is very important to have as much local and site-specific knowledge as possible.

Expert evaluation of potential impacts from light from floating or SFTB bridges

It has been documented in the literature that artificial light may affect several plant and animal species (see literature review in Follestad 2014). The effect of floating bridge lighting at a given location will depend on several factors, such as type of bridge, type of light source (lamps) and how they are designed and placed on the bridge. Lamps along the deck are used to secure a safe crossing by cars, bicycles or people, and in towers to warn ships and airplanes. Therefore, environmental impact assessment is needed on a case-by-case basis to identify problems, needs and solutions prior to the bridge planning phase. Some general conclusions can, however, be drawn:

- Lights may, under certain weather conditions, attract birds to circle around them, increasing the risk for collisions with structures of the bridge, becoming exhausted (if they continue to circle into dawn) or the risk of predation.
- Lamps may spread light to the sea surface if they do not have screens to prevent light spreading to the surroundings. This may affect migrating fishes, such as salmon smolt, or attract plankton or small fish to the surface.
- A long row of lamps may be a barrier to migration in an area where no lights have been present before the new bridge.
- It's important to obtain knowledge of the species of concern to understand the extent of the eventual interference, as animals differ in their sensitivity to different colours.

Expert evaluation of potential impacts from gateways from floating or SFTB bridges

The literature review has documented that where floating bridges replace ferries between islands and mainland, they may act as gateways for mammalian predators. If introduced to pristine areas, these predators can become a catastrophe for birds and other animals already present and not adapted to the presence of such predators. Dependent on which species are introduced in this way, ground breeding birds will be most affected in case of foxes and badgers, and birds breeding in trees in the case of pine martens. The risk for bridges acting as gateways will depend on:

- Effects of gates to prevent predators to get access to decks
- Time of the year when they cross the fjord
- Effects of take-out programs to remove predators

It is important that the necessary gates/barriers are planned during the design phase of the project to ensure their 100 % effective function.

5 Recommendations

A number of mitigating measures can be put in place to reduce negative effects of the construction process or during operations or provide positive solutions for wildlife. A number of these are listed below.

In addition, better understanding of the potential impacts of floating bridges and SFTBs through further research will open the door for new mitigation measures.

5.1 Potential mitigation measures that may be built into bridge design

1. Conduct full environmental impact assessments prior to planning, during construction and during operation phases of all structures. That includes (i) evaluation of the area, important features in geology (structures that may affect sound transfer) and biology (species presence, special ecosystem functions such as feeding or reproduction) that may affect selection and design of crossing solution, (ii) baseline study to characterise environmental features, species communities and ecosystem function and to plan the monitoring program, (iii) monitoring during the construction phase, (iv) monitoring during the operation phase.
2. Assess if the scoping part of the environmental impact assessment should be conducted as a participatory dialogue process which will give ownership to the stakeholder and reduce the level of conflicts.
3. Prevent SFTBs and bridges acting as gateways for predators into pristine areas by ensuring that wildlife exclusion measures are 100 % effective in preventing predators like foxes, badgers and pine martens.
4. Adapt lighting on bridges and pontoons to minimize impact. Use lamps with recommended design, colour spectra and temperature, based on the state-of-the-art knowledge available for species in focus.
5. Establish lighting regimes to turn off light in weather conditions that might increase collision risk of wildlife with different bridge.
6. Adapt pontoons for use by breeding birds. Secure pontoon edges (with a low fence?) so chicks do not fall into the sea (they will not be able to climb onto the pontoons). Build small structures on the pontoons to provide shelter for chicks from avian predators.
7. If birds start breeding on green pontoons, lights to warn ships should not shed light into the pontoon to reduce the risk of predation on the chicks.
8. Encircle piling activities with a bubble curtain or other noise reduction techniques to reduce sound transferred to the marine environment (Wursig et al. 2000; Verfuß, 2014).
9. Create submerged bridges in such a way that vibration from traffic movement (or movement of the structure in water?) is limited. Either by insulation or choice of construction material/method.
10. Given enough site-specific information the sound transfer properties of each potential crossing solution could be assessed for the area. For example, in areas with a lot of wind, SFTBs may be better adapted since over-water structures will move more in the wind and transfer more noise to the aquatic environment. On the other hand, areas with strong currents may cause a SFTB to create more noise.
11. Adapt the resonance frequency of a bridge to avoid key frequencies. This however, requires prior knowledge of which species are present and their key frequency sensitivities.
12. Seed structures with indigenous species, such as vegetation or invertebrates. These can occupy space that could otherwise be rapidly colonised by opportunistic non-native species and help to preserve native biodiversity.

13. Use environmentally friendly materials in structures to improve performance and durability, as well as reduce ecological stress and encourage the development of natural communities.
14. Bioprotection of the structure can also reduce the maintenance costs due to increased lifespan. Reducing the magnitude and frequency of structural maintenance improve ecological stability (reduced anthropogenic intervention), as well as reduce maintenance costs.
15. Incorporate microhabitats into artificial structures to promote diversity of species and provide refugia from both abiotic and biotic stress.
16. Integrate ecological considerations into the design of repair and maintenance of marine infrastructures during the lifetime of the structure.
17. Consider the ecological value of areas prior to planning bridge type and placement. These will be identified during the environmental impact assessment area evaluation phase.
18. Avoid construction on key areas of conservation value. Areas considered essential habitat for threatened or endangered species should be excluded from development plans.

5.2 Research areas on the environmental impacts and mitigation measures of floating bridges

Anthropogenic noise

- More research is needed into noise originating from floating bridge constructions and its impact on key species. For example, documenting the characteristics of sound (e.g. frequency range and peak frequencies) produced by bridge constructions and mapping (modeling) the distribution of noise within the affected marine area. While this ought to be conducted for each bridge construction, general research on typical fjord crossings will also provide valuable knowledge.
- Noise thresholds causing avoidance and physical injury for key species ought to be obtained from literature, or where necessary, experimentation.
- Combining mapping of distribution of bridge related noise and information on noise thresholds of key species will enable mapping of zones of impact for given species. This will enable conclusions to be drawn on whether the floating bridge construction causes physical injury (particularly during the construction process), habitat quality reduction as a result of avoidance, habitat quality as a result of masking of communication, or is a migration barrier preventing passage between two areas.
- In order to consider potential impacts relative to alternative crossing solutions such as traditional or electric ferries, studies on the character (e.g. frequencies) and extent of noise originating from these alternatives ought also to be conducted.

Light

- More research is needed on how birds will be attracted to light on bridges in weather conditions when they are known to gather around light sources like lighthouses, oil platforms or marker lights on high structures. This may cause collisions with structures, exhaustion after circling around them for a long time, or increased predation if they keep flying into sunrise (when they become more visible for predators). Knowledge on flight patterns or migration routes when crossing large bridges may help design mitigation measures, as rules for turning off light in short periods to let birds escape from the light. At the local level, knowledge of the local species is important to understand the extent of possible interference.

Habitat changes

- It is very important to gain more information on nature inclusive design/ecological engineering of the bridges, taking material composition, texture and macro-design into consideration. Optimizing these areas will strongly influence the benthic communities attaching to a floating bridge and its possible function as an artificial reef or as stepping stone for opportunistic or non-native species.
- To understand how floating bridges may function as migration barriers more research is needed on how migrating marine species (mainly mammals and fish) use the water column in this phase. Here should also the hydrodynamic changes caused by the bridge be included.
- Study how natural communities on the structures, or bio-protection, may increase durability and ecological stability and reduce maintenance cost.

Gateways

- One prerequisite for building new bridges and tunnels should be 100 % effective ports or barriers to prevent mammalian predators to get access to areas/islands where they are not present today. More research is needed on how predators behave when they meet such barriers, as well as their effectiveness by e.g. camera surveillance (infrared cameras or by developing methods for recognition by using artificial intelligence?). This could be done by using existing barriers, like those being built on the new Nordøyvegen.

Other

- More research is needed to see how birds will respond to (green) pontoons on submerged floating twin-tube bridges (SFTB) as potential breeding sites for some species, and how they should be designed (avoid chicks falling off them when running to seek shelter, but still make maintenance work possible).

Mitigation

- A consideration of the environmental aspects needs to be a part of the very early stages of a bridge construction.
- Environmental impact assessment – design a standard protocol on what needs to be investigated and how it needs to be carried out.
- Research on best design/way to carry out mitigation. E.g.
 - Best lighting regimes - how design of poles and lamps and choice of colour etc. can mitigate some effects (see section 4).
 - Best way to prevent SFTBs and bridges acting as gateways - we need to test their effectiveness and, if they can't be expected to be 100 %, establish routines for taking out predators immediately when passing of these barriers by a predator is observed.
 - Research into construction design to minimize vibration and noise generation.
 - Research existing measures to reduce the effects of the construction phase, e.g. bubble net curtains to reduce the radiation of noise produced during pile driving.
- Research into how to design structures to support native benthic communities that do not harm structures e.g. best material for native benthic community growth and how best to seed with indigenous species.

6 References

- Abadi, S., Flett, D., Berge, R., DeHaan, J., Guy, V., Qureshi, U. & Cook, M. 2018. Studying underwater sound level caused by bridge traffic in Lake Washington. *The Journal of the Acoustical Society of America* 144(3): 1808-1808.
- Acevedo-Gutierrez, A. & Cendejas-Zarelli, S. 2011. Nocturnal Haul-Out Patterns of Harbor Seals (*Phoca vitulina*) Related to Airborne Noise Levels in Bellingham, Washington, USA. *Aquatic Mammals* 37(2): 167-174. doi:10.1578/am.37.2.2011.167
- Andersson, K., Baldi, F., Brynolf, S., Lindgren, J.F., Granhag, L. & Svensson, E. 2016a. Shipping and the Environment. I: Anderson, K. (red.) Shipping and the Environment. S. 3-27.
- Andersson, M.H., Andersson, S., Ahlsen, J., Andersson, B.L., Hammar, J., Persson, L.K., Pihl, J., Sigra, P. & Wisstrom, A. 2016b. A framework for regulating underwater noise during pile driving. A technical Vindalreport. Swedish Environmental Protection Agency. <https://tethys.pnnl.gov/sites/default/files/publications/Andersson-et-al-2017-Report6775.pdf>
- Avery, M., Springer, P.F. & Cassel, J.F. 1976. The effects of a tall tower on nocturnal bird migration: a portable ceilometer study. *The Auk* 93(2): 281-291.
- Baker, H. 2020. Do seabirds have a better sense of hearing underwater than in air? <https://marinemadness.blog/2020/04/07/do-seabirds-have-a-better-sense-of-hearing-underwater-than-in-air/>.
- Baker, P., Fajans, J.S., Arnold, W.S., Ingrao, D.A., Marelli, D.C. & Baker, S.M. 2007. Range and dispersal of a tropical marine invader, the Asian green mussel, *Perna viridis*, in subtropical waters of the southeastern United States. *Journal of Shellfish Research* 26(2): 345-355. doi:10.2983/0730-8000(2007)26[345:Radoat]2.0.Co;2
- Bellebaum, J., Diederichs, A., Kube, J., Schulz, A. & Nehls, G. 2006. Flucht- und Meidedistanzen überwinternder Seetaucher und Meeressäuger gegenüber Schiffen auf See. *Ornithologischer Rundbrief Mecklenburg-Vorpommern* 45: 86-90.
- Bergan, P.I. & Giæver, M. 2002. Nordøyvegen i Haram og Sandøy kommune, Møre og Romsdal, konsekvensutredning naturmiljø. Rapport nr.: SG 559651-02. Statkraft Grøner AS, Trondheim
- Bergström, L., Sundqvist, F. & Bergström, U. 2013. Effects of an offshore wind farm on temporal and spatial patterns in the demersal fish community. *Marine Ecology Progress Series* 485: 199-210. doi:10.3354/meps10344
- Bergström, L., Kautsky, L., Malm, T., Rosenberg, R., Wahlberg, M., Capetillo, N.A. & Wilhelmsson, D. 2014. Effects of offshore wind farms on marine wildlife—a generalized impact assessment. *Environmental Research Letters* 9(3). doi:10.1088/1748-9326/9/3/034012
- Bishop, M.J., Mayer-Pinto, M., Airoidi, L., Firth, L.B., Morris, R.L., Loke, L.H.L., Hawkins, S.J., Naylor, L.A., Coleman, R.A., Chee, S.Y. & Dafforn, K.A. 2017. Effects of ocean sprawl on ecological connectivity: impacts and solutions. *Journal of Experimental Marine Biology and Ecology* 492: 7-30. doi:10.1016/j.jembe.2017.01.021
- Bjørge, A., Thompson, D., Hammond, P., Fedak, M., Bryant, E., Aarefjord, H., Roen, R. & Olsen, M. 1995. Habitat use and diving behaviour of harbour seals in a coastal archipelago in Norway. I: *Developments in marine biology*. Elsevier. S. 211-223. doi:[https://doi.org/10.1016/S0163-6995\(06\)80025-9](https://doi.org/10.1016/S0163-6995(06)80025-9)

- Bjørge, A. 2003. The Harbour Porpoise (*Phocoena phocoena*) in the North Atlantic: Variability in habitat use, trophic ecology and contaminant exposure. NAMMCO Scientific Publications 5: 223-228. doi:<https://doi.org/10.7557/3.2749>
- Blom, E.-L., Kvarnemo, C., Dekhla, I., Schöld, S., Andersson, M.H., Svensson, O. & Amorim, M.C.P. 2019. Continuous but not intermittent noise has a negative impact on mating success in a marine fish with paternal care. *Scientific reports* 9(1): 1-9.
- Bodey, T.W., Jessopp, M.J., Votier, S.C., Gerritsen, H.D., Cleasby, I.R., Hamer, K.C., Patrick, S.C., Wakefield, E.D. & Bearhop, S. 2014. Seabird movement reveals the ecological footprint of fishing vessels. *Current Biology* 24(11): R514-R515. doi:10.1016/j.cub.2014.04.041
- Bradbury, G., Trinder, M., Furness, B., Banks, A.N., Caldow, R.W.G. & Hume, D. 2014. Mapping Seabird Sensitivity to Offshore Wind Farms. *Plos One* 9(9). doi:10.1371/journal.pone.0106366
- Brandt, M.J., Diederichs, A., Betke, K. & Nehls, G. 2011. Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. *Marine Ecology Progress Series* 421: 205-216. doi:10.3354/meps08888
- Brandt, M.J., Dragon, A.C., Diederichs, A., Bellmann, M.A., Wahl, V., Piper, W., Nabe-Nielsen, J. & Nehls, G. 2018. Disturbance of harbour porpoises during construction of the first seven offshore wind farms in Germany. *Marine Ecology Progress Series* 596: 213-232. doi:10.3354/meps12560
- Bray, L., Reizopoulou, S., Voukouvalas, E., Soukissian, T., Alomar, C., Vazquez-Luis, M., Deudero, S., Attrill, M.J. & Hall-Spencer, J.M. 2016. Expected Effects of Offshore Wind Farms on Mediterranean Marine Life. *Journal of Marine Science and Engineering* 4(1). doi:10.3390/jmse4010018
- Buehler, D., Oestman, R., Reyff, J., Pommerenck, K. & Mitchell, B. 2015. Technical Guidance for Assessment and Mitigation of the Hydroacoustic Effects of Pile Driving on Fish CTHWANP-RT-15-306.01.01. California Department of Transportation
- Bulleri, F. & Airoidi, L. 2005. Artificial marine structures facilitate the spread of a non-indigenous green alga, *Codium fragile* ssp *tomentosoides*, in the north Adriatic Sea. *Journal of Applied Ecology* 42(6): 1063-1072. doi:10.1111/j.1365-2664.2005.01096.x
- Burger, C., Schubert, A., Heinanen, S., Dorsch, M., Kleinschmidt, B., Zydalis, R., Morkunas, J., Quillfeldt, P. & Nehls, G. 2019. A novel approach for assessing effects of ship traffic on distributions and movements of seabirds. *Journal of Environmental Management* 251. doi:10.1016/j.jenvman.2019.109511
- Cabrera-Cruz, S.A., Smolinsky, J.A. & Buler, J.J. 2018. Light pollution is greatest within migration passage areas for nocturnally-migrating birds around the world. *Scientific reports* 8: 3261. doi:10.1038/s41598-018-21577-6
- Carstensen, J., Henriksen, O.D. & Teilmann, J. 2006. Impacts of offshore wind farm construction on harbour porpoises: acoustic monitoring of echolocation activity using porpoise detectors (T-PODs). *Marine Ecology Progress Series* 321: 295-308. doi:10.3354/meps321295
- Celedonia, M.T., Tabor, R.A., Sanders, S., Damm, S., Lantz, D.W., Lee, T.M., Li, Z., Pratt, J.-M., Price, B.E. & Seyda, L. 2008. Movement and habitat use of Chinook salmon smolts, Northern pikeminnows, and smallmouth bass near the SR 520 Bridge. U.S. Fish and Wildlife Service
- Celi, M., Filiciotto, F., Maricchiolo, G., Genovese, L., Quinci, E.M., Maccarrone, V., Mazzola, S., Vazzana, M. & Buscaino, G. 2016. Vessel noise pollution as a human threat to fish:

- assessment of the stress response in gilthead sea bream (*Sparus aurata*, Linnaeus 1758). *Fish Physiology and Biochemistry* 42(2): 631-641. doi:10.1007/s10695-015-0165-3
- Christensen-Dalsgaard, S., Langset, M. & Anker-Nilssen, T. 2020. Offshore oil rigs—a breeding refuge for Norwegian Black-legged Kittiwakes *Rissa tridactyla*? *Seabird* 32: 20-32.
- Collins, A., Coughlin, D., Miller, J. & Kirk, S. 2015. The Production of Quick Scoping Reviews and Rapid Evidence Assessments. A How to Guide Department for Environment Food & Rural Affairs. <https://www.gov.uk/government/publications/the-production-of-quick-scoping-reviews-and-rapid-evidence-assessments>.
- Cominelli, S., Devillers, R., Yurk, H., MacGillivray, A., McWhinnie, L. & Canessa, R. 2018. Noise exposure from commercial shipping for the southern resident killer whale population. *Marine Pollution Bulletin* 136: 177-200. doi:10.1016/j.marpolbul.2018.08.050
- Connell, S.D. 2000. Floating pontoons create novel habitats for subtidal epibiota. *Journal of Experimental Marine Biology and Ecology* 247(2): 183-194. doi:10.1016/s0022-0981(00)00147-7
- Corbett, J.J. & Farrell, A. 2002. Mitigating air pollution impacts of passenger ferries. *Transportation Research Part D-Transport and Environment* 7(3): 197-211. doi:10.1016/s1361-9209(01)00019-0
- Cox, K., Brennan, L.P., Gerwing, T.G., Dudas, S.E. & Juanes, F. 2018. Sound the alarm: A meta-analysis on the effect of aquatic noise on fish behavior and physiology. *Global Change Biology* 24(7): 3105-3116. doi:10.1111/gcb.14106
- Crawford, R.L. & Engstrom, R.T. 2001. Characteristics of avian mortality at a north Florida television tower: a 29-year study. *Journal of Field Ornithology* 72(3): 380-388.
- Culloch, R.M., Anderwald, P., Brandecker, A., Haberlin, D., McGovern, B., Pinfield, R., Visser, F., Jessopp, M. & Cronin, M. 2016. Effect of construction-related activities and vessel traffic on marine mammals. *Marine Ecology Progress Series* 549: 231-242. doi:10.3354/meps11686
- Dauidsen, J.G., Rikardsen, A.H., Thorstad, E.B., Halttunen, E., Mitamura, H., Praebel, K., Skardhamar, J. & Naesje, T.F. 2013. Homing behaviour of Atlantic salmon (*Salmo salar*) during final phase of marine migration and river entry. *Canadian Journal of Fisheries and Aquatic Sciences* 70(5): 794-802. doi:10.1139/cjfas-2012-0352
- De Clippele, L.H. & Risch, D. 2021. Measuring Sound at a Cold-Water Coral Reef to Assess the Impact of COVID-19 on Noise Pollution. *Frontiers in Marine Science* 8. doi:10.3389/fmars.2021.674702
- de Jong, K., Forland, T.N., Amorim, M.C.P., Rieucan, G., Slabbekoorn, H. & Sivle, L.D. 2020. Predicting the effects of anthropogenic noise on fish reproduction. *Reviews in Fish Biology and Fisheries* 30(2): 245-268. doi:10.1007/s11160-020-09598-9
- Dehnhard, N., Skei, J., Christensen-Dalsgaard, S., May, R., Halley, D., Ringsby, T.H. & Lorentsen, S.H. 2019. Boat disturbance effects on moulting common eiders *Somateria mollissima*. *Marine Biology* 167(1). doi:10.1007/s00227-019-3624-z
- Dolman, S.J. & Jasny, M. 2015. Evolution of Marine Noise Pollution Management. *Aquatic Mammals* 41(4): 357-374. doi:10.1578/am.41.4.2015.357
- Duarte, C.M., Chapuis, L., Collin, S.P., Costa, D.P., Devassy, R.P., Eguiluz, V.M., Erbe, C., Gordon, T.A.C., Halpern, B.S., Harding, H.R., Havlik, M.N., Meekan, M., Merchant, N.D., Miksis-Olds, J.L., Parsons, M., Predragovic, M., Radford, A.N., Radford, C.A., Simpson, S.D., Slabbekoorn, H., Staaterman, E., Van Opzeeland, I.C., Winderen, J., Zhang, X.L. & Juanes,

- F. 2021. The soundscape of the Anthropocene ocean. *Science* 371(6529): eaba4658. doi:10.1126/science.aba4658
- Dähne, M., Gilles, A., Lucke, K., Peschko, V., Adler, S., Krugel, K., Sundermeyer, J. & Siebert, U. 2013. Effects of pile-driving on harbour porpoises (*Phocoena phocoena*) at the first offshore wind farm in Germany. *Environmental Research Letters* 8(2). doi:10.1088/1748-9326/8/2/025002
- Ellison, W., Southall, B., Clark, C. & Frankel, A. 2012. A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. *Conservation Biology* 26(1): 21-28.
- Erbe, C. 2012. Effects of Underwater Noise on Marine Mammals. I: Popper, A. N. & Hawkins, A. (red.) *The Effects of Noise on Aquatic Life*. Springer New York : Imprint: Springer, New York, NY
- Erbe, C., MacGillivray, A. & Williams, R. 2012. Mapping cumulative noise from shipping to inform marine spatial planning. *Journal of the Acoustical Society of America* 132(5): EL423-EL428. doi:10.1121/1.4758779
- Erbe, C. 2013. Underwater noise of small personal watercraft (jet skis). *Journal of the Acoustical Society of America* 133(4): EL326-EL330. doi:10.1121/1.4795220
- EU. 2008. Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive). *Official Journal of the European Union*, L 164, 19–40 Available at www.eur-lex.europa.eu. <https://eur-lex.europa.eu>
- Evans, W. 2010. Response to: green light for nocturnally migrating birds. *Ecology and society* 15(3): r1.
- Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C., Elvidge, C., Baugh, K., Portnov, B., Rybnikova, N. & Furgoni, R. 2016. The new world atlas of artificial night sky brightness. *Science Advances* 2(6).
- Fliessbach, K.L., Borkenhagen, K., Guse, N., Markones, N., Schwemmer, P. & Garthe, S. 2019. A Ship Traffic Disturbance Vulnerability Index for Northwest European Seabirds as a Tool for Marine Spatial Planning. *Frontiers in Marine Science* 6. doi:10.3389/fmars.2019.00192
- Follestad, A., Heggberget, T.M., Hoem, S.A., Nygård, T., Reitan, O. & Røv, N. 2005. Arealbruk på kysten påvirker dyrelivet. I: Heggberget, T. M. & Jonsson, B. (red.) *Landskapsøkologi: Arealbruk og landskapsanalyse*. NINAs strategiske instituttprogrammer 2001-2005. NINA Temahefte 32. Norsk institutt for naturforskning, Trondheim
- Follestad, A. 2014. Effekter av kunstig nattbelysning på naturmangfoldet - en litteraturstudie. NINA Rapport 1081. Norsk institutt for naturforskning
- Furness, R.W., Wade, H.M. & Masden, E.A. 2013. Assessing vulnerability of marine bird populations to offshore wind farms. *Journal of Environmental Management* 119: 56-66. doi:10.1016/j.jenvman.2013.01.025
- Garthe, S. & Huppopp, O. 2004. Scaling possible adverse effects of marine wind farms on seabirds: developing and applying a vulnerability index. *Journal of Applied Ecology* 41(4): 724-734. doi:10.1111/j.0021-8901.2004.00918.x
- Gehring, J., Kerlinger, P. & Manville, A.M. 2009. Communication towers, lights, and birds: successful methods of reducing the frequency of avian collisions. *Ecological Applications* 19(2): 505-514. doi:doi.org/10.1890/07-1708.1

- Gentry, R.L. 2002. National policy on the effects of underwater noise on marine mammals and turtles. *The Journal of the Acoustical Society of America* 111(5): 2396-2396.
- Gervaise, C., Simard, Y., Roy, N., Kinda, B. & Menard, N. 2012. Shipping noise in whale habitat: Characteristics, sources, budget, and impact on belugas in Saguenay-St. Lawrence Marine Park hub. *Journal of the Acoustical Society of America* 132(1): 76-89. doi:10.1121/1.4728190
- Ghanim, A.D., Ibrahim, M.M. & Hassan, M.A. 2021. Effect of floating bridges on bottom topography. *Ain Shams Engineering Journal* 12(1): 475-486. doi:10.1016/j.asej.2020.09.013
- Grøsvik, B.E., Prokhorova, T., Eriksen, E., Krivosheya, P., Horneland, P.A. & Prozorkevich, D. 2018. Assessment of marine litter in the Barents Sea, a part of the Joint Norwegian–Russian Ecosystem Survey. *Frontiers in Marine Science* 5: 72. doi:10.3389/fmars.2018.00072
- Hadary, T., Martínez, J.G., Sella, I. & Perkol-Finkel, S. 2022. Eco-engineering for Climate Change—Floating to the Future. I: Piątek, L., Lim, S. H., Wang, C. H. & Graaf-van Dinther, R. (red.) WCFS2020. Proceedings of the Second World Conference on Floating Solutions, Rotterdam. Springer, Singapore
- Halvorsen, M.B., Casper, B.M., Matthews, F., Carlson, T.J. & Popper, A.N. 2012a. Effects of exposure to pile-driving sounds on the lake sturgeon, Nile tilapia and hogchoker. *Proceedings of the Royal Society B-Biological Sciences* 279(1748): 4705-4714. doi:10.1098/rspb.2012.1544
- Halvorsen, M.B., Casper, B.M., Woodley, C.M., Carlson, T.J. & Popper, A.N. 2012b. Threshold for Onset of Injury in Chinook Salmon from Exposure to Impulsive Pile Driving Sounds. *Plos One* 7(6). doi:10.1371/journal.pone.0038968
- Hammond, P.S., Lacey, C., Gilles, A., Viquerat, S., Börjesson, P., Herr, H., Macleod, K., Ri-doux, V., Santos, M.B., Scheidat, M., Teilmann, J., Vingada, J. & Øien, N. 2017. Estimates of cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys. Sea Mammal Research Unit, University of St Andrews. <https://synergy.st-andrews.ac.uk/scans3/files/2017/04/SCANS-III-design-based-estimates-2017-04-28-final.pdf>
- Hastie, G.D., Russell, D.J.F., McConnell, B., Moss, S., Thompson, D. & Janik, V.M. 2015. Sound exposure in harbour seals during the installation of an offshore wind farm: predictions of auditory damage. *Journal of Applied Ecology* 52(3): 631-640. doi:10.1111/1365-2664.12403
- Holloway, M.G. & Connell, S.D. 2002. Why do floating structures create novel habitats for subtidal epibiota? *Marine Ecology Progress Series* 235: 43-52. doi:10.3354/meps235043
- Horton, K.G., Nilsson, C., Van Doren, B.M., La Sorte, F.A., Dokter, A.M. & Farnsworth, A. 2019. Bright lights in the big cities: migratory birds' exposure to artificial light. *Frontiers in Ecology and the Environment* 17(4): 209-214. doi:10.1002/fee.2029
- Jelle, Fay, R.R., Tavalga, W.N. & International Conference on the Sensory Biology of Aquatic, A. 1988. *Sensory Biology of Aquatic Animals. Papers Based on Presentations Given at an International Conference on the Sensory Biology of Aquatic Animals Held, June 24-28, 1985, at the Mote Marine Laboratory in Sarasota, Fla.* Springer New York : Imprint: Springer, New York, NY.
- Jones, J. & Francis, C.M. 2003. The effects of light characteristics on avian mortality at lighthouses. *Journal of Avian Biology* 34(4): 328-333. doi:10.1111/j.0908-8857.2003.03183.x
- Jägerbrand, A.K., Brutemark, A., Sveden, J.B. & Gren, I.M. 2019. A review on the environmental impacts of shipping on aquatic and nearshore ecosystems. *Science of the Total Environment*

695. doi:10.1016/j.scitotenv.2019.133637

- Kaplan, M.B., Mooney, T.A., Lammers, M.O. & Zang, E. 2016. Temporal and spatial variability in vessel noise on tropical coral reefs. *Proceedings of Meetings on Acoustics 4ENAL*. Proceedings
- Kastelein, R.A., van Heerden, D., Gransier, R. & Hoek, L. 2013. Behavioral responses of a harbor porpoise (*Phocoena phocoena*) to playbacks of broadband pile driving sounds. *Marine Environmental Research* 92: 206-214. doi:10.1016/j.marenvres.2013.09.020
- Kerlinger, P., Gehring, J.L., Erickson, W.P., Curry, R., Jain, A. & Guarnaccia, J. 2010. Night migrant fatalities and obstruction lighting at wind turbines in North America. *The Wilson Journal of Ornithology* 122(4): 744-754. doi:10.1676/06-075.1
- Khangaonkar, T., Nugraha, A. & Wang, T.P. 2018a. Hydrodynamic Zone of Influence Due to a Floating Structure in a Fjordal Estuary-Hood Canal Bridge Impact Assessment. *Journal of Marine Science and Engineering* 6(4). doi:10.3390/jmse6040119
- Khangaonkar, T., Nugraha, A., Xu, W.W., Long, W., Bianucci, L., Ahmed, A., Mohamedali, T. & Pelletier, G. 2018b. Analysis of Hypoxia and Sensitivity to Nutrient Pollution in Salish Sea. *Journal of Geophysical Research-Oceans* 123(7): 4735-4761. doi:10.1029/2017jc013650
- Kjellman, M. 2015. Depth use of adult Atlantic salmon during the first and last phase of the marine migration. MSc-thesis. The Arctic University of Norway
- Koschinski, S., Culik, B.M., Henriksen, O.D., Tregenza, N., Ellis, G., Jansen, C. & Kathe, G. 2003. Behavioural reactions of free-ranging porpoises and seals to the noise of a simulated 2 MW windpower generator. *Marine Ecology Progress Series* 265: 263-273. doi:10.3354/meps265263
- Kuşku, H., Yiğit, M., Ergün, S., Yiğit, Ü. & Taylor, N. 2018. Acoustic noise pollution from marine industrial activities: Exposure and impacts. *Aquatic Research* 1(4): 148-161.
- La Sorte, F.A., Fink, D., Buler, J.J., Farnsworth, A. & Cabrera-Cruz, S.A. 2017. Seasonal associations with urban light pollution for nocturnally migrating bird populations. *Global Change Biology* 23(11): 4609-4619. doi: 10.1111/gcb.13792
- Lacombe, S., Amélineau, F. & Tarroux, A. 2021. Do North Atlantic pelagic seabirds take advantage of winds and currents when they migrate? Oral presentation on 7th SEATRACK workshop/webinar 1-3 November 2021.
- Le Bot, T., Lescroel, A. & Gremillet, D. 2018. A toolkit to study seabird-fishery interactions. *Ices Journal of Marine Science* 75(5): 1513-1525. doi:10.1093/icesjms/fsy038
- Lesage, V., Hammill, M.O. & Kovacs, K.M. 1999. Functional classification of harbor seal (*Phoca vitulina*) dives using depth profiles, swimming velocity, and an index of foraging success. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 77(1): 74-87. doi:10.1139/z98-199
- Lindeboom, H.J., Kouwenhoven, H.J., Bergman, M.J.N., Bouma, S., Brasseur, S., Daan, R., Fijn, R.C., de Haan, D., Dirksen, S., van Hal, R., Lambers, R.H.R., Ter Hofstede, R., Krijgsveld, K.L., Leopold, M. & Scheidat, M. 2011. Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. *Environmental Research Letters* 6(3). doi:10.1088/1748-9326/6/3/035101
- Longcore, T., Rich, C. & Gauthreaux Jr, S.A. 2008. Height, guy wires, and steady-burning lights increase hazard of communication towers to nocturnal migrants: a review and meta-analysis. *The Auk* 125(2): 485-492. doi:10.1525/auk.2008.06253

- Luczkovich, J.J. & Keusenkothen, M.A. 2008. Can longspine squirrelfish hear bottlenose dolphin? *Bioacoustics* 17(1-3): 75-77.
- Martin, D., Bertasi, F., Colangelo, M.A., de Vries, M., Frost, M., Hawkins, S.J., Macpherson, E., Moschella, P.S., Satta, M.P., Thompson, R.C. & Ceccherelli, V.U. 2005. Ecological impact of coastal defence structures on sediment and mobile fauna: Evaluating and forecasting consequences of unavoidable modifications of native habitats. *Coastal Engineering* 52(10-11): 1027-1051. doi:10.1016/j.coastaleng.2005.09.006
- McCauley, R.D., Day, R.D., Swadling, K.M., Fitzgibbon, Q.P., Watson, R.A. & Semmens, J.M. 2017. Widely used marine seismic survey air gun operations negatively impact zooplankton. *Nature Ecology & Evolution* 1(7). doi:10.1038/s41559-017-0195
- McLaren, J.D., Buler, J.J., Schreckengost, T., Smolinsky, J.A., Boone, M., Emiel van Loon, E., Dawson, D.K. & Walters, E.L. 2018. Artificial light at night confounds broad-scale habitat use by migrating birds. *Ecology Letters* 21(3): 356-364. doi:10.1111/ele.12902
- Mendel, B., Schwemmer, P., Peschko, V., Muller, S., Schwemmer, H., Mercker, M. & Garthe, S. 2019. Operational offshore wind farms and associated ship traffic cause profound changes in distribution patterns of Loons (*Gavia* spp.). *Journal of Environmental Management* 231: 429-438. doi:10.1016/j.jenvman.2018.10.053
- Merkel, F.R. & Johansen, K.L. 2011. Light-induced bird strikes on vessels in Southwest Greenland. *Marine Pollution Bulletin* 62(11): 2330-2336. doi:10.1016/j.marpolbul.2011.08.040
- Momota, K. & Hosokawa, S. 2021. Potential impacts of marine urbanization on benthic macrofaunal diversity. *Scientific Reports* 11(1). doi:10.1038/s41598-021-83597-z
- Montgomery, J.C., Jeffs, A., Simpson, S.D., Meekan, M. & Tindle, C. 2006. Sound as an orientation cue for the pelagic larvae of reef fishes and decapod crustaceans. *Advances in marine biology* 51: 143-196.
- Mooney, T.A., Smith, A., Hansen, K.A., Larsen, O.N., Wahlberg, M. & Rasmussen, M. 2019. Birds of a feather: hearing and potential noise impacts in puffins (*Fratercula arctica*). *Proceedings of Meetings on Acoustics* 5ENAL. Proceedings
- Moore, M., Berejikian, B.A. & Tezak, E.P. 2013. A Floating Bridge Disrupts Seaward Migration and Increases Mortality of Steelhead Smolts in Hood Canal, Washington State. *Plos One* 8(9). doi:10.1371/journal.pone.0073427
- Moschella, P.S., Abbiati, M., Aberg, P., Airoidi, L., Anderson, J.M., Bacchiocchi, F., Bulleri, F., Dinesen, G.E., Frost, M., Gacia, E., Granhag, L., Jonsson, P.R., Satta, M.P., Sundelof, A., Thompson, R.C. & Hawkins, S.J. 2005. Low-crested coastal defence structures as artificial habitats for marine life: Using ecological criteria in design. *Coastal Engineering* 52(10-11): 1053-1071. doi:10.1016/j.coastaleng.2005.09.014
- Murchy, K.A., Davies, H., Shafer, H., Cox, K., Nikolich, K. & Juanes, F. 2019. Impacts of noise on the behavior and physiology of marine invertebrates: A meta-analysis. *Proceedings of Meetings on Acoustics* 5ENAL. Proceedings
- Maar, M., Bolding, K., Petersen, J.K., Hansen, J.L.S. & Timmermann, K. 2009. Local effects of blue mussels around turbine foundations in an ecosystem model of Nysted off-shore wind farm, Denmark. *Journal of Sea Research* 62(2-3): 159-174. doi:10.1016/j.seares.2009.01.008
- Nabe-Nielsen, J., Sibly, R.M., Tougaard, J., Teilmann, J. & Sveegaard, S. 2014. Effects of noise and by-catch on a Danish harbour porpoise population. *Ecological Modelling* 272: 242-251. doi:10.1016/j.ecolmodel.2013.09.025

- Nielsen, N.H., Teilmann, J., Sveegaard, S., Hansen, R.G., Sinding, M.H.S., Dietz, R. & Heide-Jorgensen, M.P. 2018. Oceanic movements, site fidelity and deep diving in harbour porpoises from Greenland show limited similarities to animals from the North Sea. *Marine Ecology Progress Series* 597: 259-272. doi:10.3354/meps12588
- Olsen, M.T., Andersen, S.M., Teilmann, J., Dietz, R., Edrén, S.M.C., Linnet, A. & Härkönen, T. 2010. Status of the harbour seal (*Phoca vitulina*) in Southern Scandinavia. *NAMMCO Scientific Publications* 8: 77-94.
- Parsons, M.J.G., Duncan, A.J., Parsons, S.K. & Erbe, C. 2020. Reducing vessel noise: An example of a solar-electric passenger ferry. *Journal of the Acoustical Society of America* 147(5): 3575-3583. doi:10.1121/10.0001264
- Perkol-Finkel, S., Zilman, G., Sella, I., Miloh, T. & Benayahu, Y. 2006. Floating and fixed artificial habitats: effects of substratum motion on benthic communities in a coral reef environment. *Marine Ecology Progress Series* 317: 9-20. doi:10.3354/meps317009
- Perkol-Finkel, S., Zilman, G., Sella, I., Miloh, T. & Benayahu, Y. 2008. Floating and fixed artificial habitats: Spatial and temporal patterns of benthic communities in a coral reef environment. *Estuarine Coastal and Shelf Science* 77(3): 491-500. doi:10.1016/j.ecss.2007.10.005
- Picken, G., Baine, M., Heaps, L. & Side, J. 2000. Rigs to reefs in the North Sea. I: Lockwood, A. P. M., Jensen, A. C. & Collins, K. J. (red.) *Artificial reefs in European seas*. Kluwer Academic, Dordrecht. S. 331-342.
- Pine, M.K., Jeffs, A.G., Wang, D. & Radford, C.A. 2016. The potential for vessel noise to mask biologically important sounds within ecologically significant embayments. *Ocean & Coastal Management* 127: 63-73.
- Poot, H., Ens, B.J., de Vries, H., Donners, M.A., Wernand, M.R. & Marquenie, J.M. 2008. Green light for nocturnally migrating birds. *Ecology and Society* 13(2): 47.
- Popper, A.N. & Hawkins, A.D. 2019. An overview of fish bioacoustics and the impacts of anthropogenic sounds on fishes. *Journal of Fish Biology* 94(5): 692-713. doi:10.1111/jfb.13948
- Purser, J. & Radford, A.N. 2011. Acoustic noise induces attention shifts and reduces foraging performance in three-spined sticklebacks (*Gasterosteus aculeatus*). *PLoS One* 6(2): e17478.
- Remage-Healey, L. & Bass, A.H. 2006. From social behavior to neural circuitry: steroid hormones rapidly modulate advertisement calling via a vocal pattern generator. *Hormones and behavior* 50(3): 432-441.
- Roberts, L. & Laidre, M.E. 2019. Noise alters chemically-mediated search behavior in a marine hermit crab: Studying cross-modal effects on behavior. *Proceedings of Meetings on Acoustics* 5ENAL. *Proceedings*
- Rooker, J.R., Dokken, Q.R., Pattengill, C.V. & Holt, G.J. 1997. Fish assemblages on artificial and natural reefs in the Flower Garden Banks National Marine Sanctuary, USA. *Coral Reefs* 16(2): 83-92. doi:10.1007/s003380050062
- Ross, D. 1993. On Ocean Ambient Noise. *Acoustics Bulletin* 18.
- Smith, M.E. & Monroe, J.D. 2016. Causes and consequences of sensoryhair cell damage and recovery in fishes. I: Sisneros, J. A. (red.) *Fish Hearing and Bioacoustics : An Anthology in Honor of Arthur N. Popper and Richard R. Fay*. Springer International Publishing : Imprint: Springer

- Schoeman, R.P., Patterson-Abrolat, C. & Plön, S. 2020. A global review of vessel collisions with marine animals. *Frontiers in Marine Science* 7: 292. doi:10.3389/fmars.2020.00292
- Schwemmer, P., Mendel, B., Sonntag, N., Dierschke, V. & Garthe, S. 2011. Effects of ship traffic on seabirds in offshore waters: implications for marine conservation and spatial planning. *Ecological Applications* 21(5): 1851-1860. doi:10.1890/10-0615.1
- Song, X.D., Zhang, X.G., Xiong, W., Guo, Z.M. & Wang, B. 2020. Experimental and numerical study on underwater noise radiation from an underwater tunnel. *Environmental Pollution* 267. doi:10.1016/j.envpol.2020.115536
- Sommerfeld, J., Mendel, B., Fock, H.O. & Garthe, S. 2016. Combining bird-borne tracking and vessel monitoring system data to assess discard use by a scavenging marine predator, the lesser black-backed gull *Larus fuscus*. *Marine Biology* 163(5). doi:10.1007/s00227-016-2889-8
- Spinuzzi, S., Schneider, K.R., Walters, L.J., Yuan, W.S. & Hoffman, E.A. 2013. Tracking the distribution of non-native marine invertebrates (*Mytella charruana*, *Perna viridis* and *Megabalanus coccopoma*) along the south-eastern USA. *Marine Biodiversity Records* 6.
- Stanley, J.A., Radford, C.A. & Jeffs, A.G. 2012. The Effects of Noise on Aquatic Life. I: Popper, A. N. & Hawkins, A. (red.) *The Effects of Noise on Aquatic Life*. Springer New York : Imprint: Springer, New York, NY. S. 371-374.
- Sørensen, K., Neumann, C., Dahne, M., Hansen, K.A. & Wahlberg, M. 2020. Gentoo penguins (*Pygoscelis papua*) react to underwater sounds. *Royal Society Open Science* 7(2). doi:10.1098/rsos.191988
- Terhune, J.M. & Killorn, D. 2021. A Method for Preliminary Assessment of the Masking Potential of Anthropogenic Noise to Baleen Whale Calls. *Aquatic Mammals* 47(3): 283-291. doi:10.1578/am.47.3.2021.283
- Tew-Kai, E., Benhamou, S., van der Lingen, C.D., Coetzee, J.C., Pichegru, L., Ryan, P.G. & Gremillet, D. 2013. Are Cape gannets dependent upon fishery waste? A multi-scale analysis using seabird GPS-tracking, hydro-acoustic surveys of pelagic fish and vessel monitoring systems. *Journal of Applied Ecology* 50(3): 659-670. doi:10.1111/1365-2664.12086
- Thingstad, P.G. & Hokstad, S. 1997. Vannfugl og marin bunndyrfauna i Kråkvågsvaet, Ørland kommune, Sør-Trøndelag. Konsekvenser av eventuell bru og veifylling over svaet. Rapport Zoologisk Serie 1997-2. NTNU Vitenskapsmuseet
- Thingstad, P.G., Hokstad, S., Frengen, O. & Strømgren, T. 1994. Vannfugl og marin bunndyrfauna i Ramsarområdet på Tautra, Nord-Trøndelag. Konsekvenser av steinmoloen over svaet. Rapport Zoologisk Serie 1994-8. Universitetet i Trondheim, Vitenskapsmuseet
- Thingstad, P.G. & Hokstad, S. 1997. Vannfugl og marin bunndyrfauna i Kråkvågsvaet, Ørland kommune, Sør-Trøndelag. Konsekvenser av eventuell bru og veifylling over svaet. Rapport Zoologisk Serie 1997-2. NTNU Vitenskapsmuseet
- Thomsen, F., Lüdemann, K., Kafemann, R. & Piper, W. 2006. Effects of offshore wind farm noise on marine mammals and fish biola. https://tethys.pnnl.gov/sites/default/files/publications/Effects_of_offshore_wind_farm_noise_on_marine-mammals_and_fish-1-.pdf
- Thomson, D.J. & Barclay, D.R. 2020. Real-time observations of the impact of COVID-19 on underwater noise. *The Journal of the Acoustical Society of America* 147(5): 3390-3396.
- Thorstad, E.B., Whoriskey, F., Uglem, I., Moore, A., Rikardsen, A.H. & Finstad, B. 2012. A critical life stage of the Atlantic salmon *Salmo salar*: behaviour and survival during the smolt and

- initial post-smolt migration. *Journal of Fish Biology* 81(2): 500-542. doi:10.1111/j.1095-8649.2012.03370.x
- Thorstad, E.B., Todd, C.D., Uglem, I., Bjorn, P.A., Gargan, P.G., Vollset, K.W., Halttunen, E., Kalas, S., Berg, M. & Finstad, B. 2016. Marine life of the sea trout. *Marine Biology* 163(3). doi:10.1007/s00227-016-2820-3
- Van Doren, B.M., Horton, K.G., Dokter, A.M., Klinck, H., Elbin, S.B. & Farnsworth, A. 2017. High-intensity urban light installation dramatically alters nocturnal bird migration. *Proceedings of the National Academy of Sciences* 114(42): 11175-11180. doi:10.1073/pnas.1708574114
- Van Renterghem, T., Botteldooren, D. & Dekoninck, L. 2014. Airborne sound propagation over sea during offshore wind farm piling. *Journal of the Acoustical Society of America* 135(2): 599-609. doi:10.1121/1.4861244
- Vandendriessche, S., Derweduwen, J. & Hostens, K. 2015. Equivocal effects of offshore wind farms in Belgium on soft substrate epibenthos and fish assemblages. *Hydrobiologia* 756(1): 19-35.
- Verfuß, T. 2014. Noise mitigation systems and low-noise installation technologies. I: Ecological Research at the Offshore Windfarm alpha ventus. Springer, Wiesbaden. S. 181-191. doi:https://doi.org/10.1007/978-3-658-02462-8_16
- Votier, S.C., Bearhop, S., Witt, M.J., Inger, R., Thompson, D. & Newton, J. 2010. Individual responses of seabirds to commercial fisheries revealed using GPS tracking, stable isotopes and vessel monitoring systems. *Journal of Applied Ecology* 47(2): 487-497. doi:10.1111/j.1365-2664.2010.01790.x
- Walker, S.J., Schlacher, T.A. & Thompson, L.M.C. 2008. Habitat modification in a dynamic environment: The influence of a small artificial groyne on macrofaunal assemblages of a sandy beach. *Estuarine Coastal and Shelf Science* 79(1): 24-34. doi:10.1016/j.ecss.2008.03.011
- Westgate, A.J., Read, A.J., Berggren, P., Koopman, H.N. & Gaskin, D.E. 1995. Diving behavior of harbor porpoises, *phocoena-phocoena*. *Canadian Journal of Fisheries and Aquatic Sciences* 52(5): 1064-1073. doi:10.1139/f95-104
- Whomersley, P. & Picken, G. 2003. Long-term dynamics of fouling communities found on offshore installations in the North Sea. *Journal of the Marine Biological Association of the United Kingdom* 83(5): 897-901.
- Wood, M.P. & Carter, L. 2008. Whale Entanglements With Submarine Telecommunication Cables. *IEEE Journal of Oceanic Engineering* 33(4): 445-450. doi:10.1109/joe.2008.2001638
- Wursig, B., Greene, C.R. & Jefferson, T.A. 2000. Development of an air bubble curtain to reduce underwater noise of percussive piling. *Marine Environmental Research* 49(1): 79-93. doi:10.1016/s0141-1136(99)00050-1
- Zhao, X., Chen, M., Wu, Z. & Wang, Z. 2014. Factors influencing phototaxis in nocturnal migrating birds. *Zoological Science* 31(12): 781-788. doi:10.2108/zs130237
- Zhao, X.B., Zhang, M., Che, X.L. & Zou, F.S. 2020. Blue light attracts nocturnally migrating birds. *Condor* 122(2). doi:10.1093/condor/duaa002
- Øigård, T.A., Frie, A.K., Nilssen, K.T. & Hammill, M.O. 2012. Modelling the abundance of grey seals (*Halichoerus grypus*) along the Norwegian coast. *Ices Journal of Marine Science* 69(8): 1436-1447. doi:10.1093/icesjms/fss103

7 Appendix: Included papers from WoS literature search

Our literature search in Web of Science on October 26th 2021 yielded 1441 articles, and all articles were screened. A systematic literature search tends to have a wide scope, and will always return a large number of irrelevant hits. The following 193 papers were considered relevant for our project:

- Amaral, J.L., Miller, J.H., Potty, G.R., Vigness-Raposa, K.J., Frankel, A.S., Lin, Y.T., Newhall, A.E., Wilkes, D.R. & Gavrilov, A.N. 2020. Characterization of impact pile driving signals during installation of offshore wind turbine foundations. *Journal of the Acoustical Society of America* 147(4): 2323-2333. doi:10.1121/10.0001035
- Andersson, M.H., Berggren, M., Wilhelmsson, D. & Ohman, M.C. 2009. Epibenthic colonization of concrete and steel pilings in a cold-temperate embayment: a field experiment. *Helgoland Marine Research* 63(3): 249-260. doi:10.1007/s10152-009-0156-9
- Aniceto, A.S., Carroll, J., Tetley, M.J. & van Oosterhout, C. 2016. Position, swimming direction and group size of fin whales (*Balaenoptera physalus*) in the presence of a fast-ferry in the Bay of Biscay. *Oceanologia* 58(3): 235-240. doi:10.1016/j.oceano.2016.02.002
- Bailey, H., Hammond, P.S. & Thompson, P.M. 2014. Modelling harbour seal habitat by combining data from multiple tracking. *Journal of Experimental Marine Biology and Ecology* 450: 30-39. doi:10.1016/j.jembe.2013.10.011
- Baltzer, J., Maurer, N., Schaffeld, T., Ruser, A., Schnitzler, J.G. & Siebert, U. 2020. Effect ranges of underwater noise from anchor vibration operations in the Wadden Sea. *Journal of Sea Research* 162. doi:10.1016/j.seares.2020.101912
- Benjamins, S., van Geel, N., Hastie, G., Elliott, J. & Wilson, B. 2017. Harbour porpoise distribution can vary at small spatiotemporal scales in energetic habitats. *Deep-Sea Research Part II: Topical Studies in Oceanography* 141: 191-202. doi:10.1016/j.dsr2.2016.07.002
- Bergström, L., Sundqvist, F. & Bergstrom, U. 2013. Effects of an offshore wind farm on temporal and spatial patterns in the demersal fish community. *Marine Ecology Progress Series* 485: 199-210. doi:10.3354/meps10344
- Bergström, L., Kautsky, L., Malm, T., Rosenberg, R., Wahlberg, M., Capetillo, N.A. & Wilhelmsson, D. 2014. Effects of offshore wind farms on marine wildlife-a generalized impact assessment. *Environmental Research Letters* 9(3). doi:10.1088/1748-9326/9/3/034012
- Bicknell, A.W.J., Sheehan, E.V., Godley, B.J., Doherty, P.D. & Witt, M.J. 2019. Assessing the impact of introduced infrastructure at sea with cameras: A case study for spatial scale, time and statistical power. *Marine Environmental Research* 147: 126-137. doi:10.1016/j.marenvres.2019.04.007
- Bochert, R. & Zettler, M.L. 2004. Long-term exposure of several marine benthic animals to static magnetic fields. *Bioelectromagnetics* 25(7): 498-502. doi:10.1002/bem.20019
- Bolgan, M., Chorazyczewska, E., Winfield, I.J., Codarin, A., O'Brien, J. & Gammell, M. 2016. First observations of anthropogenic underwater noise in a large multi-use lake. *Journal of Limnology* 75(3): 644-651. doi:10.4081/jlimnol.2016.1405
- Bouveroux, T., Waggitt, J.J., Belhadjer, A., Cazenave, P.W., Evans, P.G.H. & Kiszka, J.J. 2020. Modelling fine-scale distribution and relative abundance of harbour porpoises in the Southern Bight of the North Sea using platform-of-opportunity data. *Journal of the Marine Biological Association of the United Kingdom* 100(3): 481-489. doi:10.1017/s0025315420000326

- Brandt, M.J., Diederichs, A., Betke, K. & Nehls, G. 2011. Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. *Marine Ecology Progress Series* 421: 205-216. doi:10.3354/meps08888
- Brandt, M.J., Hoschle, C., Diederichs, A., Betke, K., Matuschek, R. & Nehls, G. 2013. Seal scarers as a tool to deter harbour porpoises from offshore construction sites. *Marine Ecology Progress Series* 475: 291-302. doi:10.3354/meps10100
- Brandt, M.J., Dragon, A.C., Diederichs, A., Bellmann, M.A., Wahl, V., Piper, W., Nabe-Nielsen, J. & Nehls, G. 2018. Disturbance of harbour porpoises during construction of the first seven offshore wind farms in Germany. *Marine Ecology Progress Series* 596: 213-232. doi:10.3354/meps12560
- Bray, L., Reizopoulou, S., Voukouvalas, E., Soukissian, T., Alomar, C., Vazquez-Luis, M., Deudero, S., Attrill, M.J. & Hall-Spencer, J.M. 2016. Expected Effects of Offshore Wind Farms on Mediterranean Marine Life. *Journal of Marine Science and Engineering* 4(1). doi:10.3390/jmse4010018
- Bray, L., Kassis, D. & Hall-Spencer, J.M. 2017. Assessing larval connectivity for marine spatial planning in the Adriatic. *Marine Environmental Research* 125: 73-81. doi:10.1016/j.marenvres.2017.01.006
- Bruderer, B., Peter, D. & Korner-Nievergelt, F. 2018. Vertical distribution of bird migration between the Baltic Sea and the Sahara. *Journal of Ornithology* 159(2): 315-336. doi:10.1007/s10336-017-1506-z
- Buck, B.H., Krause, G., Michler-Cieluch, T., Brenner, M., Buchholz, C.M., Busch, J.A., Fisch, R., Geisen, M. & Zielinski, O. 2008. Meeting the quest for spatial efficiency: progress and prospects of extensive aquaculture within offshore wind farms. *Helgoland Marine Research* 62(3): 269-281. doi:10.1007/s10152-008-0115-x
- Butkus, D., Grubliauskas, R. & Mazuolis, J. 2012. Research of equivalent and maximum value of noise generated by wind power plants. *Journal of Environmental Engineering and Landscape Management* 20(1): 27-34. doi:10.3846/16486897.2011.633337
- Cafaro, V., Piazzolla, D., Melchiorri, C., Burgio, C., Fersini, G., Conversano, F., Piermattei, V. & Marcelli, M. 2018. Underwater noise assessment outside harbor areas: The case of Port of Civitavecchia, northern Tyrrhenian Sea, Italy. *Marine Pollution Bulletin* 133: 865-871. doi:10.1016/j.marpolbul.2018.06.058
- Carstensen, J., Henriksen, O.D. & Teilmann, J. 2006. Impacts of offshore wind farm construction on harbour porpoises: acoustic monitoring of echolocation activity using porpoise detectors (T-PODs). *Marine Ecology Progress Series* 321: 295-308. doi:10.3354/meps321295
- Castro, J.J., Santiago, J.A. & Santana-Ortega, A.T. 2001. A general theory on fish aggregation to floating objects: An alternative to the meeting point hypothesis. *Reviews in Fish Biology and Fisheries* 11(3): 255-277. doi:10.1023/a:1020302414472
- Celi, M., Filiciotto, F., Maricchiolo, G., Genovese, L., Quinci, E.M., Maccarrone, V., Mazzola, S., Vazzana, M. & Buscaino, G. 2016. Vessel noise pollution as a human threat to fish: assessment of the stress response in gilthead sea bream (*Sparus aurata*, Linnaeus 1758). *Fish Physiology and Biochemistry* 42(2): 631-641. doi:10.1007/s10695-015-0165-3
- Chang, H.Y., Lin, T.H., Anraku, K. & Shao, Y.T. 2018. The Effects of Continuous Acoustic Stress on ROS Levels and Antioxidant-related Gene Expression in the Black Porgy (*Acanthopagrus schlegelii*). *Zoological Studies* 57. doi:10.6620/zs.2018.57-59

- Christel, I., Certain, G., Cama, A., Vieites, D.R. & Ferrer, X. 2013. Seabird aggregative patterns: A new tool for offshore wind energy risk assessment. *Marine Pollution Bulletin* 66(1-2): 84-91. doi:10.1016/j.marpolbul.2012.11.005
- Christie, N., Smyth, K., Barnes, R. & Elliott, M. 2014. Co-location of activities and designations: A means of solving or creating problems in marine spatial planning? *Marine Policy* 43: 254-261. doi:10.1016/j.marpol.2013.06.002
- Cleasby, I.R., Wakefield, E.D., Bearhop, S., Bodey, T.W., Votier, S.C. & Hamer, K.C. 2015. Three-dimensional tracking of a wide-ranging marine predator: flight heights and vulnerability to offshore wind farms. *Journal of Applied Ecology* 52(6): 1474-1482. doi:10.1111/1365-2664.12529
- Coleman, H.M., Kanat, G. & Turkdogan, F.I.A. 2009. Restoration of the Golden Horn Estuary (Halic). *Water Research* 43(20): 4989-5003. doi:10.1016/j.watres.2009.08.047
- Cominelli, S., Devillers, R., Yurk, H., MacGillivray, A., McWhinnie, L. & Canessa, R. 2018. Noise exposure from commercial shipping for the southern resident killer whale population. *Marine Pollution Bulletin* 136: 177-200. doi:10.1016/j.marpolbul.2018.08.050
- Cominelli, S., Leahy, M., Devillers, R. & Hall, G.B. 2019. Geovisualization tools to inform the management of vessel noise in support of species' conservation. *Ocean & Coastal Management* 169: 113-128. doi:10.1016/j.ocecoaman.2018.11.009
- Copping, A., Hanna, L., Van Cleve, B., Blake, K. & Anderson, R.M. 2015. Environmental Risk Evaluation System-an Approach to Ranking Risk of Ocean Energy Development on Coastal and Estuarine Environments. *Estuaries and Coasts* 38: S287-S302. doi:10.1007/s12237-014-9816-3
- Corbett, J.J. & Farrell, A. 2002. Mitigating air pollution impacts of passenger ferries. *Transportation Research Part D-Transport and Environment* 7(3): 197-211. doi:10.1016/s1361-9209(01)00019-0
- Culloch, R.M., Anderwald, P., Brandecker, A., Haberlin, D., McGovern, B., Pinfield, R., Visser, F., Jessopp, M. & Cronin, M. 2016. Effect of construction-related activities and vessel traffic on marine mammals. *Marine Ecology Progress Series* 549: 231-242. doi:10.3354/meps11686
- Dannheim, J., Bergstrom, L., Birchenough, S.N.R., Brzana, R., Boon, A.R., Coolen, J.W.P., Dauvin, J.C., De Mesel, I., Derweduwen, J., Gill, A.B., Hutchison, Z.L., Jackson, A.C., Janas, U., Martin, G., Raoux, A., Reubens, J., Rostin, L., Vanaverbeke, J., Wilding, T.A., Wilhelmsson, D. & Degraer, S. 2020. Benthic effects of offshore renewables: identification of knowledge gaps and urgently needed research. *Ices Journal of Marine Science* 77(3): 1092-1108. doi:10.1093/icesjms/fsz018
- De Clippele, L.H. & Risch, D. 2021. Measuring Sound at a Cold-Water Coral Reef to Assess the Impact of COVID-19 on Noise Pollution. *Frontiers in Marine Science* 8. doi:10.3389/fmars.2021.674702
- de Lima, R.L.P., Paxinou, K., Boogaard, F.C., Akkerman, O. & Lin, F.Y. 2021. In-Situ Water Quality Observations under a Large-Scale Floating Solar Farm Using Sensors and Underwater Drones. *Sustainability* 13(11). doi:10.3390/su13116421
- Debusschere, E., De Coensel, B., Bajek, A., Botteldooren, D., Hostens, K., Vanaverbeke, J., Vandendriessche, S., Van Ginderdeuren, K., Vincx, M. & Degraer, S. 2014. In Situ Mortality Experiments with Juvenile Sea Bass (*Dicentrarchus labrax*) in Relation to Impulsive Sound Levels Caused by Pile Driving of Windmill Foundations. *Plos One* 9(10). doi:10.1371/journal.pone.0109280

- Desholm, M. 2009. Avian sensitivity to mortality: Prioritising migratory bird species for assessment at proposed wind farms. *Journal of Environmental Management* 90(8): 2672-2679. doi:10.1016/j.jenvman.2009.02.005
- Didenkulova, I., Sheremet, A., Torsvik, T. & Soomere, T. 2013. Characteristic properties of different vessel wake signals. *Journal of Coastal Research*: 213-218. doi:10.2112/si65-037.1
- Dolman, S.J. & Jasny, M. 2015. Evolution of Marine Noise Pollution Management. *Aquatic Mammals* 41(4): 357-374. doi:10.1578/am.41.4.2015.357
- Dolman, S.J., Green, M., Gregerson, S. & Weir, C.R. 2016. Fulfilling EU Laws to Ensure Marine Mammal Protection During Marine Renewable Construction Operations in Scotland. I: Popper, A. N. & Hawkins, A. (red.) *Effects of Noise on Aquatic Life* li. S. 223-230. doi:10.1007/978-1-4939-2981-8_26
- Dähne, M., Gilles, A., Lucke, K., Peschko, V., Adler, S., Krugel, K., Sundermeyer, J. & Siebert, U. 2013. Effects of pile-driving on harbour porpoises (*Phocoena phocoena*) at the first offshore wind farm in Germany. *Environmental Research Letters* 8(2). doi:10.1088/1748-9326/8/2/025002
- Dähne, M., Tougaard, J., Carstensen, J., Rose, A. & Nabe-Nielsen, J. 2017. Bubble curtains attenuate noise from offshore wind farm construction and reduce temporary habitat loss for harbour porpoises. *Marine Ecology Progress Series* 580: 221-237. doi:10.3354/meps12257
- Eriksson, B.K., Sandstrom, A., Isaeus, M., Schreiber, H. & Karas, P. 2004. Effects of boating activities on aquatic vegetation in the Stockholm archipelago, Baltic Sea. *Estuarine Coastal and Shelf Science* 61(2): 339-349. doi:10.1016/j.ecss.2004.05.009
- Fang, Y.Y., Sung, P.J., Hu, W.C. & Chen, C.F. 2020. Underwater Noise Simulation of Impact Pile Driving for Offshore Wind Farm in Taiwan. *Journal of Theoretical and Computational Acoustics* 28(1). doi:10.1142/s2591728519500099
- Farfan, M.A., Duarte, J., Real, R., Munoz, A.R., Fa, J.E. & Vargas, J.M. 2017. Differential recovery of habitat use by birds after wind farm installation: A multi-year comparison. *Environmental Impact Assessment Review* 64: 8-15. doi:10.1016/j.eiar.2017.02.001
- Farr, H., Ruttenberg, B., Walter, R.K., Wang, Y.H. & White, C. 2021. Potential environmental effects of deepwater floating offshore wind energy facilities. *Ocean & Coastal Management* 207. doi:10.1016/j.ocecoaman.2021.105611
- Fernandez-Betelu, O., Graham, I.M., Brookes, K.L., Cheney, B.J., Barton, T.R. & Thompson, P.M. 2021. Far-Field Effects of Impulsive Noise on Coastal Bottlenose Dolphins. *Frontiers in Marine Science* 8. doi:10.3389/fmars.2021.664230
- Fey, D.P., Jakubowska, M., Greszkiewicz, M., Andrulewicz, E., Otremba, Z. & Urban-Malinga, B. 2019. Are magnetic and electromagnetic fields of anthropogenic origin potential threats to early life stages of fish? *Aquatic Toxicology* 209: 150-158. doi:10.1016/j.aquatox.2019.01.023
- Fijn, R.C., Krijgsveld, K.L., Poot, M.J.M. & Dirksen, S. 2015. Bird movements at rotor heights measured continuously with vertical radar at a Dutch offshore wind farm. *Ibis* 157(3): 558-566. doi:10.1111/ibi.12259
- Fowler, A.M., Jorgensen, A.M., Coolen, J.W.P., Jones, D.O.B., Svendsen, J.C., Brabant, R., Rumes, B. & Degraer, S. 2020. The ecology of infrastructure decommissioning in the North Sea: what we need to know and how to achieve it. *Ices Journal of Marine Science* 77(3): 1109-1126. doi:10.1093/icesjms/fsz143

- Fox, C.J., Benjamins, S., Masden, E.A. & Miller, R. 2018. Challenges and opportunities in monitoring the impacts of tidal-stream energy devices on marine vertebrates. *Renewable & Sustainable Energy Reviews* 81: 1926-1938. doi:10.1016/j.rser.2017.06.004
- Frades, J.L., Barba, J.G., Negro, V., Martin-Anton, M. & Soriano, J. 2020. Blue Economy: Compatibility between the Increasing Offshore Wind Technology and the Achievement of the SDG. *Journal of Coastal Research*: 1490-1494. doi:10.2112/si95-287.1
- Furness, R.W., Wade, H.M. & Masden, E.A. 2013. Assessing vulnerability of marine bird populations to offshore wind farms. *Journal of Environmental Management* 119: 56-66. doi:10.1016/j.jenvman.2013.01.025
- Furness, R.W., Garthe, S., Trinder, M., Matthiopoulos, J., Wanless, S. & Jeglinski, J. 2018. Nocturnal flight activity of northern gannets *Morus bassanus* and implications for modelling collision risk at offshore wind farms. *Environmental Impact Assessment Review* 73: 1-6. doi:10.1016/j.eiar.2018.06.006
- Gannier, A. & Marty, G. 2015. Sperm whales ability to avoid approaching vessels is affected by sound reception in stratified waters. *Marine Pollution Bulletin* 95(1): 283-288. doi:10.1016/j.marpolbul.2015.03.029
- Garcia-Soto, C. & Pingree, R.D. 2009. Spring and summer blooms of phytoplankton (SeaWiFS/MODIS) along a ferry line in the Bay of Biscay and western English Channel. *Continental Shelf Research* 29(8): 1111-1122. doi:10.1016/j.csr.2008.12.012
- Gasparatos, A., Doll, C.N.H., Esteban, M., Ahmed, A. & Olang, T.A. 2017. Renewable energy and biodiversity: Implications for transitioning to a Green Economy. *Renewable & Sustainable Energy Reviews* 70: 161-184. doi:10.1016/j.rser.2016.08.030
- Gervaise, C., Simard, Y., Roy, N., Kinda, B. & Menard, N. 2012. Shipping noise in whale habitat: Characteristics, sources, budget, and impact on belugas in Saguenay-St. Lawrence Marine Park hub. *Journal of the Acoustical Society of America* 132(1): 76-89. doi:10.1121/1.4728190
- Gill, A.B., Degraer, S., Lipsky, A., Mavraki, N., Methratta, E. & Brabant, R. 2020. Setting the context for offshore wind development effects on fish and fisheries. *Oceanography* 33(4): 118-127. doi:10.5670/oceanog.2020.411
- Gilles, A., Scheidat, M. & Siebert, U. 2009. Seasonal distribution of harbour porpoises and possible interference of offshore wind farms in the German North Sea. *Marine Ecology Progress Series* 383: 295-307. doi:10.3354/meps08020
- Goodale, M.W. & Milman, A. 2019. Assessing the cumulative exposure of wildlife to offshore wind energy development. *Journal of Environmental Management* 235: 77-83. doi:10.1016/j.jenvman.2019.01.022
- Goodale, M.W. & Milman, A. 2020. Assessing Cumulative Exposure of Northern Gannets to Offshore Wind Farms. *Wildlife Society Bulletin* 44(2): 252-259. doi:10.1002/wsb.1087
- Gordon, J., Blight, C., Bryant, E. & Thompson, D. 2019. Measuring responses of harbour seals to potential aversive acoustic mitigation signals using controlled exposure behavioural response studies. *Aquatic Conservation-Marine and Freshwater Ecosystems* 29: 157-177. doi:10.1002/aqc.3150
- Graham, I.M., Pirotta, E., Merchant, N.D., Farcas, A., Barton, T.R., Cheney, B., Hastie, G.D. & Thompson, P.M. 2017. Responses of bottlenose dolphins and harbor porpoises to impact and vibration piling noise during harbor construction. *Ecosphere* 8(5). doi:10.1002/ecs2.1793

- Grilli, A.R., Insua, T.L. & Spaulding, M. 2013. Protocol to Include Ecosystem Service Constraints in a Wind Farm Cost Model. *Journal of Environmental Engineering* 139(2): 176-186. doi:10.1061/(asce)ee.1943-7870.0000599
- Grilli, A.R. & Shumchenia, E.J. 2015. Toward wind farm monitoring optimization: assessment of ecological zones from marine landscapes using machine learning algorithms. *Hydrobiologia* 756(1): 117-137. doi:10.1007/s10750-014-2139-3
- Haelters, J., Duliere, V., Vigin, L. & Degraer, S. 2015. Towards a numerical model to simulate the observed displacement of harbour porpoises *Phocoena phocoena* due to pile driving in Belgian waters. *Hydrobiologia* 756(1): 105-116. doi:10.1007/s10750-014-2138-4
- Ham, G.S., Lahaye, E., Rosso, M., Moulins, A., Hines, E. & Tepsich, P. 2021. Predicting summer fin whale distribution in the Pelagos Sanctuary (north-western Mediterranean Sea) to identify dynamic whale-vessel collision risk areas. *Aquatic Conservation-Marine and Freshwater Ecosystems* 31(8): 2257-2277. doi:10.1002/aqc.3614
- Hammar, L., Wikstrom, A. & Molander, S. 2014. Assessing ecological risks of offshore wind power on Kattegat cod. *Renewable Energy* 66: 414-424. doi:10.1016/j.renene.2013.12.024
- Hastie, G.D., Russell, D.J.F., McConnell, B., Moss, S., Thompson, D. & Janik, V.M. 2015. Sound exposure in harbour seals during the installation of an offshore wind farm: predictions of auditory damage. *Journal of Applied Ecology* 52(3): 631-640. doi:10.1111/1365-2664.12403
- Hawkins, A.D., Pembroke, A.E. & Popper, A.N. 2015. Information gaps in understanding the effects of noise on fishes and invertebrates. *Reviews in Fish Biology and Fisheries* 25(1): 39-64. doi:10.1007/s11160-014-9369-3
- Hawkins, A.D., Hazelwood, R.A., Popper, A.N. & Macey, P.C. 2021. Substrate vibrations and their potential effects upon fishes and invertebrates. *Journal of the Acoustical Society of America* 149(4): 2782-2790. doi:10.1121/10.0004773
- He, W., Zou, C., Pang, Y.T. & Wang, X.M. 2021. Environmental noise and vibration characteristics of rubber-spring floating slab track. *Environmental Science and Pollution Research* 28(11): 13671-13689. doi:10.1007/s11356-020-11627-w
- Heery, E.C., Bishop, M.J., Critchley, L.P., Bugnot, A.B., Airoidi, L., Mayer-Pinto, M., Sheehan, E.V., Coleman, R.A., Loke, L.H.L., Johnston, E.L., Komyakova, V., Morris, R.L., Strain, E.M.A., Naylor, L.A. & Dafforn, K.A. 2017. Identifying the consequences of ocean sprawl for sedimentary habitats. *Journal of Experimental Marine Biology and Ecology* 492: 31-48. doi:10.1016/j.jembe.2017.01.020
- Heinänen, S., Zydalis, R., Kleinschmidt, B., Dorsch, M., Burger, C., Morkunas, J., Quillfeldt, P. & Nehls, G. 2020. Satellite telemetry and digital aerial surveys show strong displacement of red-throated divers (*Gavia stellata*) from offshore wind farms. *Marine Environmental Research* 160. doi:10.1016/j.marenvres.2020.104989
- Holtberget, S.H., Xiang, X., Dorum, C., Veie, J. & Minoretti, A. 2019. The Choice of Materials for the E39 Fjord Crossing Project. *Nordic Concrete Research* 61(2): 39-51. doi:10.2478/ncr-2019-0016
- Hooper, T. & Austen, M. 2014. The co-location of offshore windfarms and decapod fisheries in the UK: Constraints and opportunities. *Marine Policy* 43: 295-300. doi:10.1016/j.marpol.2013.06.011
- Hutchison, Z.L., Gill, A.B., Sigray, P., He, H.B. & King, J.W. 2020. Anthropogenic electromagnetic fields (EMF) influence the behaviour of bottom-dwelling marine species. *Scientific Reports* 10(1). doi:10.1038/s41598-020-60793-x

- Inger, R., Attrill, M.J., Bearhop, S., Broderick, A.C., Grecian, W.J., Hodgson, D.J., Mills, C., Sheehan, E., Votier, S.C., Witt, M.J. & Godley, B.J. 2009. Marine renewable energy: potential benefits to biodiversity? An urgent call for research. *Journal of Applied Ecology* 46(6): 1145-1153. doi:10.1111/j.1365-2664.2009.01697.x
- Jeong, Y., Min, S. & Paeng, D.G. 2020. Moored measurement of the ambient noise and analysis with environmental factors in the coastal sea of Jeju Island. *Journal of the Acoustical Society of Korea* 39(5): 390-399. doi:10.7776/ask.2020.39.5.390
- Johansen, P.O., Isaksen, T.E., Bye-Ingebrigtsen, E., Haave, M., Dahlgren, T.G., Kvalo, S.E., Greenacre, M., Durand, D. & Rapp, H.T. 2018. Temporal changes in benthic macrofauna on the west coast of Norway resulting from human activities. *Marine Pollution Bulletin* 128: 483-495. doi:10.1016/j.marpolbul.2018.01.063
- Jones, I.T., Peyla, J.F., Clark, H., Song, Z.C., Stanley, J.A. & Mooney, T.A. 2021. Changes in feeding behavior of longfin squid (*Doryteuthis pealeii*) during laboratory exposure to pile driving noise. *Marine Environmental Research* 165. doi:10.1016/j.marenvres.2020.105250
- Kaldellis, J.K., Apostolou, D., Kapsali, M. & Kondili, E. 2016. Environmental and social footprint of offshore wind energy. Comparison with onshore counterpart. *Renewable Energy* 92: 543-556. doi:10.1016/j.renene.2016.02.018
- Karydis, M. 2013. Public attitudes and environmental impacts of wind farms: A review. *Global Nest Journal* 15(4): 585-604.
- Kastelein, R.A., van Heerden, D., Gransier, R. & Hoek, L. 2013. Behavioral responses of a harbor porpoise (*Phocoena phocoena*) to playbacks of broadband pile driving sounds. *Marine Environmental Research* 92: 206-214. doi:10.1016/j.marenvres.2013.09.020
- Kastelein, R.A., Huybrechts, J., Covi, J. & Helder-Hoek, L. 2017. Behavioral Responses of a Harbor Porpoise (*Phocoena phocoena*) to Sounds from an Acoustic Porpoise Deterrent. *Aquatic Mammals* 43(3): 233-244. doi:10.1578/am.43.3.2017.233
- Kastelein, R.A., Helder-Hoek, L., Kommeren, A., Covi, J. & Gransier, R. 2018a. Effect of pile-driving sounds on harbor seal (*Phoca vitulina*) hearing. *Journal of the Acoustical Society of America* 143(6): 3583-3594. doi:10.1121/1.5040493
- Kastelein, R.A., Van de Voorde, S. & Jennings, N. 2018b. Swimming Speed of a Harbor Porpoise (*Phocoena phocoena*) During Playbacks of Offshore Pile Driving Sounds. *Aquatic Mammals* 44(1): 92-99. doi:10.1578/am.44.1.2018.92
- Kastelein, R.A., Helder-Hoek, L., Booth, C., Jennings, N. & Leopold, M. 2019a. High Levels of Food Intake in Harbor Porpoises (*Phocoena phocoena*): Insight into Recovery from Disturbance. *Aquatic Mammals* 45(4): 380-388. doi:10.1578/am.45.4.2019.380
- Kastelein, R.A., Huijser, L.A.E., Cornelisse, S., Helder-Hoek, L., Jennings, N. & de Jong, C.A.F. 2019b. Effect of Pile-Driving Playback Sound Level on Fish-Catching Efficiency in Harbor Porpoises (*Phocoena phocoena*). *Aquatic Mammals* 45(4): 398-410. doi:10.1578/am.45.4.2019.398
- Khangaonkar, T., Nugraha, A. & Wang, T.P. 2018. Hydrodynamic Zone of Influence Due to a Floating Structure in a Fjordal Estuary-Hood Canal Bridge Impact Assessment. *Journal of Marine Science and Engineering* 6(4). doi:10.3390/jmse6040119
- Kitazawa, D., Tabeta, S., Fujino, M. & Kato, T. 2010. Assessment of environmental variations caused by a very large floating structure in a semi-closed bay. *Environmental Monitoring and Assessment* 165(1-4): 461-474. doi:10.1007/s10661-009-0959-9

- Kok, A.C.M., Bruil, L., Berges, B., Sakinan, S., Debusschere, E., Reubens, J., de Haan, D., Norro, A. & Slabbekoorn, H. 2021. An echosounder view on the potential effects of impulsive noise pollution on pelagic fish around windfarms in the North Sea. *Environmental Pollution* 290. doi:10.1016/j.envpol.2021.118063
- Koschinski, S., Culik, B.M., Henriksen, O.D., Tregenza, N., Ellis, G., Jansen, C. & Kathe, G. 2003. Behavioural reactions of free-ranging porpoises and seals to the noise of a simulated 2 MW windpower generator. *Marine Ecology Progress Series* 265: 263-273. doi:10.3354/meps265263
- Kraus, C. & Carter, L. 2018. Seabed recovery following protective burial of subsea cables - Observations from the continental margin. *Ocean Engineering* 157: 251-261. doi:10.1016/j.oceaneng.2018.03.037
- Krone, R., Gutow, L., Brey, T., Dannheim, J. & Schroder, A. 2013. Mobile demersal megafauna at artificial structures in the German Bight - Likely effects of offshore wind farm development. *Estuarine Coastal and Shelf Science* 125: 1-9. doi:10.1016/j.ecss.2013.03.012
- Kumar, J.C.R., Kumar, D.V., Baskar, D., Arunsi, B.M., Jenova, R. & Majid, M.A. 2021. Offshore wind energy status, challenges, opportunities, environmental impacts, occupational health, and safety management in India. *Energy & Environment* 32(4): 565-603. doi:10.1177/0958305x20946483
- Langston, R.H.W. 2013. Birds and Wind Projects Across the Pond: A UK Perspective. *Wildlife Society Bulletin* 37(1): 5-18. doi:10.1002/wsb.262
- Lesage, V., Barrette, C., Kingsley, M.C.S. & Sjare, B. 1999. The effect of vessel noise on the vocal behavior of Belugas in the St. Lawrence River estuary, Canada. *Marine Mammal Science* 15(1): 65-84. doi:10.1111/j.1748-7692.1999.tb00782.x
- Leung, D.Y.C. & Yang, Y. 2012. Wind energy development and its environmental impact: A review. *Renewable & Sustainable Energy Reviews* 16(1): 1031-1039. doi:10.1016/j.rser.2011.09.024
- Leunissen, E.M. & Dawson, S.M. 2018. Underwater noise levels of pile-driving in a New Zealand harbour, and the potential impacts on endangered Hector's dolphins. *Marine Pollution Bulletin* 135: 195-204. doi:10.1016/j.marpolbul.2018.07.024
- Lin, H., Xiang, Y.Q., Yang, Y. & Chen, Z.Y. 2018. Dynamic response analysis for submerged floating tunnel due to fluid-vehicle-tunnel interaction. *Ocean Engineering* 166: 290-301. doi:10.1016/j.oceaneng.2018.08.023
- Lin, M., Wang, R.K. & Lin, W. 2020. Artificial Island Construction Using Large Steel Cylinders. *Structural Engineering International* 30(4): 484-492. doi:10.1080/10168664.2020.1726255
- Lin, Y.T., Newhall, A.E., Miller, J.H., Potty, G.R. & Vigness-Raposa, K.J. 2019. A three-dimensional underwater sound propagation model for offshore wind farm noise prediction. *Journal of the Acoustical Society of America* 145(5): EL335-EL340. doi:10.1121/1.5099560
- Lindeboom, H.J., Kouwenhoven, H.J., Bergman, M.J.N., Bouma, S., Brasseur, S., Daan, R., Fijn, R.C., de Haan, D., Dirksen, S., van Hal, R., Lambers, R.H.R., Ter Hofstede, R., Krijgsveld, K.L., Leopold, M. & Scheidat, M. 2011. Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. *Environmental Research Letters* 6(3). doi:10.1088/1748-9326/6/3/035101
- Linder, H.L., Horne, J.K. & Ward, E.J. 2017. Modeling baseline conditions of ecological indicators: Marine renewable energy environmental monitoring. *Ecological Indicators* 83: 178-191. doi:10.1016/j.ecolind.2017.07.015

- Luis, A.R., Couchinho, M.N. & dos Santos, M.E. 2014. Changes in the acoustic behavior of resident bottlenose dolphins near operating vessels. *Marine Mammal Science* 30(4): 1417-1426. doi:10.1111/mms.12125
- Madsen, P.T., Wahlberg, M., Tougaard, J., Lucke, K. & Tyack, P. 2006. Wind turbine underwater noise and marine mammals: implications of current knowledge and data needs. *Marine Ecology Progress Series* 309: 279-295. doi:10.3354/meps309279
- Malmgren, E., Brynolf, S., Fridell, E., Grahn, M. & Andersson, K. 2021. The environmental performance of a fossil-free ship propulsion system with onboard carbon capture - a life cycle assessment of the HyMethShip concept. *Sustainable Energy & Fuels* 5(10): 2753-2770. doi:10.1039/d1se00105a
- Mann, J. & Teilmann, J. 2013. Environmental impact of wind energy. *Environmental Research Letters* 8(3). doi:10.1088/1748-9326/8/3/035001
- Margaritou, M.D. & Tzannatos, E. 2018. A Multi-Criteria Optimization Approach for Solar Energy and Wind Power echnologies in Shipping. *Fme Transactions* 46(3): 374-380. doi:10.5937/fmet1803374M
- Martin, S.B. & Barclay, D.R. 2019. Determining the dependence of marine pile driving sound levels on strike energy, pile penetration, and propagation effects using a linear mixed model based on damped cylindrical spreading. *Journal of the Acoustical Society of America* 146(1): 109-121. doi:10.1121/1.5114797
- Masden, E.A., Haydon, D.T., Fox, A.D. & Furness, R.W. 2010. Barriers to movement: Modelling energetic costs of avoiding marine wind farms amongst breeding seabirds. *Marine Pollution Bulletin* 60(7): 1085-1091. doi:10.1016/j.marpolbul.2010.01.016
- Masden, E.A., Reeve, R., Desholm, M., Fox, A.D., Furness, R.W. & Haydon, D.T. 2012. Assessing the impact of marine wind farms on birds through movement modelling. *Journal of the Royal Society Interface* 9(74): 2120-2130. doi:10.1098/rsif.2012.0121
- Matabos, M., Aguzzi, J., Robert, K., Costa, C., Menesatti, P., Company, J.B. & Juniper, S.K. 2011. Multi-parametric study of behavioural modulation in demersal decapods at the VENUS cabled observatory in Saanich Inlet, British Columbia, Canada. *Journal of Experimental Marine Biology and Ecology* 401(1-2): 89-96. doi:10.1016/j.jembe.2011.02.041
- May, R., Nygard, T., Rd, N. & Bevanger, K. 2013. Habitat Utilization in White-Tailed Eagles (*Haliaeetus albicilla*) and the Displacement Impact of the Smola Wind-Power Plant. *Wildlife Society Bulletin* 37(1): 75-83. doi:10.1002/wsb.264
- May, R., Astrom, J., Hamre, O. & Dahl, E.L. 2017. Do birds in flight respond to (ultra)violet lighting? *Avian Research* 8. doi:10.1186/s40657-017-0092-3
- May, R., Nygard, T., Falkdalen, U., Astrom, J., Hamre, O. & Stokke, B.G. 2020. Paint it black: Efficacy of increased wind turbine rotor blade visibility to reduce avian fatalities. *Ecology and Evolution* 10(16): 8927-8935. doi:10.1002/ece3.6592
- May, R., Jackson, C.R., Middel, H., Stokke, B.G. & Verones, F. 2021. Life-cycle impacts of wind energy development on bird diversity in Norway. *Environmental Impact Assessment Review* 90. doi:10.1016/j.eiar.2021.106635
- McWhinnie, L.H., O'Hara, P.D., Hilliard, C., Le Baron, N., Smallshaw, L., Pelot, R. & Canessa, R. 2021. Assessing vessel traffic in the Salish Sea using satellite AIS: An important contribution for planning, management and conservation in southern resident killer whale critical habitat. *Ocean & Coastal Management* 200. doi:10.1016/j.ocecoaman.2020.105479

- Methratta, E.T. & Dardick, W.R. 2019. Meta-Analysis of Finfish Abundance at Offshore Wind Farms. *Reviews in Fisheries Science & Aquaculture* 27(2): 242-260. doi:10.1080/23308249.2019.1584601
- Miles, J., Martin, T. & Goddard, L. 2017. Current and wave effects around windfarm monopile foundations. *Coastal Engineering* 121: 167-178. doi:10.1016/j.coastaleng.2017.01.003
- Miller, J.H., Potty, G.R., Vigness-Raposa, K.J., Casagrande, D., Miller, L.A., Nystuen, J.A., Scheifele, P.M. & Clark, J.G. 2012. Environmental Assessment of Offshore Wind Power Generation: Effect on a Noise Budget. I: Popper, A. N. & Hawkins, A. (red.) *Effects of Noise on Aquatic Life*. S. 519-522. doi:10.1007/978-1-4419-7311-5_118
- Millidine, K.J., Malcolm, I.A., McCartney, A., Laughton, R., Gibbins, C.N. & Fryer, R.J. 2015. The influence of wind farm development on the hydrochemistry and ecology of an upland stream. *Environmental Monitoring and Assessment* 187(8). doi:10.1007/s10661-015-4750-9
- Mitamura, H., Nishizawa, H., Mitsunaga, Y., Tanaka, K., Takagi, J., Noda, T., Tsujimura, H., Omi, H., Sakurai, R., Sato, M., Arai, N. & Hori, M. Attraction of an artificial reef: a migratory demersal flounder remains in shallow water under high temperature conditions in summer. *Environmental Biology of Fishes*. doi:10.1007/s10641-021-01153-0
- Mooney, T.A., Andersson, M.H. & Stanley, J. 2020. Acoustic impacts of offshore wind energy on fishery resources. *An Evolving Source and Varied Effects Across a Wind Farm's Lifetime*. *Oceanography* 33(4): 82-95. doi:10.5670/oceanog.2020.408
- Moore, M., Berejikian, B.A. & Tezak, E.P. 2013. A Floating Bridge Disrupts Seaward Migration and Increases Mortality of Steelhead Smolts in Hood Canal, Washington State. *Plos One* 8(9). doi:10.1371/journal.pone.0073427
- Morrisey, D.J., Cole, R.G., Davey, N.K., Handley, S.J., Bradley, A., Brown, S.N. & Madarasz, A.L. 2006. Abundance and diversity of fish on mussel farms in New Zealand. *Aquaculture* 252(2-4): 277-288. doi:10.1016/j.aquaculture.2005.06.047
- Munehara, H., Tanaka, Y. & Futamura, T. 2009. Novel sledge net system employing propulsion vehicles for sampling demersal organisms on sandy bottoms. *Estuarine Coastal and Shelf Science* 83(3): 371-377. doi:10.1016/j.ecss.2009.04.004
- Nabe-Nielsen, J., Sibly, R.M., Tougaard, J., Teilmann, J. & Sveegaard, S. 2014. Effects of noise and by-catch on a Danish harbour porpoise population. *Ecological Modelling* 272: 242-251. doi:10.1016/j.ecolmodel.2013.09.025
- Nabe-Nielsen, J., van Beest, F.M., Grimm, V., Sibly, R.M., Teilmann, J. & Thompson, P.M. 2018. Predicting the impacts of anthropogenic disturbances on marine populations. *Conservation Letters* 11(5). doi:10.1111/conl.12563
- Narita, H. 1995. Coastal marine transportation and floating structures. *Marine Technology Society Journal* 29(3): 50-57.
- Nastasi, M., Fredianelli, L., Bernardini, M., Teti, L., Fidecaro, F. & Licitra, G. 2020. Parameters Affecting Noise Emitted by Ships Moving in Port Areas. *Sustainability* 12(20). doi:10.3390/su12208742
- Nehls, G., Rose, A., Diederichs, A., Bellmann, M. & Pehlke, H. 2016. Noise Mitigation During Pile Driving Efficiently Reduces Disturbance of Marine Mammals. I: Popper, A. N. & Hawkins, A. (red.) *Effects of Noise on Aquatic Life* Ii. S. 755-762. doi:10.1007/978-1-4939-2981-8_92
- Neuman, D.G., Tapio, E., Haggard, D., Laws, K.E. & Bland, R.W. 2001. Observation of long waves generated by ferries. *Canadian Journal of Remote Sensing* 27(4): 361-370. doi:10.1080/07038992.2001.10854878

- Nielsen, L.P. 2016. Ecology: Electrical Cable Bacteria Save Marine Life. *Current Biology* 26(1): R32-R33. doi:10.1016/j.cub.2015.11.014
- Ohman, M.C., Sigraý, P. & Westerberg, H. 2007. Offshore windmills and the effects electromagnetic fields an fish. *Ambio* 36(8): 630-633. doi:10.1579/0044-7447(2007)36[630:Owateo]2.0.Co;2
- Papathanasopoulou, E., Beaumont, N., Hooper, T., Nunes, J. & Queiros, A.M. 2015. Energy systems and their impacts on marine ecosystem services. *Renewable & Sustainable Energy Reviews* 52: 917-926. doi:10.1016/j.rser.2015.07.150
- Parsons, M.J.G., Duncan, A.J., Parsons, S.K. & Erbe, C. 2020. Reducing vessel noise: An example of a solar-electric passenger ferrya). *Journal of the Acoustical Society of America* 147(5): 3575-3583. doi:10.1121/10.0001264
- Pearce-Higgins, J.W., Stephen, L., Langston, R.H.W., Bainbridge, I.P. & Bullman, R. 2009. The distribution of breeding birds around upland wind farms. *Journal of Applied Ecology* 46(6): 1323-1331. doi:10.1111/j.1365-2664.2009.01715.x
- Perkol-Finkel, S., Zilman, G., Sella, I., Miloh, T. & Benayahu, Y. 2006. Floating and fixed artificial habitats: effects of substratum motion on benthic communities in a coral reef environment. *Marine Ecology Progress Series* 317: 9-20. doi:10.3354/meps317009
- Perrow, M.R., Skeate, E.R., Lines, P., Brown, D. & Tomlinson, M.L. 2006. Radio telemetry as a tool for impact assessment of wind farms: the case of Little Terns *Sterna albifrons* at Scroby Sands, Norfolk, UK. *Ibis* 148: 57-75. doi:10.1111/j.1474-919X.2006.00508.x
- Perrow, M.R., Gilroy, J.J., Skeate, E.R. & Tomlinson, M.L. 2011. Effects of the construction of Scroby Sands offshore wind farm on the prey base of Little tern *Sternula albifrons* at its most important UK colony. *Marine Pollution Bulletin* 62(8): 1661-1670. doi:10.1016/j.marpolbul.2011.06.010
- Peschko, V., Mendel, B., Muller, S., Markones, N., Mercker, M. & Garthe, S. 2020. Effects of offshore windfarms on seabird abundance: Strong effects in spring and in the breeding season. *Marine Environmental Research* 162. doi:10.1016/j.marenvres.2020.105157
- Peschko, V., Mendel, B., Mercker, M., Dierschke, J. & Garthe, S. 2021. Northern gannets (*Morus bassanus*) are strongly affected by operating offshore wind farms during the breeding season. *Journal of Environmental Management* 279. doi:10.1016/j.jenvman.2020.111509
- Pezy, J.P., Raoux, A. & Dauvin, J.C. 2020. The environmental impact from an offshore windfarm: Challenge and evaluation methodology based on an ecosystem approach. *Ecological Indicators* 114. doi:10.1016/j.ecolind.2020.106302
- Picciulin, M., Sebastianutto, L., Codarin, A., Farina, A. & Ferrero, E.A. 2010. In situ behavioural responses to boat noise exposure of *Gobius cruentatus* (Gmelin, 1789; fam. Gobiidae) and *Chromis chromis* (Linnaeus, 1758; fam. Pomacentridae) living in a Marine Protected Area. *Journal of Experimental Marine Biology and Ecology* 386(1-2): 125-132. doi:10.1016/j.jembe.2010.02.012
- Pinn, E.H., Macleod, K. & Tasker, M.L. 2021. Conservation of transnational species: The tensions between legal requirements and best scientific evidence. *Aquatic Conservation-Marine and Freshwater Ecosystems*. doi:10.1002/aqc.3693
- Pizzol, M. 2019. Deterministic and stochastic carbon footprint of intermodal ferry and truck freight transport across Scandinavian routes. *Journal of Cleaner Production* 224: 626-636. doi:10.1016/j.jclepro.2019.03.270

- Pollock, C.J., Lane, J.V., Buckingham, L., Garthe, S., Jeavons, R., Furness, R.W. & Hamer, K.C. 2021. Risks to different populations and age classes of gannets from impacts of offshore wind farms in the southern North Sea. *Marine Environmental Research* 171. doi:10.1016/j.marenvres.2021.105457
- Popper, A.N. & Hawkins, A.D. 2019. An overview of fish bioacoustics and the impacts of anthropogenic sounds on fishes. *Journal of Fish Biology* 94(5): 692-713. doi:10.1111/jfb.13948
- Pradhan, S., Kumar, S., Mohanty, S. & Nayak, S.K. 2019. Environmentally Benign Fouling-Resistant Marine Coatings: A Review. *Polymer-Plastics Technology and Materials* 58(5): 498-518. doi:10.1080/03602559.2018.1482922
- Reeder, D.B., Joseph, J.E. & Rago, T.A. 2020. Underwater sound generated by motor vehicle traffic in an underwater tunnel. *Journal of the Acoustical Society of America* 148(3): EL215-EL220. doi:10.1121/10.0001805
- Reubens, J.T., Braeckman, U., Vanaverbeke, J., Van Colen, C., Degraer, S. & Vincx, M. 2013. Aggregation at windmill artificial reefs: CPUE of Atlantic cod (*Gadus morhua*) and pouting (*Trisopterus luscus*) at different habitats in the Belgian part of the North Sea. *Fisheries Research* 139: 28-34. doi:10.1016/j.fishres.2012.10.011
- Rossington, K., Benson, T., Lepper, P. & Jones, D. 2013. Eco-hydro-acoustic modeling and its use as an EIA tool. *Marine Pollution Bulletin* 75(1-2): 235-243. doi:10.1016/j.marpolbul.2013.07.024
- Rotllant, G., Aguzzi, J., Sarria, D., Gisbert, E., Sbragaglia, V., Del Rio, J., Simeo, C.G., Manuel, A., Molino, E., Costa, C. & Sarda, F. 2015. Pilot acoustic tracking study on adult spiny lobsters (*Palinurus mauritanicus*) and spider crabs (*Maja squinado*) within an artificial reef. *Hydrobiologia* 742(1): 27-38. doi:10.1007/s10750-014-1959-5
- Russell, D.J.F., Hastie, G.D., Thompson, D., Janik, V.M., Hammond, P.S., Scott-Hayward, L.A.S., Matthiopoulos, J., Jones, E.L. & McConnell, B.J. 2016. Avoidance of wind farms by harbour seals is limited to pile driving activities. *Journal of Applied Ecology* 53(6): 1642-1652. doi:10.1111/1365-2664.12678
- Saidur, R., Rahim, N.A., Islam, M.R. & Solangi, K.H. 2011. Environmental impact of wind energy. *Renewable & Sustainable Energy Reviews* 15(5): 2423-2430. doi:10.1016/j.rser.2011.02.024
- Sara, G., Dean, J.M., D'Amato, D., Buscaino, G., Oliveri, A., Genovese, S., Ferro, S., Buffa, G., Lo Martire, M. & Mazzola, S. 2007. Effect of boat noise on the behaviour of bluefin tuna *Thunnus thynnus* in the Mediterranean Sea. *Marine Ecology Progress Series* 331: 243-253. doi:10.3354/meps331243
- Schaffeld, T., Schnitzler, J.G., Ruser, A., Woelfing, B., Baltzer, J. & Siebert, U. 2020. Effects of multiple exposures to pile driving noise on harbor porpoise hearing during simulated flights- An evaluation tool. *Journal of the Acoustical Society of America* 147(2): 685-697. doi:10.1121/10.0000595
- Scheidat, M., Tougaard, J., Brasseur, S., Carstensen, J., Petel, T.V., Teilmann, J. & Reijnders, P. 2011. Harbour porpoises (*Phocoena phocoena*) and wind farms: a case study in the Dutch North Sea. *Environmental Research Letters* 6(2). doi:10.1088/1748-9326/6/2/025102
- Seike, K., Sassa, S., Shirai, K. & Kubota, K. 2018. Lasting Impact of a Tsunami Event on Sediment-Organism Interactions in the Ocean. *Journal of Geophysical Research-Oceans* 123(2): 1376-1392. doi:10.1002/2017jc013746

- Skeate, E.R., Perrow, M.R. & Gilroy, J.J. 2012. Likely effects of construction of Scroby Sands offshore wind farm on a mixed population of harbour Phoca vitulina and grey Halichoerus grypus seals. *Marine Pollution Bulletin* 64(4): 872-881. doi:10.1016/j.marpolbul.2012.01.029
- Slavik, K., Lemmen, C., Zhang, W.Y., Kerimoglu, O., Klingbeil, K. & Wirtz, K.W. 2019. The large-scale impact of offshore wind farm structures on pelagic primary productivity in the southern North Sea. *Hydrobiologia* 845(1): 35-53. doi:10.1007/s10750-018-3653-5
- Solan, M., Hauton, C., Godbold, J.A., Wood, C.L., Leighton, T.G. & White, P. 2016. Anthropogenic sources of underwater sound can modify how sediment-dwelling invertebrates mediate ecosystem properties. *Scientific Reports* 6. doi:10.1038/srep20540
- Song, X.D., Zhang, X.G., Xiong, W., Guo, Z.M. & Wang, B. 2020. Experimental and numerical study on underwater noise radiation from an underwater tunnel. *Environmental Pollution* 267. doi:10.1016/j.envpol.2020.115536
- Stober, U. & Thomsen, F. 2019. Effect of impact pile driving noise on marine mammals: A comparison of different noise exposure criteria. *Journal of the Acoustical Society of America* 145(5): 3252-3259. doi:10.1121/1.5109387
- Stokke, B.G., Nygard, T., Falkdalen, U., Pedersen, H.C. & May, R. 2020. Effect of tower base painting on willow ptarmigan collision rates with wind turbines. *Ecology and Evolution* 10(12): 5670-5679. doi:10.1002/ece3.6307
- Terhune, J.M. & Killorn, D. 2021. A Method for Preliminary Assessment of the Masking Potential of Anthropogenic Noise to Baleen Whale Calls. *Aquatic Mammals* 47(3): 283-291. doi:10.1578/am.47.3.2021.283
- Thompson, P.M., Lusseau, D., Barton, T., Simmons, D., Rusin, J. & Bailey, H. 2010. Assessing the responses of coastal cetaceans to the construction of offshore wind turbines. *Marine Pollution Bulletin* 60(8): 1200-1208. doi:10.1016/j.marpolbul.2010.03.030
- Thompson, P.M., Hastie, G.D., Nedwell, J., Barham, R., Brookes, K.L., Cordes, L.S., Bailey, H. & McLean, N. 2013. Framework for assessing impacts of pile-driving noise from offshore wind farm construction on a harbour seal population. *Environmental Impact Assessment Review* 43: 73-85. doi:10.1016/j.eiar.2013.06.005
- Tougaard, J., Henriksen, O.D. & Miller, L.A. 2009. Underwater noise from three types of offshore wind turbines: Estimation of impact zones for harbor porpoises and harbor seals. *Journal of the Acoustical Society of America* 125(6): 3766-3773. doi:10.1121/1.3117444
- Tougaard, J., Hermannsen, L. & Madsen, P.T. 2020. How loud is the underwater noise from operating offshore wind turbines? *Journal of the Acoustical Society of America* 148(5): 2885-2893. doi:10.1121/10.0002453
- Tsouvalas, A. 2020. Underwater Noise Emission Due to Offshore Pile Installation: A Review. *Energies* 13(12). doi:10.3390/en13123037
- Vaissiere, A.C., Levrel, H., Pioch, S. & Carlier, A. 2014. Biodiversity offsets for offshore wind farm projects: The current situation in Europe. *Marine Policy* 48: 172-183. doi:10.1016/j.marpol.2014.03.023
- Valsecchi, E., Arcangeli, A., Lombardi, R., Boyse, E., Carr, I.M., Galli, P. & Goodman, S.J. 2021. Ferries and Environmental DNA: Underway Sampling From Commercial Vessels Provides New Opportunities for Systematic Genetic Surveys of Marine Biodiversity. *Frontiers in Marine Science* 8. doi:10.3389/fmars.2021.704786

- van Berkel, J., Burchard, H., Christensen, A., Mortensen, L.O., Petersen, O.S. & Thomsen, F. 2020. The Effects of Offshore Wind Farms on Hydrodynamics and Implications for Fishes. *Oceanography* 33(4): 108-117. doi:10.5670/oceanog.2020.410
- van der Molen, J., Smith, H.C.M., Lepper, P., Limpenny, S. & Rees, J. 2014. Predicting the large-scale consequences of offshore wind turbine array development on a North Sea ecosystem. *Continental Shelf Research* 85: 60-72. doi:10.1016/j.csr.2014.05.018
- Van Renterghem, T., Botteldooren, D. & Dekoninck, L. 2014. Airborne sound propagation over sea during offshore wind farm piling. *Journal of the Acoustical Society of America* 135(2): 599-609. doi:10.1121/1.4861244
- Verfuss, U.K., Sparling, C.E., Arnot, C., Judd, A. & Coyle, M. 2016. Review of Offshore Wind Farm Impact Monitoring and Mitigation with Regard to Marine Mammals. I: Popper, A. N. & Hawkins, A. (red.) *Effects of Noise on Aquatic Life* li. S. 1175-1182. doi:10.1007/978-1-4939-2981-8_147
- Wahlberg, M. & Westerberg, H. 2005. Hearing in fish and their reactions to sounds from offshore wind farms. *Marine Ecology Progress Series* 288: 295-309. doi:10.3354/meps288295
- Wang, T., Yu, W.W., Zou, X.Q., Zhang, D.J., Li, B.J., Wang, J.J. & Zhang, H. 2018a. Zooplankton Community Responses and the Relation to Environmental Factors from Established Offshore Wind Farms within the Rudong Coastal Area of China. *Journal of Coastal Research* 34(4): 843-855. doi:10.2112/jcoastres-d-17-00058.1
- Wang, T., Zou, X.Q., Li, B.J., Yao, Y.L., Li, J.S., Hui, H.J., Yu, W.W. & Wang, C.L. 2018b. Microplastics in a wind farm area: A case study at the Rudong Offshore Wind Farm, Yellow Sea, China. *Marine Pollution Bulletin* 128: 466-474. doi:10.1016/j.marpolbul.2018.01.050
- Whyte, K.F., Russell, D.J.F., Sparling, C.E., Binnerts, B. & Hastie, G.D. 2020. Estimating the effects of pile driving sounds on seals: Pitfalls and possibilities. *Journal of the Acoustical Society of America* 147(6): 3948-3958. doi:10.1121/10.0001408
- Wilber, D.H., Carey, D.A. & Griffin, M. 2018. Flatfish habitat use near North America's first offshore wind farm. *Journal of Sea Research* 139: 24-32. doi:10.1016/j.seares.2018.06.004
- Wilhelmsson, D., Yahya, S.A.S. & Ohman, M.C. 2006. Effects of high-relief structures on cold temperate fish assemblages: A field experiment. *Marine Biology Research* 2(2): 136-147. doi:10.1080/17451000600684359
- Wilhelmsson, D. & Malm, T. 2008. Fouling assemblages on offshore wind power plants and adjacent substrata. *Estuarine Coastal and Shelf Science* 79(3): 459-466. doi:10.1016/j.ecss.2008.04.020
- Wilkens, S.L., Stanley, J.A. & Jeffs, A.G. 2012. Induction of settlement in mussel (*Perna canaliculus*) larvae by vessel noise. *Biofouling* 28(1): 65-72. doi:10.1080/08927014.2011.651717
- Wilson, J.C., Elliott, M., Cutts, N.D., Mander, L., Mendao, V., Perez-Dominguez, R. & Phelps, A. 2010. Coastal and Offshore Wind Energy Generation: Is It Environmentally Benign? *Energies* 3(7): 1383-1422. doi:10.3390/en3071383
- Wood, M.P. & Carter, L. 2008. Whale Entanglements With Submarine Telecommunication Cables. *IEEE Journal of Oceanic Engineering* 33(4): 445-450. doi:10.1109/joe.2008.2001638
- Zagubien, A. 2017. The Results of the Measurements and Analyses of Impact of Wind Farms on Acoustic Climate. *Rocznik Ochrona Srodowiska* 19: 527-539.

The Norwegian Institute for Nature Research, NINA, is as an independent foundation focusing on environmental research, emphasizing the interaction between human society, natural resources and biodiversity.

NINA was established in 1988. The headquarters are located in Trondheim, with branches in Tromsø, Lillehammer, Bergen and Oslo. In addition, NINA owns and runs the aquatic research station for wild fish at lms in Rogaland and the arctic fox breeding center at Oppdal.

NINA's activities include research, environmental impact assessments, environmental monitoring, counselling and evaluation. NINA's scientists come from a wide range of disciplinary backgrounds that include biologists, geographers, geneticists, social scientists, sociologists and more. We have a broad-based expertise on the genetic, population, species, ecosystem and landscape level, in terrestrial, freshwater and coastal marine ecosystems.

ISSN: 1504-3312
ISBN: 978-82-426-4840-2

Norwegian Institute for Nature Research

NINA head office

Postal address: P.O. Box 5685 Torgarden,
NO-7485 Trondheim, NORWAY

Visiting address: Høgskoleringen 9, 7034 Trondheim

Phone: +47 73 80 14 00

E-mail: firmapost@nina.no

Organization Number: 9500 37 687

<http://www.nina.no>



Cooperation and expertise for a sustainable future