

Consequences of "Natural" Disasters on Aquatic Life and Habitats

Journal:	<i>Environmental Reviews</i>
Manuscript ID	er-2022-0050.R1
Manuscript Type:	Review
Date Submitted by the Author:	06-Jul-2022
Complete List of Authors:	Cooke, Steven; Carleton University Department of Biology, Biology Galassi, Diana M.P.; University of L'Aquila, Department of Life, Health & Environmental Sciences Gillanders, Bronwyn M.; University of Adelaide Landsman, Sean J.; Carleton University, Department of Biology Hammerschlag, Neil; University of Miami, RSMAS Gallagher, Austin J.; Beneath The Waves Eliason, Erika J.; University of California Santa Barbara, Department of Ecology, Evolution and Marine Biology Kraft, Clifford E.; Cornell University, Department of Natural Resources and Environment Taylor, Mark K.; Parcs Canada, Banff Field Unit Crisafulli, Charlie M.; United States Forest Service Shugar, Dan H.; University of Calgary, Earth Sciences Lennox, Robert; Norwegian Institute for Nature Research
Is this manuscript invited for consideration in a Special Issue? :	Not applicable (regular submission)
Keyword:	Geophysical disasters, natural disasters, natural hazards, aquatic ecosystems, ecosystem services

Consequences of “Natural” Disasters on Aquatic Life and Habitats

Steven J. Cooke^{1,*}, Diana M.P. Galassi², Bronwyn M. Gillanders³, Sean J. Landsman¹, Neil Hammerschlag⁴,
Austin J. Gallagher⁵, Erika J. Eliason⁶, Clifford E. Kraft⁷, Mark K. Taylor⁸, Charlie M. Crisafulli⁹, Dan H.
Shugar¹⁰ and Robert J. Lennox¹¹

¹ Department of Biology and Institute of Environmental and Interdisciplinary Science, Carleton University, 1125 Colonel By Dr., Ottawa, ON, K1S 5B6, Canada

² Department of Life, Health & Environmental Sciences, University of L’Aquila, via Vetoio, 67100 L’Aquila, Italy

³ School of Biological Sciences and Environment Institute, University of Adelaide, SA 5005, Australia

⁴ Rosenstiel School of Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Cswy., Miami, FL, 33149, USA

⁵ Beneath the Waves, Herndon, VA, 20172, USA

⁶ Department of Ecology, Evolution and Marine Biology, University of California, Santa Barbara, Santa Barbara, CA, 93106, USA

⁷ Department of Natural Resources and Environment, Cornell University, Ithaca, NY, 14853, USA

⁸ Parks Canada Agency, Banff Field Unit, Banff, AB, T1L 1K2, Canada

⁹ United States Forest Service (retired), Pacific Northwest Research Station, Olympia, WA, 98512, USA

¹⁰ Water, Sediment, Hazards, and Earth-surface Dynamics (waterSHED) Lab, Department of Geoscience, University of Calgary, Calgary, AB, T2N 1N4, Canada

¹¹ Norwegian Institute for Nature Research (NINA) and NORCE, Trondheim, 7034, Norway

*Author for correspondence: Steven.Cooke@carleton.ca

Keywords: Geophysical disasters, natural disasters, natural hazards, aquatic ecosystems, ecosystem services, threats.

29 **Abstract**

30 “Natural” disasters (also known as geophysical disasters) involve physical processes that have a direct or
31 indirect impact on humans. These events occur rapidly and may have severe consequences for resident
32 flora and fauna as their habitat undergoes dramatic and sudden change. Although most studies have
33 focused on the impact of natural disasters on humans and terrestrial systems, geophysical disasters can
34 also impact aquatic ecosystems. Here we provide a synthesis on the effects of the most common and
35 destructive geophysical disasters on aquatic systems (life and habitat). Our approach spanned realms
36 (i.e., freshwater, estuarine, marine) and taxa (i.e., plants, vertebrates, invertebrates, microbes) and
37 included floods, droughts, wildfires, hurricanes/cyclones/typhoons, tornadoes, dust storms, ice storms,
38 avalanches (snow), landslides, volcanic eruptions, earthquakes (including limnic eruptions), tsunamis,
39 and cosmic events. Many geophysical disasters have dramatic effects on aquatic systems. The evidence
40 base is somewhat limited for some natural disasters because transient events (e.g., tornadoes, floods)
41 are difficult to study. Most natural disaster studies focus on geology/geomorphology and hazard
42 assessment for humans and infrastructure. However, the destruction of aquatic systems can impact
43 humans indirectly through loss of food security, cultural services or livelihoods. Many geophysical
44 disasters interact in complex ways (e.g., wildfires often lead to landslides and flooding) and can be
45 magnified or otherwise mediated by human activities. Our synthesis reveals that geophysical events
46 influence aquatic ecosystems, often in negative ways, yet systems can be resilient provided that effects
47 are not compounded by anthropogenic stressors. It is difficult to predict or prevent geophysical
48 disasters but understanding how aquatic ecosystems are influenced by geophysical events is important
49 given the inherent connection between peoples and aquatic ecosystems.

50

51 Introduction

52 “Natural” disasters can be defined as “some rapid, instantaneous or profound impact of the natural
53 environment upon the socio-economic system” (Alexander 2018) usually with a restricted temporal and
54 spatial scale (i.e., rarely global; Turner 1976). These events are often of great magnitude and can thus
55 be further defined as “any manifestations in a geophysical system (i.e., lithosphere, biosphere, or
56 atmosphere) which differs substantially or significantly from the mean [state of the system]” (Alexander
57 2018). The term “natural disaster” is, of course, a misnomer, because every aspect of the impacts -
58 what translates a hazardous event into a disaster - are heavily conditioned by human factors (Smith
59 2006). Thus, herein we use the term *geophysical disaster*. For there to be a disaster, there must be a
60 geohazard, whereby the physical process has a direct or indirect impact on humans. In its most obvious
61 form, one could think about the direct harm of a hurricane where humans are killed and injured, key
62 infrastructure is destroyed or damaged, and essential services such as water, electricity, and
63 communications are disrupted. As the human population has expanded and settled in diverse
64 environments around the globe, the risk posed by geophysical disasters has risen (e.g. Donner and
65 Rodriguez 2008). Encroachment on coastlines, floodplains, forests, and mountains all represent
66 increased exposure of humans and human infrastructure to hazardous phenomena such as floods,
67 landslides, and tsunamis. The United Nations declared 1990-2000 the Decade for Natural Disaster
68 Reduction (Mitchell 1988) emphasizing the heavy toll of disasters on society (Pelling 2001) and the
69 economy (Benson and Clay 2004) along with the recognition that through better planning the impacts
70 from such hazards could be mitigated (Oaks and Bender 1990; Board on Natural Disasters 1999).

71 Given the massive toll of geophysical disasters on humans (average annual mortality of ~ 60,000 people;
72 [Ritchie and Roser 2019], hundreds of billions in economic costs, [Guha-Sapir et al. 2013]) and the
73 apparent growing frequency and magnitude of these events (some of which is driven by anthropogenic
74 climate change; [Van Aalst 2006; Banholzer et al. 2014] – or human activities such as drilling; [Ellsworth
75 2013]), it is not surprising that there has been a surge of research on the topic with a focus on
76 prediction, preparedness, management, and mitigation (Sahil and Sood 2021). However, geophysical
77 disasters also have direct and indirect effects on plants and animals. The environment is constantly
78 changing for plants and animals; slow changes such as gradual temperature warming or changes in
79 oxygen and salinity may be described as ramp stressors (Bender et al. 1984). Many geophysical disasters
80 occur rapidly (Niemi et al. 1990) and require animals to adopt emergency life history stages (Wingfield
81 et al. 1998) as their environment undergoes dramatic and sudden change. For example, wildfires,
82 avalanches, and landslides can displace or kill terrestrial organisms and these effects have been well
83 documented (e.g., reviewed in Zhang et al. 2018; Kaur et al. 2019; Rondeau et al. 2020). However,
84 geophysical disasters can also impact aquatic ecosystems. These disasters have an important role in the
85 succession of ecosystems and the maintenance of biological diversity, but the increasing frequency and
86 severity is troubling given that much of the human infrastructure system is not designed or prepared for
87 increasing severity or frequency of disastrous geophysical conditions. There is much great interest in
88 documenting the response of these aquatic systems to disasters as well as understanding the capacity
89 for species to detect or prepare for sudden changes in their environment.

90 Water covers 71% of the earth and some geophysical disasters (e.g., hurricanes, floods, tsunamis) by
91 definition involve interaction with aquatic systems. Aquatic systems contain much life that is often
92 cryptic yet generates numerous ecosystem services including those of direct benefit to humans including
93 nutritional security and supporting livelihoods (Peterson and Lubchenco 1997; Lynch et al. 2016). This is

94 particularly the case in developing countries, which is also where disproportionate effects of geophysical
95 disasters are often felt (Benson and Clay 2000). Aquatic species, especially in freshwater (Woodward et
96 al. 2010; Doney et al. 2012) or in specific marine habitats (e.g., coral reefs [Hoegh-Guldberg 1999],
97 estuaries [Dyer 2021]), are known to be among the most vulnerable taxa on the planet to ongoing
98 climatic change, which may be exacerbated by stressors that result in sudden and potentially irreversible
99 changes to the stable states of lakes, rivers, estuaries, or marine environments. However, to our
100 knowledge there has yet to be a review of what is known about the effects of geophysical disasters on
101 aquatic systems. To that end, we provide a synthesis on the effects of the most common and
102 destructive geophysical disasters on aquatic life and habitat. Our approach spans realms (i.e.,
103 freshwater, estuarine, marine) and taxa (i.e., plants, vertebrates, invertebrates, microbes). We focused
104 on floods, droughts, wildfires, hurricanes/cyclones/typhoons, tornadoes, dust storms, ice storms,
105 avalanches (snow), landslides, volcanic eruptions, earthquakes (including limnic eruptions), and
106 tsunamis (See Figure 1). We have excluded discussion of cold and warm snaps and routine weather
107 events (e.g., snow, lightning, hail) given that there is much written about the effects of temperature and
108 short-term environmental variability on aquatic life (e.g., Fry 1971; Bhaud et al. 1995; Tittensor et al.
109 2010). Epidemics, although timely and relevant to aquatic life (i.e., the COVID-19 pandemic; Cooke et al.
110 2021), are also outside the scope of this paper given that they have a largely biological basis unlike the
111 other disasters that we cover here.

112 **Disaster Types and Consequences on Aquatic Life and Habitats**

113 Disaster types are organized in what we deem a logical order, focusing initially on events that are largely
114 driven by climate (e.g., floods, drought, hurricanes, etc) and then moving on towards geologically-driven
115 events (e.g., volcanic eruptions, earthquakes) and ending on cosmic events which is somewhat unique.
116 We acknowledge that the evidence base associated with these topics is highly variable. For example,
117 floods, droughts, and wildfires have been quite well studied with respect to their impacts on aquatic
118 systems while for others very little is known. Nonetheless, we attempt to provide equal coverage but
119 necessarily refer readers to other focused syntheses for some of the more studied disasters (see Table 1
120 for list of key syntheses where relevant).

121 **Drought**

122 Drought generally encompasses prolonged dry periods where there is a shortage of water often as a
123 result of a lack of precipitation (Kallis 2008). While drought is a natural process, it has been exacerbated
124 by humans through effects on global warming, construction of impervious surfaces, and water
125 extraction, which affect the frequency, severity, and duration of drought (Bond et al. 2008). Impacts
126 occur across spatial and temporal scales with most studies focusing on short duration droughts (months
127 to a year) in local stream reaches (Matthews and Marsh-Matthews 2003). In standing water, drought
128 associated with water abstraction may decrease lake levels and alter habitat for organisms (Glassic and
129 Gaeta 2019). In flowing water, drought may also reduce habitat availability (Bond et al. 2008). Drought
130 not only affects freshwater systems, but alterations to freshwater inflow also impacts estuarine and
131 marine ecosystems (Gillanders and Kingsford 2002; Lennox et al. 2019). Droughts contribute
132 cumulatively to other extreme environmental perturbations (floods, cyclones, heat waves, fire, dust
133 storms). Severe drought can increase conditions suitable for wildfires. For example, severe drought led
134 to fires in a tributary of the Amazon River that killed most of the floodplain forest trees with little
135 evidence of regeneration even after a decade of recovery (Flores et al. 2014).

136 Reduced freshwater inflow leads to low allochthonous organic matter and nutrient inputs which may
137 impact nearshore phytoplankton (Wetz and Yoskowitz 2013). For example, reduced freshwater
138 discharge into the Nile delta, which was similar to prolonged drought, meant that seasonal
139 phytoplankton blooms did not occur, impacting Mediterranean fisheries (Oczkowski et al. 2009). It was
140 not until anthropogenic nutrient loadings through fertiliser and sewage discharge increased that the
141 fishery started to recover (Oczkowski et al. 2009). Stream organisms including fish and invertebrates are
142 impacted by changes to flow and subsequent impacts on water quality (Bond et al. 2008). Food
143 resources for organisms may also be depleted, increasing potential for biotic interactions (competition
144 and predation). In coastal marine systems, habitats including mangroves, saltmarsh, and seagrass have
145 all been impacted by drought (Duke et al. 2017). Saltmarsh dieback associated with drought has been
146 recorded along the Gulf Coast and South Atlantic Bight in the USA during the early 2000s (Hughes et al.
147 2012; McKee et al. 2004; Silliman et al. 2005). Studies have also reported reductions in seagrass
148 associated with drought linked to reductions in freshwater and nutrient inputs (Hirst et al. 2016), high
149 salinity (Wilson and Dunton 2018), or failure of a facultative mutualistic relationship (de Fouw et al.
150 2016). Reductions in extent of habitats will presumably impact aquatic organisms.

151

152 **Floods**

153 Rivers flood when water input exceeds the capacity for the system to drain or buffer the sudden
154 increase in water input, either from rapid melting of snow or ice, torrential rain, or sudden changes in
155 regulation by dams. Floods cause numerous changes to fluvial systems including increasing water depth,
156 accelerating water velocity, widening channels, regrading sediment, and reducing water temperatures.
157 There may also be potential changes to turbidity, pH, conductivity, salinity, and concentration of
158 pollutants. Lateral expansion of the water into the floodplain may draw pollutants into the water, as was
159 recorded in the 2002 flood on the River Elbe (Einsporn et al. 2005). Extreme flooding has also been
160 implicated as a major catalyst for invasions of non-native species where rivers connect with artificial
161 ponds, rearing facilities, or impoundments (e.g., Kumar et al. 2019). Freshwater effluent in estuaries will
162 also suddenly freshen these ecotones and alter coastal marine communities in the interim (Pollack et al.
163 2011; Bailey and Secor 2016). The incredible force of water to move silt, sand, gravel, and even boulders
164 has a crucial role in the form and function of rivers (Gupta and Fox 1974; Hauer and Habersack 2009)
165 and estuary mouths, which drives the ecology of the species that live there.

166 When flooding occurs, animals may take refuge and survive in eddies, backchannels, or tributaries
167 where the rapid increase in flow is buffered, eventually recolonizing when flows diminish (Koizumi et al.
168 2013) unless they have been washed beyond barriers that restrict upstream passage. For counterparts
169 remaining in the main channel exposed to the full force of the flood, including larvae and eggs, many will
170 be carried downstream and will likely die in harsh environments or become stranded in areas that
171 dewater when floodwaters recede (Nagrodski et al. 2012; Death et al. 2015). Immobile species such as
172 plants and sessile invertebrates will be forced to adapt to the conditions or die. For example, coral
173 bleaching in Hawai'i was exacerbated by the synergistic effect of warm temperatures and freshwater
174 input from flooding in 2014 (Bahr et al. 2015). Eastern oysters (*Perkinsus marinus*) in a Texas estuary
175 responded to flooding with reduced oyster abundance and growth but their rapid life cycle facilitated
176 rapid recolonization (Pollack et al. 2011). As mobile species recolonize following catastrophic floods,
177 they are likely to find themselves occupying new habitat as the river channels have changed course, with
178 re-graded sediments and new structure adopted from the flood debris (Death et al. 2015) as well as
179 shifting delta and estuary structure (Cooper 2002).

180 **Wildfire**

181 Fire is a natural disturbance in many forests across the globe; however, uncontrollable and high intensity
182 wildfire can have catastrophic impacts on social, economic, and environmental systems (UNEP 2022).
183 The effect of wildfire on aquatic ecosystems will depend on the fire characteristics (extent, duration, and
184 severity) as well as local factors (e.g. physical, biochemical and chemical elements of the watershed and
185 previous evolutionary exposure to fire regimes; Gresswell 1999; Bixby et al. 2015; UNEP 2022). Short-
186 term, immediate effects of wildfire can include a reduction in plant biomass and canopy cover,
187 increased runoff and erosion, altered soil and sediment dynamics, nutrient mobilization, and ash
188 deposition (Bixby et al 2015). Wildfires that occur near human activities such as homes and mining
189 increase the risk of toxicants (e.g. benzene, arsenic, lead) accumulating in the watershed (Burke et al.
190 2013; Murphy et al. 2020a). Collectively, these acute wildfire effects tend to negatively impact aquatic
191 life. Fish, amphibian, and invertebrate populations as well as algal cover and biomass all commonly
192 decrease immediately after wildfire (Gresswell 1999; Bixby et al. 2015; Silva et al. 2020; Verkaik et al.
193 2013). Over the longer term (years), stream biota recovery dynamics from wildfire are highly variable
194 (Robinne et al. 2020) and have been linked to the burn severity of the riparian vegetation, whether
195 debris flows occurred, and the connectivity of the watershed (Verkaik et al. 2013; Minshall 2003;
196 Dunham et al. 2003). Severely burned riparian vegetation results in canopy removal and thus increased
197 light and altered temperature regimes (Verkaik et al. 2013). Heavy rains after a fire can cause floods and
198 scouring, resulting in channel reorganization and sediment deposition (Tuckett and Koetsier 2018). More
199 severely impacted streams may have delayed recovery and altered macroinvertebrate aquatic
200 community composition, even 10 years after a fire (Tuckett and Koetsier 2018). However, in connected
201 watersheds, fish recovery can occur within a few years (Dunham et al. 2003).

202 **Hurricanes/Cyclones/Typhoons**

203 Hurricanes or tropical cyclones are destructive storms characterized by excessive wind speeds (typically
204 with winds exceeding 119 km/h). Geophysical effects of hurricanes on aquatic systems include rapid
205 fluctuations in temperature, drops in barometric pressure, and heavy rains as well as wind-driven
206 disturbances to surface and subsurface waters (e.g., currents, waves, surge), substrate and adjacent
207 land. Such geophysical effects can have direct and indirect biological impacts on aquatic life. Storm
208 impacts are greatest for immobile species or life-stages and/or those confined to closed systems, which
209 cannot flee the path of a hurricane. For example, coral reefs can experience breakage and dislodgement
210 with the extent of damage related to wind intensity (Fabricius et al. 2008). In Florida, USA, the passing
211 of a hurricane shifted macroinvertebrate and benthic fish communities via reduction to water salinity
212 and increases in depth (Zink et al. 2020) while a hurricane passing over saltwater marshes in Galveston
213 Bay, Texas reduced nekton abundance but increased diversity (Oakley and Guillen 2020). Peierls et al.
214 (2003) documented increases in phytoplankton following hurricanes with return to normal coinciding
215 with when salinity increased to pre-storm levels more than 1 year after disturbance. Off the west coast
216 of Florida, 69% mortality to pre-hatchling sea turtles was attributed to drowning from storm surge
217 flooding of nests (Milton et al. 1994). Mobile species, able to flee in the wake of the storm, are
218 seemingly less vulnerable to hurricanes. For example, when Hurricane Irene hit the US Mid-Atlantic
219 Bight, many satellite-tagged juvenile and adult loggerhead sea turtles (*Caretta caretta*) moved north of
220 their pre-storm foraging grounds (Crowe et al. 2020). In response to dropping barometric pressure in
221 advance of a hurricane, common snook (*Centropomus undecimalis*) have been found to move down
222 river, potentially exiting the river for deeper marine water that is less exposed to physical disturbance
223 (Massie et al. 2020). Similarly, juvenile blacktip (*Carcharhinus limbatus*) sharks have been found to flee

224 shallow water nursery habitats for adjacent deep-water areas in response to a dropping barometric
225 pressure in the wake of a hurricane (Heupel et al. 2003). However, some juvenile sharks failed to return
226 to their nursery areas following the hurricane, possibly due to predation. Unlike smaller sharks,
227 Gutowsky et al. (2021) found variable responses of large sharks to hurricanes, with adult tiger sharks
228 (*Galeocerdo cuvier*) remaining in shallow waters of the Bahamas, even when the eye of a hurricane
229 passed overhead, with numbers of these apex predators at the site increasing for two weeks following
230 the storm. The study hypothesized that these tiger sharks may have been scavenging on animals that
231 died from the storm (Gutowsky et al. 2021). Teasing apart the relative consequences of physical
232 damage versus changes in water chemistry arising from flooding have proved challenging but will be
233 important for determining how to best mitigate future hurricane damage (Liu et al. 2021).

234 **Tornadoes**

235 Tornadoes and waterspouts (defined as tornadoes that cross from land onto water or originate over
236 water) can have profound impacts on aquatic ecosystems and life within them. For example, people
237 have been recording instances of fauna – including snails, jellyfish, crabs, frogs, and fishes – falling from
238 the sky ('animal rain') for hundreds of years (Gudger 1929), a phenomenon largely attributed to
239 waterspouts and other strong wind events (Morgan 2012; Allaby 2014). Such events have become the
240 focus of books (e.g., Dennis 2013) and are frequently documented in the popular media. 'Animal rain'
241 events, primarily of fish, have also been documented in the academic literature (e.g., James 1894;
242 Bajkov 1949; Dees 1961; Pigg and Gibbs 1998; Roberts 1999). Yet, our understanding of this mode of
243 dispersal is poor (Unmack 2001). Wind-induced movement may be an important method of dispersal
244 over short distances for some taxa such as zooplankton (Havel and Shurin 2004). The extreme wind
245 velocities and intense rain associated with tornadoes can cause physical damage to aquatic and riparian
246 habitats including aggradation (Pierce 2016), boulder dislodgement (de Lange et al. 2006) and possible
247 rockslides that could damage aquatic life and habitat; vegetation removal (Nelson et al. 2008) that could
248 allow more solar radiation to reach the water's surface and thereby increase water temperatures; and
249 changes in water levels associated with the seiche effect (Chaston 1979). Although there are several
250 potential negative consequences associated with tornadoes, they can also create new opportunities for
251 plants and animals to thrive. For example, Nelson et al. (2008) observed both increases to woody stem
252 density following soil disturbance and more shade-tolerant and -intolerant plant species with canopy
253 removal. Enhanced plant growth in riparian zones may benefit aquatic systems by creating more shade
254 and thus reducing water temperatures. Blowdowns from tornadoes can also deposit coarse woody
255 debris, which serves as important habitat for fishes (see Harmon et al. 1986) and increases the amount
256 of allochthonous nutrient inputs. In some cases, tornadoes can create habitats such as breeding pools
257 for Cope's gray tree frogs *Hyla chrysoscelis* (Smith 2013) and new sediment layers (Card 1997). Overall,
258 there is a paucity of research on the impact of tornadoes on aquatic life and habitats, but the impact of
259 tornadoes will likely be variable, localized, and short-term.

260

261 **Dust Storms**

262 Dust or sand storms involve major meteorological processes (e.g., wind) that move inordinate amounts
263 of particles such as dust and sand, especially in the drylands of north Africa and the Arabian peninsula
264 (Goudie 2009). Dust storms transfer massive volumes of particulate matter into lakes and seas given
265 that sediment chronology can be used to detect such events going back as far as 2000 years ago (e.g.,
266 Chen et al. 2013). In addition to transporting dust particles, dust storms can disperse metals

267 (Gunawardena et al. 2013), nutrients (Shi et al. 2013), prokaryotic and eukaryotic algae (Rahav et al.
268 2016), and even pathogenic microorganisms (Gonzalez-Martin et al. 2014) all of which could impact
269 aquatic systems. Algal blooms in marine systems are reasonably common following dust storms as a
270 result of a pulse of nutrient inputs (Bali et al. 2019). Hallegraeff et al. (2014) documented fungal blooms
271 in coastal water following a major dust storm in Australia which did not prove harmful (e.g., non-toxic to
272 fish, minor impacts on algae and coral symbionts), but raised the possibility of other dust storm-driven
273 pathogens that are more pathogenic being transported in future dust storms. Given the transient
274 nature of these events, there is little known about effects on most aquatic animals (Goudie 2009).
275 Interestingly, the creation of reservoirs in areas that typically have dry stream beds have led to
276 reductions in dust storm activity in Mexico (Jáuregui 1989). There is no doubt that dust storms play
277 important roles in distributing nutrients and sedimentary materials (on a global scale) that are key
278 elements of freshwater and marine systems yet in general any negative effects (e.g., algal blooms,
279 microbe deposition) tend to be rather localized and short lived.

280 **Ice Storms**

281 Ice storms occur at locations with precipitation events along frontal transition zones separating warm
282 and cold air masses. Most attention to the biotic impacts of ice storms has focused on the extensive
283 breakage of tree canopy branches in forested ecosystems (Rhoads et al. 2002) that subsequently
284 deposits wood throughout forested watersheds, including stream and lake shoreline environments
285 (Kraft et al. 2002; Millward et al. 2010). As a result, this wood accumulation has substantial impacts on
286 biota within these aquatic ecosystems. Accumulations of wood in streams – commonly referred to as
287 debris dams – are more frequent in number and larger in volume following ice storms (Kraft et al. 2002).
288 These large accumulations of wood have been shown to increase the diversity and abundance of aquatic
289 invertebrates in forested headwater streams (particularly taxa such as Ephemeroptera, Plecoptera, and
290 Trichoptera) compared to riffle habitats (Baillie et al. 2019). Debris dams have been consistently
291 demonstrated to provide habitat for many species of fish (Bisson et al. 1987), and brook trout
292 abundance in forested stream habitats increased in response to the presence of debris dams resulting
293 from an intense ice storm in the northeastern U.S. (Warren and Kraft 2003).

294 Another set of well-studied biotic impacts from ice storm disturbance events has focused on the
295 intersection of tree damage and altered forest processes that reduce nutrient uptake, thereby
296 increasing primary production in downslope streams. This was extensively documented at the Hubbard
297 Brook Experimental Forest in New Hampshire, USA, where increased downstream export of dissolved
298 inorganic nitrogen was observed following an intense ice storm (Houlton et al 2003). Tree canopy
299 damage also increases light availability, reducing light limitation of primary production in streams within
300 forested watersheds (Stovall et al. 2009). Although the Intergovernmental Panel on Climate Change
301 places low confidence in predicting whether climate change will increase the frequency of ice storms
302 (Ranasinghe et al. 2021), IPCC model projections indicate that the location of freezing rain events
303 responsible for ice storms will change in Europe and North America (Kämäräinen et al. 2018; Ning and
304 Bradley 2015).

305 **Landslides**

306 Landslides involve disturbances in the stability of a slope and are often triggered by other disasters (e.g.,
307 droughts, earthquakes) whereas mudslides tend to be induced by the rapid accumulation of water in the

308 ground. These events are sufficiently well researched that there is a realm of study known as “landslide
309 ecology” (Walker and Shiels 2012) with diverse environmental effects well documented (Geertsema et
310 al. 2009). Slides are most common in areas with steep slopes such as river valleys. A major landslide on
311 the Fraser River of British Columbia around 1911 (the Hells Gate slide; see Evenden 2004) temporarily
312 impeded the upstream spawning migration of Pacific salmon in the largest salmon producing river in
313 Canada. A second major landslide on the Fraser River in 2018 (the Big Bar slide; Murphy et al. 2020b) is
314 still being addressed as of January 2022, but only with efforts to remove both the 1911 and 2018 slides
315 has fish passage been somewhat restored. Nonetheless, the effects of the Hells Gate slide are still
316 evident today given the collapse of some populations (Hobbs and Wolfe 2008) and the legacy of the
317 reach being the most hydraulically complex and physiological challenging in the Fraser (Hinch et al.
318 1996). Landslides are known for their recruitment of coarse woody debris to fluvial systems (Ruiz-
319 Villanueva et al. 2014) which helps to create diverse and complex habitats that benefit invertebrates and
320 fish (Harmon et al. 1986; Gurnell et al. 1995). However, they can also mobilize sediment which can
321 degrade habitat and smother spawning grounds for lithophilic fish (Schuster et al. 1989). In some alpine
322 areas, slides can lead to full damming of fluvial systems such that they lead to the development of lakes
323 (Butler and Malanson 1993; Shapley et al. 2019) which represents a major aquatic transition (i.e., from
324 lotic to lentic) but also creates new habitats that can be exploited by some species and cause phase
325 shifts in the plant and animal communities (see Logan and Schuster 1991).

326 Landslides can also impact marine life. An inland landslide in California during the Pleistocene not only
327 impeded the migration of anadromous fish leading to genetic change, but also impeded the
328 downstream transport of sediment to the estuary and offshore areas (Mackey et al. 2011). Landslides
329 instigated by storms have been attributed with the transport of sediment and coarse woody debris to
330 the ocean (West et al. 2011) while coastal landslides have led to pulses in sedimentation (Hapke et al.
331 2003) and contributed to coastal degradation (Nichols et al. 2019) by smothering plants such as macro
332 algae (including kelp forests; Oliver et al. 1999). Indeed, marine coastal organisms are often impacted
333 by the so-called “triad of sediment inputs from landslide activity: direct burial, sediment scouring, and
334 suspended sediment plumes” (Schuster and Highland 2003) that can impede respiration (e.g., of fish or
335 invertebrates) and light penetration (Oliver et al. 1999). Slides can also happen entirely underwater
336 (i.e., submarine landslides) but they are often at great depth and we know more about their physical
337 properties than their biological consequences (Schuster and Highland 2003).

338 **Avalanches (snow)**

339 Similar to landslides, snow avalanches are a slope disturbance and a source of debris. Snow avalanches
340 are a common disturbance where avalanche paths intersect with water, but to our knowledge there has
341 been no research on the effects of snow avalanches on aquatic systems. Nevertheless, we propose
342 direct and indirect effects of these disturbances on aquatic systems below. An example of a direct effect
343 of snow avalanches is the abrupt natural damming of mountain rivers (Butler 1989; Richardson 2000).
344 Outburst floods caused by failure of snow avalanche dams have been reported for several mountain
345 ranges (Andes of Argentina [King 1934]; the European Alps [Allix 1924]; northern Scandinavia [Rapp
346 1960]; the Himalayas [Richardson and Reynolds 2000]; the New Zealand Alps [Ackroyd 1987]) and may
347 have profound impacts on aquatic organisms and their habitats.

348 Indirect effects of snow avalanches on aquatic life likely occur based on the physical disturbance
349 avalanches cause to wetted and riparian habitats in streams, lakes, and marine coastlines. For example,

350 snow avalanches structure riparian vegetation communities in their paths and termini (“snow avalanche
351 ecology”; Butler 1979; Johnson 1987). Landform creation and modification is another indirect link
352 between snow avalanches and aquatic systems given the important role of geomorphology in the
353 maintenance of aquatic habitat for vertebrates and fishes (Vannote et al. 1980). For example, snow
354 avalanches that terminate in stream channels redistribute and add debris on the stream bed (Luckman
355 1978). Boulders and woody debris stabilize channels and create pools that provide cover for fish and
356 invertebrates (Lanka et al. 1987; Ackroyd 1987). The recurring action of avalanches at the foot of slopes
357 can also form erosional depressions that become water-filled ponds known as snow avalanche impact
358 pools (Johnson and Smith 2010). Snow avalanches that terminate in lakes or reservoirs contribute to
359 sediment budgets (Vasskog et al. 2011). Disturbances from snow avalanches are likely a contributing
360 factor to the vertical and horizontal heterogeneities found in the substrates of mountain lakes.
361 Avalanches that terminate in the ocean occur in mountainous coastlines (e.g. Iceland; Johannesson
362 2001). The literature on marine-terminating snow avalanches is focused on hazards to public safety, but
363 snow avalanches could modify the physical structure of coastlines, potentially affecting marine
364 invertebrates, fishes, corals, and plants. We suggest that snow avalanches have a subsidiary role in
365 modifying aquatic life because they have a localized spatial distribution (Luckman 1978) and many snow
366 avalanches contain no debris because they do not come into contact with underlying ground and the
367 ground is typically frozen (Rapp 1960). The contribution of snow avalanches to the physical landscape
368 tends to be masked by the more obvious geomorphic processes involved in debris transport such as
369 landslides/rockfalls, mud-debris flows, and fluvial activity (Luckman 1978).

370

371 **Volcanic Eruption**

372 Volcanoes are incisions in the Earth’s crust where gas and magma can move between the lithosphere
373 and the biosphere. Volcanic eruptions, when gas and matter are discharged from subterranean
374 chambers onto land or into the sea, involve numerous geophysical processes including lava flows, sector
375 collapses and debris avalanches, pyroclastic density currents, lahars (volcanic mudflows), tephra falls,
376 dome building, and chemical plumes, and all or a subset of these may occur during any single eruption
377 event (Swanson and Major 2005; Crisafulli et al. 2015). Volcanism may stand apart from many other
378 forms of natural disturbance because of its diversity of disturbance processes, large spatial extent (10s-
379 1000s km), and potentially interacting physicochemical processes that can generate massive impacts
380 and enduring effects. The majority of scientific inquiry on biotic response to volcanism has been in
381 terrestrial ecosystems, with a strong emphasis on vegetation and arthropods and succession following
382 volcanic disturbance (Crisafulli et al. 2015). Nonetheless, there is a growing body of literature on
383 riverine, lacustrine, and marine environments to complement this knowledge and establish a broader
384 understanding of how the physical and chemical impacts of volcanism affect water and its inhabitants.

385 A primary driver of change in both lakes and rivers is inputs of inorganic ejecta (tephra fall) or flow
386 material (debris avalanche, lahar, pyroclastic flow deposits; matter subsidies), as well as inputs of
387 biological constituents of the surrounding terrestrial biota (i.e. nutrient subsidies; Crisafulli et al. 2015;
388 Dale et al. 2005a,b). The effects of these inputs range from highly ephemeral to protracted in duration,
389 and from relatively minor to profound in volume and extent. Volcanic effects on biota vary between
390 flowing and non-flowing aquatic systems. For example, the deposition of volcanic products into lakes
391 and oceans often have a fertilization effect from nutrient and mineral enrichment of the allochthonous

392 matter (Olgun et al. 2013) that has a bottom-up effect on community structure. On the other hand,
393 large tephra fall events may reduce phytoplankton primary productivity by reducing light transmission
394 because of suspended ash or by extensive mats of floating pumice (Carrillo and Díaz-Villanueva 2021).
395 Whether enrichment or alteration of optical properties, these effects are relatively ephemeral
396 compared to lakes that experience changes in basin morphometry, gross biogeochemical
397 transformations, and enormous log rafts as happened at Spirit Lake following the 1980 eruption of
398 Mount St. Helens, USA (Dale et al. 2005a). For river systems, sediment is the primary driver of ecological
399 responses and may be a protracted problem lasting years to decades in the case of sector collapses from
400 eruptions (Major et al. 2018) or >10m thick pyroclastic deposits (Hayes et al. 2002) versus thinner
401 deposits (<1 m) established by tephra fall events (Arnalds 2013). Sediment directly kills biota through
402 abrasion of the epidermis or smothering of the respiratory organs and indirectly through habitat
403 alteration by filling stream bed interstices and changes to channel morphometry (Bisson et al. 2005). In
404 high-gradient streams, fines are moved through the system during freshets or floods, exposing coarse
405 substrates and altering habitat conditions. Many aquatic organisms are vagile (e.g., insects, amphibians,
406 fish) and quickly recolonize depopulated systems once the physical conditions improve via local
407 immigration. However, volcanic blockages may create barriers to dispersal leading to long-term
408 community reorganization. In the case of both lakes and rivers, different volcanic processes can lead to
409 enormous recruitment of wood or partial to complete removal, with long-term consequences for biotic
410 reassembly and community structure.

411 **Earthquakes**

412 Earthquakes trigger changes of the hydrological regime of springs, streams, lakes, groundwaters, and
413 marine waters (Wang and Manga 2010a; Lubick 2011; Zhang et al. 2021), and are mainly related to the
414 dynamic responses associated with seismic waves (Wang and Manga 2010b). Marine earthquakes alter
415 the structure and function of deep-sea ecosystems (Chunga-Llauce and Pacheco 2021). However,
416 knowledge of their effects on mobile large vertebrates are scant, with the exception of the fin whale
417 (*Balaenoptera physalus*) for which disturbance from strong sounds produced by earthquakes may kill
418 individuals, inducing high speed swimming as a seismic-escape response (Gallo-Reynoso et al. 2011).
419 Sperm whales (*Physeter macrocephalus*) in New Zealand altered their spatial distribution and diving
420 behaviour following habitat changes that occurred as a result of deep-sea canyon “flushing” triggered by
421 the Kaikoura earthquake in 2016 (Guerra et al. 2020). In addition, canyon “flushing” and a resulting
422 turbidity current triggered a massive eradication of benthic invertebrates (Mountjoy et al. 2018). The
423 Kaikoura earthquake affected the intertidal marine vegetation with a massive disruption of the habitat-
424 forming fucoid seaweeds. Epiphytes associated with seaweeds became functionally extinct after the
425 earthquake with less than 0.1 % of the population surviving. The same occurred among seaweed-
426 associated invertebrates (Thomsen et al. 2020). The meiofauna (Giere 2009) seems to be very sensitive
427 to habitat alterations generated by ground shaking, as observed in the marine meiobenthos, mostly
428 represented by the small-sized crustacean copepods after the 2011 Tohoku earthquake in Japan
429 (Kitahashi et al. 2014) likely due to the combination of an increase in organic matter in the surface layers
430 and siltation among sediment particles where these invertebrates live.

431 In inland freshwaters, different biological alterations were related to “earthquake hydrology” (see Mohr
432 et al. 2017; Ingebritsen and Manga 2019). For example, the Mw 6.3 Christchurch earthquake in New
433 Zealand altered the spawning habitat requirements (i.e., salinity gradients) of ‘īnanga’ (*Galaxias*

434 maculatus), an anadromous riparian-spawning fish. The species was forced to migrate 2 km upstream in
435 rivers where several anthropogenic land uses threatened the populations (Orchard et al. 2018). In urban
436 streams, invertebrate taxonomic richness decreased, and benthic Ephemeroptera, Plecoptera, and
437 Trichoptera disappeared due to post-earthquake siltation. Fish richness and density decreased
438 significantly, with fish absent from some heavily silted streams (Harding and Jellyman 2015). Brancelj
439 (2021) observed species replacement in the zooplankton of Slovenian lakes after earthquakes, with
440 severe changes in the zooplankton biomass likely related to a pulse input of nutrients and resuspension
441 of fine sediments. After earthquakes, the increase in groundwater discharge was documented on many
442 occasions (Mohr et al. 2017). The co-seismic aquifer strain *biotriggered* a flushing of the groundwater
443 meiofauna, with a dramatic decrease in subterranean species abundance (Galassi et al. 2014; Fattorini
444 et al. 2018). The repercussions of this event were impressive, given the low resilience of the
445 subterranean communities, being the species characterized by considerable longevity and low fertility.
446 In groundwater-fed springs, a lower abundance of obligate groundwater-dweller microcrustaceans four
447 years after the main shock was observed, together with a higher post-seismic niche overlap among
448 groundwater- and surface-water species at the spring outlets (Fattorini et al. 2017).

449 **Limnic eruptions**

450 A limnic eruption is not necessarily related to a volcanic eruption and it does not refer to the expulsion
451 of magma, but to the “explosion” in some lakes of dissolved gases which are toxic for humans, aquatic
452 and terrestrial invertebrates and vertebrates. For this reason these crater lakes (Kusakabe 2017) are
453 called “killer lakes” or “exploding lakes” (Shanklin 1989). They are somehow silently toxic under various
454 conditions that occur suddenly. The gas originates from magma at great depth, and dissolves into
455 groundwater near the Earth’s surface (Kling et al. 2005). The CO₂-enriched water enters the lake
456 bottoms through springs. These rare events are well known from Lake Nyos and Lake Manoun in
457 Cameroon, despite similar events being known from Lake Averno in Italy (Tassi et al. 2018) and Lake Kivu
458 (4,000 years ago) at the border between Rwanda and the Democratic Republic of the Congo (Hirslund
459 2020). In Lake Nyos and Lake Manoun water mixing is very limited, either because they are tropical lakes
460 where the temperature remains high year-round, and show also a chemocline, where the deep water is
461 more dense due to the presence of CO₂, CH₄, other volcanic gases, and total dissolved salts. The
462 condition at the lake surface may remain relatively stable until unexpected events such as earthquakes
463 or landslides occur and trigger the rupture of the stratification (lake overturn), thus allowing toxic gases
464 to reach the surface waters and the atmosphere around the lake. Degassing through the lake surface
465 occurs by bubbling or by diffusion through the water/air interface (Hernández et al. 2021).

466 Very little is known about their communities both in the planktonic habitat and in the deeper benthic
467 layers. In both lakes, Proteobacteria for Bacteria and Crenarchaea for Archaea were dominant and
468 present at all depths but in different proportions. In these meromictic lakes pH, O₂ or CO₂
469 concentrations, ions and nutrients would affect the abundance, activity and diversity of bacterial and
470 archaeal populations (Nana et al. 2020). Fish are routinely introduced with breeding populations but
471 whenever this has happened a subsequent lake overturn deoxygenated the surface water and killed all
472 the invertebrates and vertebrates living there. This may be also the case of the planktonic cladoceran
473 crustacean recorded from this lake (Green and Kling 1988). Interestingly, gaps in plankton fossil
474 recording at the bottom of Lake Kivu suggest that such sudden events occurred more than once in
475 several lakes with similar characteristics in the last 5,000 years (Nayar 2009). Lake Kivu is known to host

476 more than 28 fish species and a diversified planktonic community (Sarmiento et al. 2006; Darchambeau
477 et al. 2012). The lake biodiversity is likely supported by its higher altitude if compared to other African
478 lakes with the same characteristics, and its greater depth, thus determining cooler temperatures at the
479 surface that may support a stronger stratification between the oxic and predominantly autotrophic
480 (Borges et al. 2014) mixolimnion up to 70 m and the deep monimolimnion rich in CH₄ and CO₂. From
481 data of recent studies, it seems that methane and carbon dioxide concentrations in Lake Kivu are
482 currently close to a steady state (Bärenbold et al. 2020).

483

484 **Tsunamis**

485 Tsunamis are massive and powerful waves that are most commonly generated by earthquakes, and less
486 commonly by submarine (or terrestrial) landslides or aquatic cosmic impacts. The magnitude of tsunami
487 effects on aquatic life have been relatively well documented in a few specific regions, due to the
488 geographically isolated extent in which they occur. Due to the damaging physical contact of a tsunami
489 wave, coastal marine plants are often ripped out, destabilizing substrates and leading to drastic changes
490 in community structure. Seagrass coverage decreased rapidly following a tsunami that hit Sumatra,
491 Indonesia (Nakaoka et al. 2007). The coverage of corals and mangroves decreased ~10% and ~47% after
492 a tsunami hit India in 2004 (Majumdar et al. 2018). The species composition of coastal fish communities
493 was affected by this tsunami, yet overall fish diversity did not change (Sathianandan et al. 2012). The
494 movements of marine sediments produced by tsunamis have the potential to make toxic metals
495 bioavailable by stirring up sediments and resulted in an increase in the metal content in the muscle
496 tissue of mollusks off of Chile (Tapia et al. 2019).

497 Some of the most comprehensive data for tsunami impacts on aquatic life resulted from research
498 following the Great East Japan earthquake (magnitude 9.0) of March 2011, which generated a wave
499 extending up to 20 m at maximum height as it struck the northeastern part of Honshu Island of Japan.
500 Following the tsunami, the community structure of seagrass beds (*Zostera marina*) showed a decrease in
501 vegetation coverage, as did the biomass and abundance of seagrass-associated fish species, relative to
502 pre-tsunami levels (Shoji and Morimoto 2016). Sea urchin densities decreased rapidly after the event,
503 leading to an indirect increase in kelp abundance, however these impacts were not seen at sites that
504 were afforded greater protection from the wave impact (Muraoka et al. 2017). In other locations, the
505 abundance, diversity, and species composition of shallow demersal fish assemblages did not appear to
506 change significantly, which may be explained by the translocation or movements of more mobile fishes,
507 enabling high survival rates (Okazaki et al. 2017). Juvenile growth rates of a regional flounder species off
508 Japan showed no change up to two years after the event (Kurita et al. 2017). Populations of Pacific cod
509 (*Gadus macrocephalus*) off northeastern Japan showed a remarkable four-fold increase in the three
510 years after the tsunami, which was suggested to be linked to lower mortality resulting from a marked
511 decrease in fishing mortality arising from damage to the fishing fleet (Narimatsu et al. 2017). In some
512 areas, affected aquatic communities actually showed a gradual recovery to pre-tsunami levels (seven
513 years; Shoji et al. 2021), suggesting that shallow shores may have long term resilience to tsunamis.
514 However, clear community-shifts in dominant fish species have been detected, suggesting local ecology
515 and productivity may be profoundly affected at small spatial scales, leading to legacy effects (Shoji and
516 Morimoto 2016).

517 **Cosmic events**

518 Lesser-known forms of geophysical disasters are of cosmic origin. The most extreme would be a strike
519 from an asteroid or comet, which is credited with the Cretaceous–Paleogene mass extinction event that
520 included aquatic organisms (D’Hondt 2005). In due course (e.g., the consequences of the strike on the
521 atmospheric conditions such as temperature), such an event would likely be globally catastrophic to
522 most forms of life (Toon et al. 1995) even if the strike occurred in the oceans (Patchett et al. 2016;
523 Rampino et al. 2019). Meteors are smaller and much more common than asteroids and comets. We
524 failed to identify any studies of meteors on aquatic life. Similarly, the impacts of solar flares on
525 biodiversity of any sort have been little studied although such events could lead to rapid changes in
526 temperature (Somov 1991), which would presumably impact aquatic life.

527 Geomagnetic storms are a temporary disturbance of the Earth's magnetosphere and occur relatively
528 frequently (several per year). Given the reliance of aquatic wildlife (ranging from microbes to fish to
529 mammals) on the magnetosphere to assist with navigation via magnetoreception (Wiltshcko and
530 Wiltshcko 2005; Monteil and Lefevre 2020), there is a reasonably large body of work on the effects of
531 geomagnetic storms on aquatic life. For example, geomagnetic storm conditions are known to disrupt
532 circadian biochemical processes which are believed to be mediated through melatonin and
533 cryptochrome (Close 2012; Krylov 2017). Simulated geomagnetic storms alter the behaviour of fish (Fitak et
534 al. 2020) and crabs (Muraveiko et al. 2013) and have been implicated in the mass stranding of cetaceans
535 (Pulkkinen et al. 2020; Zellar et al. 2021). There are a number of studies that document developmental
536 effects on fish embryos if exposures occur during early developmental stages; using rudd as a model,
537 researchers have documented reductions in condition indices and morphological abnormalities (Krylov
538 et al. 2017, 2019) which may be a result of alterations in digestive function (Golovanova et al. 2015).
539 There are also documented effects on zooplankton life-history traits (via maternal effects; Krylov and
540 Osipova 2019). Although research is still in its infancy, of all the cosmic events, geomagnetic storms
541 appear to be highly relevant to aquatic life.

542 **Cascading Disasters**

543 Some major disasters commonly involve more than a single geophysical driver or component, while
544 others are made worse by preconditioning from previous events. Referred to variably as cascading,
545 compound, or complex disasters (see Cutter 2018), they appear to be growing in frequency (Kumasaki et
546 al. 2016). For example, the atmospheric river that affected much of southwest British Columbia and
547 Washington State in late 2021 resulted in flooding in lowlands, as well as substantial mass wasting in
548 upland terrain; it was the debris flows that severed multiple highways, leading to supply chain
549 disruptions across much of Canada. While no research has yet been published, it is plausible that some
550 of these debris flows originated in areas burned by wildfires in recent years (Gillett et al. 2022). In South
551 Korea, wildfires and heavy rains caused landslides that extirpated a number of stream fish including rare
552 and Endangered species (Cho et al. 2003). Similarly, flooding in Australia in 2020 following heavy
553 precipitation was likely exacerbated by intense wildfires the previous summer. The floods resulted in
554 substantial erosion and increased turbidity in streams, impoverishing water quality (Kemter et al. 2021).
555 In coastal British Columbia, a complicated hazards cascade in November 2020 involving a landslide,
556 which triggered a tsunami in a lake, and an outburst flood from that lake that resulted in intense
557 scouring of about 8.5 km of salmon spawning habitat in Southgate River and Elliot Creek (Geertsema et
558 al. 2022).

559 Assessing multi-hazard interrelationships is challenging, particularly in a predictive context (Tilloy et al.
560 2019). Nonetheless, given that water flows downstream, it is not surprising that such hazards can have
561 diverse consequences on aquatic ecosystems when they involve inland aquatic systems. Threats to
562 aquatic life compound, often in synergistic ways to yield outcomes that are somewhat unpredictable
563 (Folt et al. 1999). Given that the intersection of hazards can occur over long time scales (e.g., wildfire or
564 volcanic ash may not intersect with floods until years later when there is a large precipitation event) the
565 actual and potential hazard risk to aquatic ecosystems can vary over time.

566 **Human-Mediated Disasters**

567 Given that we are in the Anthropocene it is unsurprising that human activities are increasingly mediating
568 disasters and their consequences on aquatic systems. For example, freshwater, estuarine and marine
569 ecosystems are already experiencing many anthropogenic threats. When hazards occur it is on top of
570 existing threats that can lead to exceedance of tipping points (Stelzer et al. 2021). Disasters themselves
571 may be spurred by human activities. Rapid deforestation (and the installation of logging roads) can have
572 dramatic effects on freshwater life (e.g., changes in hydrology, water temperature, nutrient fluxes) in
573 diverse systems ranging from Boreal (Kreutzweiser et al. 2008) to tropical (Chapman and Chapman
574 2002) forests. However, logging can also set the conditions for hazards to develop including floods and
575 landslides (Jakob 2000; Schuster and Highland 2007), exacerbating the impacts of logging on aquatic
576 systems (Hartman et al. 1996). In the coming decades, climate change will increasingly be both a driver
577 of hazards (e.g., fires, dust storms, ice storms, hurricanes) but will also pressure aquatic systems
578 because the impacts of hazards will be exacerbated as aquatic ecosystems become less resilient (Klein et
579 al. 2003). Dale et al. (2001) noted how climate change can impact forests by altering the frequency,
580 intensity, duration, and timing of fire, drought, introduced species, insect and pathogen outbreaks,
581 hurricanes, windstorms, ice storms, or landslides. Forests are essential for aquatic systems and thus it is
582 apparent how climate change may lead to additional hazards for aquatic life. Climate change may also
583 contribute to more cascading hazards (Lawrence et al. 2020; see above). The full extent of climate
584 change on natural hazards and cascading effects on aquatic systems are poorly understood but it is
585 almost certain to make things worse.

586 **Synthesis**

587 Natural hazards (See Figure 1) have been demonstrated to alter aquatic ecosystems and biodiversity in
588 diverse ways. Of course, the intensity, scale (time and space) and frequency of natural disasters is highly
589 variable and may be changing with altered climate systems. Similarly, not all ecosystems and biota will
590 respond to disasters in the same way. Freshwater systems seem to be more vulnerable than marine
591 systems when considering the impacts of natural disasters on aquatic life, however the relative
592 susceptibility of these two realms to different types of disasters is highly variable such that direct
593 contrasts are challenging. For example, tsunamis are unlikely to have major impacts on freshwater
594 systems (albeit some evidence of lake tsunamis; see Lockridge 1990) whereas avalanches are
595 presumably irrelevant to marine systems). What is clear is that the evidence base remains diffuse with
596 few studies that have occurred for a given disaster type where responses have been examined using a
597 variety of biological endpoints that transcend the individual to the assemblage. Examples such as those
598 we have drawn upon for this synthesis are rarely replicated (a function of disasters rarely being
599 predictable) and often lack experimental designs that include relevant comparators (e.g., using a before-
600 after control-impact design; Smith 2002). Nonetheless, in most studies of the effects of natural hazards

601 on aquatic systems a decline in biodiversity was detected usually in the form of reduced population
602 sizes. Yet, there is little data at the level of the assemblage or changes in species' interactions and
603 ecosystem functions. Moreover, sublethal effects (e.g., behavioural alterations in animals, changes in
604 plant respiration) have rarely been studied for all but a few disaster types (e.g., fires, floods, hurricanes,
605 some cosmic events).

606 The review we conducted here was not systematic and there are certainly evidence clusters for some
607 disaster types and endpoints (e.g., flood impacts on riverine fish populations; Rytwinski et al. 2020) that
608 have enabled formal systematic review and meta-analysis. Unfortunately, we are decades away from
609 being able to do that for most disaster types and endpoints. Building a robust evidence base that
610 wherever possible employs an experimental approach (making use of so-called natural experiments) is
611 needed recognizing the reality that many of the studies we referred to above were opportunistic. On
612 occasion a disaster can lead to the development of a long term research program, best exemplified by
613 the extensive body of work on the biophysical impacts of the Mount St. Helens volcanic eruption in
614 Washington State, USA (see Crisafulli and Dale 2018). However, this is relatively uncommon. There are
615 certainly researchers that devote their lives to studying the ecological consequences of some disasters
616 such as floods, droughts, and fires on aquatic systems. Yet, in most instances it is our suspicion that
617 researchers drawn into such work are responding to a local need or opportunity. A good example arises
618 from the work of several of the authors on this paper who were conducting a telemetry study on the
619 spatial ecology of sharks in the Caribbean when several notable hurricanes happened to move through
620 the area where the work was underway. Although not part of the initial study plan, the authors were
621 able to assess behavioural responses of sharks to hurricanes using their dataset (see Gutowsky et al.
622 2021). The opportunistic nature of such research has likely contributed to the evidence base being
623 disparate and diffuse. To be clear, that is not a critique of any individual study but a recognition that
624 much of the work done in this space occurs as discrete projects rather than as research programs.
625 Laboratory experiments can be conducted for some disasters (e.g., simulated flood, drought and
626 barometric pressure conditions, exposing organisms to electromagnetic fields to simulate cosmic events,
627 exposing organisms to fire or volcanic ash) which are a useful complement to more opportunistic field
628 studies. Research that combined lab simulations with field mesocosm research and studies of real
629 events would be profitable.

630 **Future Research Directions**

631 Besides the aforementioned discussion about the evidence base and limitations with some of the
632 studies, there are some notable research questions that need to be addressed. Each of these could
633 represent many careers and research programs given the need for research on different disasters, in
634 different ecosystems, in different regions, focused on different endpoints. We note that to tackle these
635 questions in a fulsome way will require collaboration among disciplines including ecology,
636 limnology/oceanography, geosciences, physics, engineering, chemistry, and even social science.
637 Questions are ordered in what we consider a logical progression but their order does not imply any
638 prioritization.

- 639 • How does disaster type, magnitude, frequency, and spatial scale influence ecosystem resilience
640 to an event?
- 641 • What is the relative resilience of different aquatic ecosystems (e.g., inland waters, estuaries,
642 coastal marine, offshore marine) and biological assemblages to various natural disasters?

- 643 • How does spatial and temporal extent of a given disaster influence aquatic ecosystems and
644 biological assemblages?
- 645 • What are the mechanistic links between natural disasters and observed changes at the level of
646 the organism, population or assemblage?
- 647 • Can geophysical disasters lead to regime shifts in aquatic ecosystems?
- 648 • How do natural hazards intersect with anthropogenic stressors (e.g., land use change, invasive
649 species, pollution) to influence the scope and severity of impacts on aquatic ecosystems?
- 650 • How will climate change influence the frequency, severity, and consequences of natural hazards
651 on aquatic ecosystems?
- 652 • To what extent are ecosystem services to humans directly or indirectly dependent on and/or
653 disrupted by geophysical disasters to aquatic life?
- 654 • What are the best ways to prepare for natural disasters in ways that contribute to the resilience
655 of aquatic systems?
- 656 • What is the potential of applying a Social Ecological Systems framework (sensu Ostrom 2009) to
657 geophysical disaster responses in order to increase resiliency and mitigate potential negative outcomes
658 to both humans and aquatic life?
- 659 • How may aquatic systems make communities and societies more resilient to natural disasters?
660 (see Eriksson et al. 2017 for example on the role of fish and fisheries in enabling community recovery
661 from natural hazard on Vanuatu).

662 **Conclusions**

663 We documented diverse examples of how various natural disasters influence aquatic life at different
664 levels of biological organization spanning the individual to the ecosystem. Effects were highly variable
665 with different spatial and temporal impacts. In some cases there was evidence of resilience to disasters.
666 In other cases, there was recovery ranging from short periods (days, weeks) to longer periods (years to
667 decades). Given the patchwork of research on different taxa and levels of organization, it is difficult to
668 draw strong conclusions. Some forms of natural disaster are well studied in terms of aquatic impacts
669 (e.g., wildfires, flood, drought, volcanic eruptions) whereas for some others (avalanches, cosmic events,
670 and tornadoes), the evidence base is sparse (Table 1). As noted above, many research gaps remain.
671 There is no doubt that natural disasters will continue to occur. Moreover, some types of disasters may
672 become more common as a result of human activities and anthropogenic climate change. Given that
673 many aquatic systems are already under threat as a result of pollution, invasive species, habitat
674 alteration and exploitation, with associated losses in freshwater (Reid et al. 2019), estuarine (Kennish
675 2002), and marine (Crain et al. 2009) biodiversity, it is probable that the effects of natural disasters on
676 aquatic life may become more pronounced, as they interact with the aforementioned stressors.
677 Although it is difficult if not impossible to prevent many of the natural disasters discussed here, in areas
678 where they are common (e.g., flood prone systems, areas subject to frequent hurricanes), some
679 planning can be done in an effort to ensure that where possible, efforts are taken to attempt to mitigate
680 impacts. This may be best achieved through win-win scenarios such as protecting mangroves along
681 shoreline to mitigate effects of tsunamis and hurricanes, typhoons and cyclones on both humans and

682 aquatic life. Such actions are also regarded as nature-based solutions to climate change (Seddon et al.
683 2020). There are also opportunities to explore various restoration strategies as has been done
684 extensively for volcanic eruptions and wildfires in an attempt to expedite recovery. Of course,
685 addressing some of the aforementioned anthropogenic stressors and restoring aquatic ecosystems and
686 biodiversity would help to make aquatic systems more resilient to natural disasters.

687 **Acknowledgements**

688 We thank Richard Amos for comments on an early draft of the manuscript. The figure was created by
689 Hiram Henriquez from H2H Graphics & Design Inc in a pay-for-service arrangement. Mike Dusevic and
690 Gillian Zorn assisted with formatting references. We are grateful to two referees for thoughtful
691 comments on our manuscript.

692 **Funding Statement:** Cooke was supported by the Natural Sciences and Engineering Research Council of
693 Canada via an NSERC Discovery Grant.

694 **Competing Interests Statement:** Cooke is on the editorial board of Environmental Reviews but was not
695 involved in handling the manuscript.

696 **Data Availability Statement:** This is a narrative review paper and does not include any empirical data or
697 analysis given a weak evidence base for most topics explored here.

698

699 **References**

700

701 Ackroyd P. 1987. Erosion by Snow Avalanche and Implications for Geomorphic Stability, Torlesse Range,
702 New Zealand, *Arctic and Alpine Research* 19:65-70

703 Alexander D. 2018. *Natural disasters*. Routledge, NY.

704 Allaby M. 2014. *Tornadoes*. Infobase Publishing, NY.

705 Allix A. 1924. Avalanches. *Geographical Review* 14:519-56

706 Arnalds O. 2013. The influence of volcanic tephra (ash) on ecosystems. *Advances in Agronomy* 121:331-
707 380

708 Bahr KD, Jokiel PL, Rodgers KS. 2015. The 2014 coral bleaching and freshwater flood events in Kāneʻohe
709 Bay, Hawaiʻi. *PeerJ* 3:1136

710 Bailey H, Secor DH. 2016. Coastal evacuations by fish during extreme weather events. *Scientific reports*
711 6:1-9

712 Baillie BR, Hicks BJ, Hogg ID, Van den Heuvel MR, Kimberley MO. 2019. Debris dams as habitat for
713 aquatic invertebrates in forested headwater streams: A large-scale field experiment. *Marine and*
714 *Freshwater Research* 70:734–744

715 Bajkov AD. 1949. Do fish fall from the sky? *Science* 109:402-402

- 716 Bali K, Mishra AK, Singh S, Chandra S, Lehahn Y. 2019. Impact of dust storm on phytoplankton bloom
717 over the Arabian Sea: a case study during March 2012. *Environmental Science and Pollution Research*
718 26:11940-11950
- 719 Banholzer S, Kossin J, Donner S. 2014. The impact of climate change on natural disasters. In A. Singh & Z.
720 Zommers (Eds.), *Reducing disaster: Early warning systems for climate change* (pp. 21–49). Dordrecht,
721 the Netherlands: Springer, Netherlands.
- 722 Bärenbold F, Boehrer B, Grilli R, Mugisha A, von Tümpling W, Umutoni A. 2020. No increasing risk of a
723 limnic eruption at Lake Kivu: Intercomparison study reveals gas concentrations close to steady state.
724 *PLoS ONE* 15:8
- 725 Bender EA, Case TJ, Gilpin ME. 1984. Perturbation experiments in community ecology: theory and
726 practice. *Ecology* 65:1–15
- 727 Benson C, Clay EJ. 2000. Developing countries and the economic impacts of natural disasters. In
728 *Managing disaster risk in emerging economies*. A. Kreimer and M. Arnold, Eds. Pp 11–21. World Bank,
729 Washington, DC.
- 730 Benson C, Clay EJ. 2004. *Understanding the economic and financial impacts of natural disasters* (No. 4).
731 World Bank Publications, Washington, DC.
- 732 Bhaud M, Cha JH, Duchene JC, Nozais C. 1995. Influence of temperature on the marine fauna: what can
733 be expected from a climatic change. *Journal of Thermal Biology* 20:91-104
- 734 Bisson PA, Bilby RE, Bryant MD, Dolloff CA, Grette GB, House RA, Murphy ML, Koski KV, Sedell JR. 1987.
735 Large woody debris in forested streams in the Pacific Northwest: past, present, and future. *In*
736 *Streamside management: forestry and fishery interactions*. Edited by E.O. Salo and T.W. Cundy.
737 University of Washington, Seattle. 143–190 pp.
- 738 Bisson PA, Crisafulli CM, Fransen BR, Lucas RE, Hawkins CP. 2005. Responses of fish to the 1980 eruption
739 of Mount St. Helens. Pp 163–182 in Dale VH, Swanson FJ, Crisafulli CM, eds. *Ecological Responses to the*
740 *1980 Eruption of Mount St. Helens*. New York: Springer.
- 741 Bixby RJ, Cooper SD, Gresswell RE, Brown LE, Dahm CN, Dwire KA. 2015. Fire effects on aquatic
742 ecosystems: an assessment of the current state of the science. *Freshwater Science* 34:1340-1350
- 743 Board on Natural Disasters. 1999. Mitigation emerges as major strategy for reducing losses caused by
744 natural disasters. *Science* 284:1943-1947
- 745 Bond NR, Lake PS, AH. Arthington. 2008. The impacts of drought on freshwater ecosystems: an
746 Australian perspective. *Hydrobiologia* 600:3-16
- 747 Borges AV, Morana C, Bouillon S, Servais P, Descy J-P, Darchambeau F. 2014. Carbon Cycling of Lake Kivu
748 (East Africa): Net Autotrophy in the Epilimnion and Emission of CO₂ to the Atmosphere Sustained by
749 Geogenic Inputs. *PLoS ONE* 9:109500

- 750 Brancelj A. 2021. Shifts in zooplankton communities in high-mountain lakes induced by singular events
751 (fish stocking, earthquakes): evidence from a 20-year survey in Slovenia (Central Europe). *Aquat Ecol.*
752 55:1253–1271
- 753 Burke MP, Hogue TS, Kinoshita AM, Barco J, Wessel C, Stein ED. 2013. Pre-and post-fire pollutant loads
754 in an urban fringe watershed in Southern California. *Environmental monitoring and assessment*
755 185:10131-10145
- 756 Butler DR. 1979. Snow avalanche path terrain and vegetation, Glacier National Park, Montana. *Arctic*
757 *and Alpine Research* 11:17-32
- 758 Butler DR. 1989. Snow avalanche-dams and resultant hazards in Glacier National Park, *Northwest*
759 *Science* 63:109–115
- 760 Butler DR, Malanson GP. 1993. Characteristics of two landslide-dammed lakes in a glaciated alpine
761 environment. *Limnology and Oceanography* 38(2): 441.
- 762 Card VM. 1997. Varve-counting by the annual pattern of diatoms accumulated in the sediment of Big
763 Watab Lake, Minnesota, AD 1837–1990. *Boreas* 26:103-112
- 764 Carrillo U, Díaz-Villanueva V. 2021. Impacts of volcanic eruptions and early recovery in freshwater
765 environments and organisms. *Biological Reviews* 96(6): 2546-2560
- 766 Chapman LJ, Chapman CA. 2002. Tropical forest degradation and aquatic ecosystems: our current state
767 of knowledge. Conservation of freshwater fishes: options for the future (MJ Collares-Pereira, IG Cowx &
768 MM Coelho, ed.). Blackwell Science, Oxford 237-249 pp.
- 769 Chaston PR. 1979. An unusual weather phenomenon: The Rochester Seiche. *Weatherwise* 32:211-211
- 770 Chen F, Qiang M, Zhou A, Xiao S, Chen J, Sun D. 2013. A 2000-year dust storm record from Lake Suga
771 in the dust source area of arid China. *Journal of Geophysical Research: Atmospheres* 118:2149-2160
- 772 Cho GI, Lee CW, Kim GY, Joo GJ. 2003. The Impacts of the Fire and Landslide of the Mountain Streams on
773 the Fish in the Northeastern Park of S. Korea. In Proceedings of the Zoological Society Korea Conference.
774 The Korean Society for Integrative Biology 153-3 pp.
- 775 Chunga-Llauce JA, Pacheco AS. 2021. Impacts of earthquakes and tsunamis on marine benthic
776 communities: A review. *Marine Environmental Research* 171:105481
- 777 Close J. 2012. Are stress responses to geomagnetic storms mediated by the cryptochrome compass
778 system?. Proceedings of the Royal Society B: *Biological Sciences* 279:2081-2090
- 779 Cooke SJ, Twardek WM, Lynch AJ, Cowx IG, Olden JD, Funge-Smith S, Britton JR. 2021. A global
780 perspective on the influence of the COVID-19 pandemic on freshwater fish biodiversity. *Biological*
781 *Conservation* 253:108932
- 782 Cooper JAG. 2002. The role of extreme floods in estuary-coastal behaviour: contrasts between river-and
783 tide-dominated microtidal estuaries. *Sedimentary Geology* 150:123-137
- 784 Crain CM, Halpern BS, Beck MW, Kappel CV. 2009. Understanding and managing human threats to the
785 coastal marine environment. *Annals of the New York Academy of Sciences* 1162(1): 39-62

- 786 Crisafulli CM, Dale VH. 2018. Ecological responses at Mount St. Helens: revisited 35 years after the 1980
787 eruption. New York: *Springer* 47-111 pp.
- 788 Crisafulli CM, Swanson FJ, Halvorson JJ, Clarkson B. 2015. Volcano ecology: Disturbance characteristics
789 and assembly of biological communities. in: H. Sigurdsson, B. Houghton, S. McNutt, H. Rymer, and J. Stix,
790 eds. *Encyclopedia of Volcanoes 2nd Edition*. London: Elsevier Publishing. 1265–1284 pp.
- 791 Crowe LM, Hatch JM, Patel SH, Smolowitz RJ, Haas HL. 2020. Riders on the storm: loggerhead sea turtles
792 detect and respond to a major hurricane in the Northwest Atlantic Ocean. *Movement Ecology* 1-13 pp.
- 793 Cutter SL. 2018. Compound, cascading, or complex disasters: what's in a name? *Environment: Science
794 and Policy for Sustainable Development* 60:16-25
- 795 Dale VH, Crisafulli CM, Swanson FJ. 2005a. 25 years of ecological change at Mount St. Helens. *Science
796* 308(5724): 961-962
- 797 Dale VH, Joyce LA, McNulty S, Neilson RP, Ayres MP, Flannigan MD, Wotton, BM. 2001. Climate change
798 and forest disturbances: climate change can affect forests by altering the frequency, intensity, duration,
799 and timing of fire, drought, introduced species, insect and pathogen outbreaks, hurricanes, windstorms,
800 ice storms, or landslides. *BioScience* 51:723-734
- 801 Dale VH, Swanson FJ, Crisafulli CM. 2005b. Disturbance, survival, and succession: understanding
802 ecological responses to the 1980 eruption of Mount St. Helens. In *Ecological responses to the 1980
803 eruption of Mount St. Helens* (pp. 3-11). Springer, New York, NY.
- 804 Darchambeau F, Isumbisho M, Descy JP. 2012. Zooplankton of Lake Kivu. In: J.-P. Descy, F. Darchambeau,
805 M. Schmid (Eds.), *Lake Kivu — Limnology and biogeochemistry of a tropical great lake: Aquatic Ecology
806 Series*, vol. 5, Springer, Dordrecht. pp. 107-126
- 807 Death RG, Fuller IC, Macklin MG. 2015. Resetting the river template: The potential for climate-related
808 extreme floods to transform river geomorphology and ecology. *Freshwater Biology* 60: 2477-2496
- 809 Dees LT. 1961. Rains of fishes. US Fish. Wildl. Serv. Bur. Commercial Fisheries. Fisheries Leaflet 513.
- 810 de Fouw J, Govers LL, van de Koppel J, van Belzen J, Dorigo W, Cheikh MAS, Christianen MJA, van der
811 Reijden KJ, van der Geest M, Piersma T, Smolders AJP, Olf H, Lamers LPM, van Gils JA, van der Heide T.
812 2016. Drought, mutualism breakdown, and landscape-scale degradation of seagrass beds. *Current
813 Biology* 26:1051-1056
- 814 de Lange WP, de Lange PJ, Moon VG. 2006. Boulder transport by waterspouts: an example from Aorangi
815 Island, New Zealand. *Marine Geology* 230:115-125
- 816 Dennis J. 2013. *It's raining frogs and fishes: four seasons of natural phenomena and oddities of the sky
817* (Vol. 1). Diversion Books, New York.
- 818 D'Hondt S. 2005. Consequences of the Cretaceous/Paleogene mass extinction for marine ecosystems.
819 *Annu. Rev. Ecol. Evol. Syst.* 36:295-317
- 820 Doney SC, Ruckelshaus M, Emmett Duffy J, Barry JP, Chan F, English CA, Talley LD. 2012. Climate change
821 impacts on marine ecosystems. *Annual review of marine science* 4:11-37

- 822 Donner W, Rodríguez H. 2008. Population composition, migration and inequality: The influence of
823 demographic changes on disaster risk and vulnerability. *Social forces* 87:1089-1114
- 824 Duke NC, Kovacs JM, Griffiths AD, Preece L, Hill DJE, van Oosterzee P, Mackenzie J, Morning HS, Burrows
825 D. 2017. Large-scale dieback of mangroves in Australia's Gulf of Carpentaria: a severe ecosystem
826 response, coincidental with an unusually extreme weather event. *Marine and Freshwater Research*
827 68:1816-1829
- 828 Dunham JB, Young MK, Gresswell RE, Rieman BE. 2003. Effects of fire on fish populations: landscape
829 perspectives on persistence of native fishes and nonnative fish invasions. *Forest Ecology and*
830 *Management* 178:183-196
- 831 Dyer KR. 2021. Response of estuaries to climate change. In Eisma, D. (Ed.), *Climate Change: Impact on*
832 *Coastal Habitation*. CRC Press, Boca Raton, pp. 85–110
- 833 Einsporn S, Broeg K, Köhler A. 2005. The Elbe flood 2002—toxic effects of transported contaminants in
834 flatfish and mussels of the Wadden Sea. *Marine Pollution Bulletin* 50:423-429
- 835 Ellsworth WL. 2013. Injection-induced earthquakes. *Science* 341:1225942
- 836 Eriksson H, Albert J, Albert S, Warren R, Pakoa K, Andrew N. 2017. The role of fish and fisheries in
837 recovering from natural hazards: Lessons learned from Vanuatu. *Environmental Science & Policy* 76: 50-
838 58
- 839 Evenden MD. 2004. *Fish Versus Power: An Environmental History of the Fraser River*. Cambridge
840 University Press, New York. 309 pp
- 841 Fabricius KE, De'Ath G, Puotinen ML, Done T, Cooper TF, Burgess SC. 2008. Disturbance gradients on
842 inshore and offshore coral reefs caused by a severe tropical cyclone. *Limnology and Oceanography*
843 53:690-704
- 844 Fattorini S, Di Lorenzo T, Galassi DMP. 2018. Earthquake impacts on microcrustacean communities
845 inhabiting groundwater-fed springs alter species-abundance distribution patterns. *Sci Rep.* 8:1501
- 846 Fattorini S, Lombardo P, Fiasca B, Di Cioccio A, Di Lorenzo T, Galassi DMP. 2017. Earthquake-Related
847 Changes in Species Spatial Niche Overlaps in Spring Communities. *Sci Rep.* 7:443
- 848 Fitak RR, Wheeler BR, Johnsen S. 2020. Effect of a magnetic pulse on orientation behavior in rainbow
849 trout (*Oncorhynchus mykiss*). *Behavioural Processes* 172:104058
- 850 Flores BM, Piedade MTF, Nelson BW. 2014. Fire disturbance in Amazonian blackwater floodplain forests.
851 *Plant Ecology & Diversity* 7:319-327
- 852 Folt CL, Chen CY, Moore MV, Burnaford J. 1999. Synergism and antagonism among multiple stressors.
853 *Limnology and Oceanography* 44: 864-877
- 854 Freund F, Stolc V. 2013. Nature of pre-earthquake phenomena and their effects on living organisms.
855 *Animals* 3(2):513-531.
- 856 Fry FEJ. 1971. The effect of environmental factors on the physiology of fish. In *Fish Physiology* Vol. 6 (ed.
857 Hoar W. S., Randall D. J.), New York: Academic Press 1-98 pp.

- 858 Galassi D, Lombardo P, Fiasca B, et al. 2014. Earthquakes trigger the loss of groundwater biodiversity.
859 *Scientific Reports* 4:6273
- 860 Galassi DMP, De Laurentiis P, Dole-Olivier MJ. 1999. Phylogeny and biogeography of the genus
861 *Pseudectinosoma*, and description of *P. janineae* sp. n. (Crustacea, Copepoda, Ectinosomatidae).
862 *Zoologica Scripta* 28:289-303
- 863 Gallo-Reynoso JP, Égido-Villarreal J, Martínez-Villalba G. 2011. Reaction of fin whales Balaenoptera
864 physalus to an earthquake. *Bioacoustics* 20:317-329
- 865 Geertsema M, Highland L, Vaugeouis L. 2009. Environmental impact of landslides. In: Sassa K, Canuti P
866 (Eds) Landslides—disaster risk reduction. Springer, Berlin, pp 589–607
- 867 Geertsema M, Menounos B, Bullard G, Carrivick JL, Clague JJ, Dai C, et al. 2022. The 28 November 2020
868 landslide, tsunami, and outburst flood—a hazard cascade associated with rapid deglaciation at Elliot
869 Creek, British Columbia, Canada. *Geophysical Research Letters* 49(6): e2021GL096716
- 870 Geertsema M, Pojar JJ. 2007. Influence of landslides on biophysical diversity—a perspective from British
871 Columbia. *Geomorphology* 89(1-2):55-69.
- 872 Giere O. 2009 *Meiobenthology: the microscopic motile fauna of aquatic sediments*. 2nd edition.
873 Springer-Verlag, Berlin 1-527 pp.
- 874 Gillanders BM, Kingsford MJ. 2002. Impact of changes in flow of freshwater on estuarine and open
875 coastal habitats and the associated organisms. *Oceanography and Marine Biology: An Annual Review*
876 40:233-309
- 877 Gillett N, Cannon A, Malinina E, Schnorbus M, Anslow F, Sun Q, Kirchmeier-Young M, et al. 2022. Human
878 Influence on the 2021 British Columbia Floods. Available at SSRN: <https://ssrn.com/abstract=4025205> or
879 <http://dx.doi.org/10.2139/ssrn.4025205>.
- 880 Glassic HC, Gaeta JW. 2019. Littoral habitat loss caused by multiyear drought and the response of an
881 endemic fish species in a deep desert lake. *Freshwater Biology* 64:421-432
- 882 Golovanova IL, Philippov AA, Chebotareva YV, Izyumov YG, Krylov VV. 2015. Impact of simulated
883 geomagnetic storm on activity of digestive glycosidases in roach *Rutilus rutilus* underyearlings. *Journal of*
884 *Ichthyology* 55:590-595
- 885 Gomez Isaza DF, Cramp RL, Franklin CE. 2022. Fire and rain: A systematic review of the impacts of
886 wildfire and associated runoff on aquatic fauna. *Global Change Biology* 28(8): 2578-2595
- 887 Gonzalez-Martin C, Teigell-Perez N, Valladares B, Griffin DW. 2014. The global dispersion of pathogenic
888 microorganisms by dust storms and its relevance to agriculture. *Advances in Agronomy* 127:1-41
- 889 Goudie AS. 2009. Dust storms: Recent developments. *Journal of Environmental Management* 90:89-94
- 890 Green J, Kling GW. 1988. The genus *Daphnia* in Cameroon, West Africa. *Hydrobiologia* 160:257–261
- 891 Greening H, Doering P, Corbett C. 2006. Hurricane impacts on coastal ecosystems. *Estuaries and Coasts*
892 29:877–879

- 893 Gresswell RE. 1999. Fire and aquatic ecosystems in forested biomes of North America. *Transactions of*
894 *the American fisheries society* 128:193-221
- 895 Griffin DW, Kellogg CA. 2004. Dust storms and their impact on ocean and human health: dust in Earth's
896 atmosphere. *EcoHealth* 1(3): 284-295.
- 897 Gudger EW. 1929. I.—More rains of fishes. *Annals and Magazine of Natural History* 3:1-26
- 898 Guerra M, Dawson S, Sabadel A, Slooten E, Somerford T, Williams R, Wing L, Rayment W. 2020 Changes
899 in habitat use by a deep-diving predator in response to a coastal earthquake, Deep Sea Research Part I:
900 *Oceanographic Research Papers* 158:10322
- 901 Guha-Sapir D, Santos I, Borde A. 2013. The economic impacts of natural disasters. Oxford University
902 Press, NY.
- 903 Gunawardena J, Ziyath AM, Bostrom TE, Bekessy LK, Ayoko GA, Egodawatta P, Goonetilleke A. 2013.
904 Characterisation of atmospheric deposited particles during a dust storm in urban areas of Eastern
905 Australia. *Science of the Total Environment* 461:72-80
- 906 Gupta A, Fox H. 1974. Effects of high-magnitude floods on channel form: A case study in Maryland
907 Piedmont. *Water Resources Research* 10:499-509
- 908 Gurnell AM, Gregory KJ, Petts GE. 1995. The role of coarse woody debris in forest aquatic habitats:
909 implications for management. *Aquatic Conservation: Marine and Freshwater Ecosystems* 5143-166
- 910 Gutowsky LFG, Rider MJ, Roemer RP, Gallagher AJ, Heithaus M. R, Cooke SJ, Hammerschlag N. 2021.
911 Large sharks exhibit varying behavioral responses to major hurricanes. *Estuarine, Coastal and Shelf*
912 *Science* 256:107373
- 913 Hallegraef G, Coman F, Davies C, Hayashi A, McLeod D, Slotwinski A, Richardson AJ. 2014. Australian
914 dust storm associated with extensive *Aspergillus sydowii* fungal "bloom" in coastal waters. *Applied and*
915 *Environmental Microbiology* 80:3315-3320
- 916 Hapke C, Dallas K, Green K. 2003. Estimated sediment yield from coastal landslides and active slope
917 distribution along the Big Sur coast and Addendum: Coastal cliff Erosion Rates, Big Sur, CA. A Report of
918 Findings for the Coastal Highway Management Plan, California Department of Transportation, District, 5.
- 919 Harding S, Jellyman PG. 2015. Earthquakes, catastrophic sediment additions and the response of urban
920 stream communities. *Marine and Freshwater Research* 49:346-355
- 921 Harmon ME, Franklin JF, Swanson FJ, Sollins P, Gregory SV, Lattin JD, Cummins KW. 1986. Ecology of
922 coarse woody debris in temperate ecosystems. *Advances in Ecological Research* 15:133-302
- 923 Hartman GF, Scrivener JC, Miles MJ. 1996. Impacts of logging in Carnation Creek, a high-energy coastal
924 stream in British Columbia, and their implication for restoring fish habitat. *Canadian Journal of Fisheries*
925 *and Aquatic Sciences* 53:237-251
- 926 Hauer C, Habersack H. 2009. Morphodynamics of a 1000-year flood in the Kamp River, Austria, and
927 impacts on floodplain morphology. *Earth Surface Processes and Landforms* 34:654-682

- 928 Havel JE, Shurin JB. 2004. Mechanisms, effects, and scales of dispersal in freshwater zooplankton.
929 *Limnology and Oceanography* 49:1229-1238
- 930 Hayes SK, Montgomery DR, Newhall CG. 2002. Fluvial sediment transport and deposition following the
931 1991 eruption of Mount Pinatubo. *Geomorphology* 45(3-4): 211-224
- 932 Hernández P A, Nogami K, Padrón E, Somoza L, Amonte C, Mori T, Melián GV, Sumino H, Kikawada Y,
933 Pérez Nemesio M. 2021. Hydrochemical and Hydroacoustic Investigation of the Yugama Acid Crater
934 Lake, Kusatsu-Shirane, Japan. *Frontiers in Earth Science* 9:1180
- 935 Heupel MR, Simpfendorfer CA, Hueter RE, 2003. Running before the storm: blacktip sharks respond to
936 falling barometric pressure associated with Tropical Storm Gabrielle. *Journal of Fish Biology* 63: 1357-
937 1363
- 938 Hinch SG, Diewert RE, Lissimore TJ, Prince AM, Healey MC, Henderson MA 1996. Use of electromyogram
939 telemetry to assess difficult passage areas for river-migrating adult sockeye salmon. *Transactions of the*
940 *American Fisheries Society* 125:253-260
- 941 Hirslund F. 2020. A single limnic eruption at the origin of today's large-scale density structure of Lake
942 Kivu. *African Earth Sciences* 161:103614
- 943 Hirst AJ, Longmore AR, Ball D, Cook PLM, Jenkins GP. 2016. Linking nitrogen sources utilised by seagrass
944 in a temperate marine embayment to patterns of seagrass change during drought. *Marine Ecology*
945 *Progress Series* 549:79-88
- 946 Hobbs WO, Wolfe AP. 2008. Recent paleolimnology of three lakes in the Fraser River Basin (BC, Canada):
947 no response to the collapse of sockeye salmon stocks following the Hells Gate landslides. *Journal of*
948 *Paleolimnology* 40:295-308
- 949 Hoegh-Guldberg O. 1999. Climate change, coral bleaching and the future of the world's coral reefs.
950 *Marine and Freshwater Research* 50(8): 839-866
- 951 Houlton BZ, Driscoll CT, Fahey TJ, Likens GE, Groffman PM, Bernhardt ES, Buso DC. 2003. Nitrogen
952 dynamics in ice storm-damaged forest ecosystems: implications for nitrogen limitation theory.
953 *Ecosystems* 6:431-43
- 954 Hughes ALH, Wilson AM, Morris JT. 2012. Hydrologic variability in a salt marsh: Assessing the links
955 between drought and acute marsh dieback. *Estuarine Coastal and Shelf Science* 111:95-106
- 956 Ingebritsen SE, Manga M. 2019. Earthquake hydrogeology. *Water Resources Research* 55:5212- 5216
- 957 Jakob M. 2000. The impacts of logging on landslide activity at Clayoquot Sound, British Columbia. *Catena*
958 38:279-300
- 959 James FH. 1894. Raining Worms and Frogs. *Science* 65-65
- 960 Jáuregui E. 1989. The dust storms of Mexico City. *International Journal of Climatology* 9:169-180
- 961 Johannesson T. 2001. Run-up of two avalanches on the deflecting dams at Flateyri, northwestern
962 Iceland. *Annals of Glaciology* 32:350-354

- 963 Johnson A L, Smith DJ. 2010. Geomorphology of snow avalanche impact landforms in the southern
964 Canadian Cordillera. *Canadian Geographer* 54:87–103
- 965 Johnson EA. 1987. The relative importance of snow avalanche disturbance and thinning on the growth-
966 habit of canopy plant populations. *Ecology* 68:43-53
- 967 Kallis G. 2008. Droughts. *Annual Review of Environment and Resources* 33:85-118
- 968 Kämäräinen M, Hyvärinen O, Vajda A, Nikulin G, Meijgaard, EV, Teichmann C, Jacob D, Gregow H, Jylhä
969 K. 2018. Estimates of present-day and future climatologies of freezing rain in Europe based on CORDEX
970 regional climate models. *Journal of Geophysical Research: Atmospheres*. 123
- 971 Kaur H, Habibullah MS, Nagaratnam S. 2019. Impact of natural disasters on biodiversity: evidence using
972 quantile regression approach. *Jurnal Ekonomi Malaysia* 53:67-81
- 973 Kemter M, Fischer M, Luna LV, Schönfeldt E, Vogel J, Banerjee A, Thonicke K. 2021. Cascading hazards in
974 the aftermath of Australia's 2019/2020 Black Summer wildfires. *Earth's Future* 9:1884
- 975 Kennish MJ. 2002. Environmental threats and environmental future of estuaries. *Environmental*
976 *Conservation* 29(1): 78-107.
- 977 King VD VO. 1934. The Mendoza River flood of 10-11 January 1934-Argentina. *The Geographical Journal*
978 84:321-326
- 979 Kitahashi T, Jenkins RG, Nomaki H, Shimanaga M, Fujikura K, Kojima S. 2014. Effect of the 2011 Tohoku
980 Earthquake on deep-sea meiofaunal assemblages inhabiting the landward slope of the Japan Trench.
981 *Marine Geology* 358:128-137
- 982 Klein RJ, Nicholls RJ, Thomalla F. 2003. Resilience to natural hazards: How useful is this concept?. Global
983 environmental change part B: *environmental hazards* 5:35-45
- 984 Kling G. W., Evans W. C., Tanyileke G., Kusakabe M., Ohba T., Yoshida Y., Hell J V. (2005) Degassing Lakes
985 Nyos and Monoun: Defusing certain disaster. *Proceedings of the National Academy of Sciences*
986 102:14185-14190
- 987 Koizumi I, Kanazawa Y, Tanaka Y. 2013. The fishermen were right: experimental evidence for tributary
988 refuge hypothesis during floods. *Zoological Science* 30:375-379
- 989 Kraft CE, Schneider RL, Warren DR. 2002. Ice storm impacts on woody debris and debris dam formation
990 in northeastern U.S. streams. *Canadian Journal of Fisheries and Aquatic Sciences* 59:1677-1684
- 991 Kreuzweiser DP, Hazlett PW, Gunn JM. 2008. Logging impacts on the biogeochemistry of boreal forest
992 soils and nutrient export to aquatic systems: A review. *Environmental Reviews* 16:157-179
- 993 Krylov VV. 2017. Biological effects related to geomagnetic activity and possible mechanisms.
994 *Bioelectromagnetics* 38:497-510
- 995 Krylov VV, Chebotareva YV, Izyumov YG. 2019. Delayed consequences of the influence of simulated
996 geomagnetic storms on roach *Rutilus rutilus* embryos. *Journal of Fish Biology* 95:1422-1429

- 997 Krylov VV, Osipova EA. 2019. The response of *Daphnia magna* Straus to long-term exposure to simulated
998 geomagnetic storms. *Life Sciences in Space Research* 21:83-88
- 999 Krylov VV, Osipova EA, Pankova NA, Talikina MG, Chebotareva YV, Izyumov YG, Nepomnyashchikh VA.
1000 2017. The effect of a temporal shift in diurnal geomagnetic variation on roach *Rutilus rutilus* L. embryos:
1001 a comparison with effects of simulated geomagnetic storms. *Biophysics* 62:675-681
- 1002 Kumar AB, Raj S, Arjun CP, Katwate U, Raghavan R. 2019. Jurassic invaders: flood-associated occurrence
1003 of arapaima and alligator gar in the rivers of Kerala. *Current Science* 116: 1628-1630
- 1004 Kumasaki M, King M, Arai M, Yang L. 2016. Anatomy of cascading natural disasters in Japan: main modes
1005 and linkages. *Natural Hazards* 80:1425-1441
- 1006 Kurita Y, Uehara S, Okazaki Y, Sakami T, Nambu R, Tomiyama T. 2017. Impact of the great tsunami in
1007 2011 on the quality of nursery grounds for juvenile Japanese flounder *Paralichthys olivaceus* in Sendai
1008 Bay, Japan. *Fisheries Oceanography* 26(2): 165-180
- 1009 Kusakabe M. 2017. Lakes Nyos and Monoun gas disasters (Cameroon)—Limnic eruptions caused by
1010 excessive accumulation of magmatic CO₂ in crater lakes. *Geochemical Monograph Series* 1:1–50
- 1011 Lake PS. 2011. Drought and aquatic ecosystems: effects and responses. John Wiley & Sons, NY.
- 1012 Lake PS 2003. Ecological effects of perturbation by drought in flowing waters. *Freshwater Biology*
1013 48(7):1161-1172
- 1014 Lanka RP, Hubert WA, Wesche TA. 1987. Relations of geomorphology to stream habitat and trout
1015 standing stock in small Rocky Mountain streams. *Transactions of the American Fisheries Society* 116:21–
1016 28
- 1017 Lawrence J, Blackett P, Cradock-Henry NA. 2020. Cascading climate change impacts and implications.
1018 *Climate Risk Management* 29:100234
- 1019 Lennox RJ, Crook DA, Moyle PB, Struthers DP, Cooke SJ. 2019. Toward a better understanding of
1020 freshwater fish responses to an increasingly drought-stricken world. *Reviews in Fish Biology and*
1021 *Fisheries* 29:71-92
- 1022 Liu H, Gilmartin J, Li C, Li K. 2021. Detection of time-varying pulsed event effects on estuarine pelagic
1023 communities with ecological indicators after catastrophic hurricanes. *Ecological Indicators* 123: 107327
- 1024 Lockridge PA. 1990. Nonseismic phenomena in the generation and augmentation of tsunamis. *Natural*
1025 *Hazards* 3(4): 403-412
- 1026 Logan RL, Schuster RL. 1991. Lakes divided – the origin of Lake Sutherland and Lake Crescent, Clallam
1027 County, Washington. *Wash. Geol., Div. of Geol. and Earth Resources, Olympia.* 19(1):38-42
- 1028 Lubick N. Earthquakes from the ocean: Danger zones. *Nature* 476:391–392
- 1029 Luckman BH. 1978. Geomorphic Work of Snow Avalanches in the Canadian Rocky Mountains. *Arctic and*
1030 *Alpine Research* 10:261-276

- 1031 Lynch AJ, Cooke SJ, Deines AM, Bower SD, Bunnell DB, Cowx IG, Beard Jr TD. 2016. The social, economic,
1032 and environmental importance of inland fish and fisheries. *Environmental Reviews* 24:115-121
- 1033 Mackey BH, Roering JJ, Lamb MP. 2011. Landslide-dammed paleolake perturbs marine sedimentation
1034 and drives genetic change in anadromous fish. *Proceedings of the National Academy of Sciences*
1035 108:18905-18909
- 1036 Major JJ, Mosbrucker AR, Spicer KR. 2018. Sediment erosion and delivery from Toutle River basin after
1037 the 1980 eruption of Mount St. Helens: A 30-year perspective, in C.M. Crisafulli and V.H. Dale (eds.)
1038 Ecological Responses at Mount St. Helens: Revisited 35 years after the 1980 Eruption, Springer
1039 Science+Business, New York. 19-44 pp.
- 1040 Majumdar SD, Hazra S, Giri S, Chanda A, Gupta K, Mukhopadhyay A, Roy SD. 2018. Threats to coral reef
1041 diversity of Andaman Islands, India: A review. *Regional Studies in Marine Science* 24: 237-250
- 1042 Mallin MA, Corbett CA. 2006. How hurricane attributes determine the extent of environmental effects:
1043 multiple hurricanes and different coastal systems. *Estuaries and Coasts* 29(6): 1046-1061
- 1044 Massie JA, Strickland BA, Santos RO, Hernandez J, Viadero N, Boucek RE, Willoughby H, Heithaus MR,
1045 Rehage JS. 2020. Going Downriver: Patterns and cues in hurricane-driven movements of common snook
1046 in a subtropical coastal river. *Estuaries and Coasts* 1158-1173 pp.
- 1047 Matthews WJ, Marsh-Matthews E. 2003. Effects of drought on fish across axes of space, time and
1048 ecological complexity. *Freshwater Biology* 48:1232-1253
- 1049 McKee KL, Mendelssohn IA, Materne MD. 2004. Acute salt marsh dieback in the Mississippi River deltaic
1050 plain: a drought-induced phenomenon? *Global Ecology and Biogeography* 13:65-73
- 1051 Merz B, Blöschl G, Vorogushyn S, Dottori F, Aerts JC, Bates P, et al. 2021. Causes, impacts and patterns
1052 of disastrous river floods. *Nature Reviews Earth & Environment* 2(9): 592-609
- 1053 Millward AA, Kraft CE, Warren DR. 2010. Ice-storm damage greater along the terrestrial-aquatic
1054 interface in forested landscapes. *Ecosystems* 13:249–260
- 1055 Milton SL, Leone-Kabler S, Schulman AA, Lutz PL. 1994. Effects of Hurricane Andrew on the sea turtle
1056 nesting beaches of South Florida. *Bulletin of Marine Science* 974-981 pp.
- 1057 Minshall GW. 2003. Responses of stream benthic macroinvertebrates to fire. *Forest Ecology and*
1058 *Management* 178:155-161
- 1059 Mitchell JK. 1988. Confronting natural disasters: An international decade for natural hazard reduction.
1060 *Environment: Science and Policy for Sustainable Development* 30:25-29
- 1061 Mohr CH, Manga M, Wang CY, Korup O. 2017. Regional changes in streamflow after a megathrust
1062 earthquake. *Earth and Planetary Science Letters* 458:418-428
- 1063 Monteil CL, Lefevre CT. 2020. Magnetoreception in microorganisms. *Trends in microbiology* 28:266-275
- 1064 Morgan G. 2012. An attempt to explain the falls of fish and other strange objects from the sky. *Weather*
1065 67:87-87

- 1066 Mountjoy JJ, Howarth JD, Orpin AR, Barnes PM, Bowden DA, Rowden AA, Schimel ACG, Holden C,
1067 Horgan HJ, Nodder SD, Patton JR, Lamarche G, Gerstenberger M, Micallef A, Pallentin A, Kane T. 2018.
1068 *Sci. Adv.* 4:374
- 1069 Muller KA, Straub PN. 2016. A review of avalanche ecology: Forest habitat structure and wildlife
1070 biodiversity. In Proceedings, International Snow Science Workshop, Breckenridge, Colorado (pp. 1389-
1071 1391) Available from: https://arc.lib.montana.edu/snow-science/objects/ISSW16_P4.52.pdf
- 1072 Muraoka D, Tamaki H, Takami H, Kurita Y, Kawamura T. 2017. Effects of the 2011 Great East Japan
1073 Earthquake and tsunami on two kelp bed communities on the Sanriku coast. *Fisheries Oceanography*
1074 26(2): 128-140
- 1075 Muraveiko VM, Stepanyuk IA, Zenzerov VS. 2013. The response of the crab *Paralithodes camtschaticus*
1076 (*Tilesius*, 1815) to geomagnetic storms. Springer Nature BV. In *Doklady Biological Sciences* 10:448
- 1077 Murphy I, Johnson S, Hatfield T. 2020b. Big Bar Landslide Southern Endowment Fund Science Workshop
1078 Summary. Consultant's report prepared for Pacific Salmon Commission and Fisheries and Oceans
1079 Canada by Ecofish Research Ltd.
- 1080 Murphy SF, McCleskey RB, Martin DA, Holloway JM, Writer JH, 2020a. Wildfire-driven changes in
1081 hydrology mobilize arsenic and metals from legacy mine waste. *Science of the Total Environment*
1082 743:140635
- 1083 Nagrodski A, Raby GD, Hasler CT, Taylor MK, Cooke SJ. 2012. Fish stranding in freshwater systems:
1084 sources, consequences, and mitigation. *Journal of environmental management* 103:133-141
- 1085 Nakaoka M, Tanaka Y, Mukai H, Suzuki T, Aryuthaka C. 2007. Tsunami impacts on biodiversity of
1086 seagrass communities in the Andaman Sea, Thailand:(1) Seagrass abundance and diversity. *Publications*
1087 *of the Seto Marine Biological Laboratory, Special Publication Series* 8:49-56
- 1088 Nana P, Nola M, Bricheux G, Fokam Z, Ngassam P, Enah D, Colombet J, Vellet A, Mone A, Ravet V,
1089 Debros D, Sime-Ngando T. 2020. Diversity and Structure of the Prokaryotic Communities Indigenous to
1090 Two Volcanic Lakes: Nyos and Monoun in Cameroon. *Open Journal of Ecology* 10:632-650
- 1091 Narimatsu Y, Shibata Y, Hattori T, Yano T, Nagao J. 2017. Effects of a marine-protected area occurred
1092 incidentally after the Great East Japan Earthquake on the Pacific cod (*Gadus macrocephalus*) population
1093 off northeastern Honshu, Japan. *Fisheries Oceanography* 26(2): 181-192
- 1094 Nayar A. 2009. A lakeful of trouble: Africa's Lake Kivu contains vast quantities of gas, which makes it
1095 both dangerous and valuable. Anjali Nayar asks whether it is possible to tap the gas without causing a
1096 disaster. *Nature* 460:7253
- 1097 Nelson JL, Groninger JW, Battaglia LL, Ruffner CM. 2008. Bottomland hardwood forest recovery
1098 following tornado disturbance and salvage logging. *Forest Ecology and Management* 256:388-395
- 1099 Nichols CR, Zinnert J, Young DR. 2019. Degradation of coastal ecosystems: causes, impacts and
1100 mitigation efforts. In L. D. Wright & C. R. Nichols (Eds.), *Tomorrow's coasts: complex and impermanent*
1101 (pp. 119–136). Springer Nature.

- 1102 Niemi GJ, DeVore P, Detenbeck N, Taylor D, Lima A, Pastor J, et al. 1990. Overview of case studies on
1103 recovery of aquatic systems from disturbance. *Environmental Management* 14(5):571-587
- 1104 Ning L, Bradley RS. 2015. Snow occurrence changes over the central and eastern United States under
1105 future 36 warming scenarios. *Scientific Reports* 5:17073
- 1106 Oakley JW, Guillen GJ. 2020. Impact of hurricane Harvey on Galveston bay saltmarsh nekton
1107 communities. *Estuaries and Coasts* 43:984-992.
- 1108 Oaks SD, Bender SO. 1990. Hazard reduction and everyday life: Opportunities for integration during the
1109 decade for natural disaster reduction. *Natural Hazards* 3:87-89
- 1110 Oczkowski AJ, Nixon SW, Granger SL, El-Sayed AFM, McKinney RA. 2009. Anthropogenic enhancement of
1111 Egypt's Mediterranean fishery. *Proceedings of the National Academy of Sciences of the United States of*
1112 *America* 106:1364-1367
- 1113 Okazaki Y, Kurita Y, Uehara S. 2017. Limited effect of the massive tsunami caused by the 2011 Great
1114 East Japan Earthquake on the shallow sandy shore demersal fish assemblages in Sendai Bay. *Fisheries*
1115 *Oceanography* 26(2): 155-164
- 1116 Olgun N, Duggen S, Langmann B, Hort M, Waythomas CF, Hoffmann L, Croot P. 2013. Geochemical
1117 evidence of oceanic iron fertilization by the Kasatochi volcanic eruption in 2008 and the potential
1118 impacts on Pacific sockeye salmon. *Marine Ecology Progress Series* 488: 81-88
- 1119 Oliver J, Carney D, Okden J. 1999. Ecological Impacts of the Landslide Manipulations at the McWay Slide,
1120 Big Sur Coast, and Management Recommendations, Report to Caltrans (Calif. Dept. of Transp.), Moss
1121 Landing Marine Laboratories, Moss Landing, Calif., 40 pp.
- 1122 Orchard S, Hickford MJH, Schiel DR. 2018. Earthquake-induced habitat migration in a riparian spawning
1123 fish has implications for conservation management. *Aquatic Conservation* 28:702–712
- 1124 Ostrom E. 2009. A general framework for analyzing sustainability of social-ecological systems. *Science*
1125 325(5939): 419-422
- 1126 Patchett J, Samsel F, Tsai KC, Gisler GR, Rogers DH, Abram GD, Turton TL. 2016. Visualization and
1127 analysis of threats from asteroid ocean impacts. Los Alamos National Laboratory
- 1128 Peierls BL, Christian RR, Paerl HW. 2003. Water quality and phytoplankton as indicators of hurricane
1129 impacts on large estuarine ecosystems. *Estuaries* 26:1329–1343
- 1130 Pelling M. 2001. Natural disasters. Social nature: Theory, Practice, and Politics, Blackwell Publishers, Inc.,
1131 Malden, MA, 170-189 pp.
- 1132 Peterson CH, Lubchenco J. 1997. Marine ecosystem services. Island Press, Washington, DC. 177-195 pp.
- 1133 Pierce II GH. 2016. *Geomorphic response to tornado impact in Abrams Creek, Smoky Mountains National*
1134 *Park, Tennessee*. The University of Alabama. Master's thesis. 34 pp.
- 1135 Pigg J, Gibb R. 1998. Wind-caused fish kills. In *Proceedings of the Oklahoma Academy of Science* 78:117

- 1136 Pollack JB, Kim HC, Morgan EK, Montagna PA. 2011. Role of flood disturbance in natural oyster
1137 (*Crassostrea virginica*) population maintenance in an estuary in South Texas, USA. *Estuaries and Coasts*
1138 34:187-197
- 1139 Pulkkinen A, Moore K, Zellar R, Uritskaya O, Karaköylü E M, Uritsky V, Reeb D. 2020. Statistical analysis
1140 of the possible association between geomagnetic storms and cetacean mass strandings. *Journal of*
1141 *Geophysical Research: Biogeosciences* 10:125
- 1142 Rahav E, Paytan A, Chien CT, Ovadia G, Katz T, Herut B. 2016. The impact of atmospheric dry deposition
1143 associated microbes on the southeastern Mediterranean Sea surface water following an intense dust
1144 storm. *Frontiers in Marine Science* 3:127
- 1145 Rampino MR, Caldeira K, Prokoph A. 2019. What causes mass extinctions? Large asteroid/comet
1146 impacts, flood-basalt volcanism, and ocean anoxia—Correlations and cycles. *Geological Society of*
1147 *America Special Paper* 542:271-302
- 1148 Ranasinghe, R., Ruane, A. C., Vautard, R., Arnell, N., Coppola, E., Cruz, F. A., et al. (2021). Climate change
1149 information for regional impact and for risk assessment. In P. Zhai, A. Pirani, S. L. Connors, C. Péan, S.
1150 Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T.
1151 K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zhou (Eds.), *Climate change 2021: The physical science*
1152 *basis. Contribution of working group I to the sixth assessment report of the intergovernmental Panel on*
1153 *climate change [Masson-Delmotte, V.]*. Cambridge University Press, UK.
- 1154 Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PT, et al. 2019. Emerging threats and
1155 persistent conservation challenges for freshwater biodiversity. *Biological Reviews* 94(3): 849-873
- 1156 Rhoads AG, Hamburg SP, Fahey TJ, Siccama TG, Hane EN, Battles J, Cogbill C, Randall J, Wilson G. 2002.
1157 Effects of an intense ice storm on the structure of a northern hardwood forest. *Canadian Journal of*
1158 *Forest Research* 32:1763–1775
- 1159 Richardson SD, Reynolds JM. 2000. An overview of glacial hazards in the Himalayas. *Quaternary*
1160 *International* 65:31-47
- 1161 Ritchie H, Roser M. 2019. Natural disasters. Published online at OurWorldInData.org. Retrieved from
1162 <https://ourworldindata.org/naturaldisasters> [Online Resource]
- 1163 Roberts SK. 1999. Raining cats and frogs. *Weather* 54.4:126-126
- 1164 Rondeau D, Perry B, Grimard F. 2020. The consequences of COVID-19 and other disasters for wildlife and
1165 biodiversity. *Environmental and Resource Economics* 76:945-961
- 1166 Rouwet D, Tassi F, Mora-Amador R, Sandri L, Chiarini V. 2014. Past, present and future of volcanic lake
1167 monitoring. *Journal of Volcanology and Geothermal Research* 272:78-97
1168
- 1169 Ruiz-Villanueva V, Díez-Herrero A, Ballesteros JA, Bodoque JM. 2014. Potential large woody debris
1170 recruitment due to landslides, bank erosion and floods in mountain basins: a quantitative estimation
1171 approach. *River Research and Applications* 30:81-97

- 1172 Rytwinski T, Harper M, Taylor JJ, Bennett JR, Donaldson LA, Smokorowski K E, Cooke SJ. 2020. What are
1173 the effects of flow-regime changes on fish productivity in temperate regions? A systematic map.
1174 *Environmental Evidence* 9:1-26
- 1175 Sahil, Sood, SK. 2021. Scientometric analysis of natural disaster management research. *Natural Hazards*
1176 *Review* 22:04021008
- 1177 Sarmiento H, Isumbisho M, Descy J-P. 2006. Phytoplankton ecology of Lake Kivu (eastern Africa). *Journal*
1178 *of Plankton Research* 28:815–829
- 1179 Sathianandan TV, Mohamed KS, Vivekanandan E. 2012. Species diversity in fished taxa along the
1180 southeast coast of India and the effect of the Asian Tsunami of 2004. *Marine Biodiversity* 42(2):179-187
- 1181 Schuster RL, Chleborad AF, Hays WH. 1989. The White Bluffs landslides. in R. Galster (ed.), *Engineering*
1182 *Geology in Washington, Wash. Div. Geol. and Earth Resources Bull., Olympia.* 911-920 pp.
- 1183 Schuster RL, Highland LM. 2003. Impact of landslides and innovative landslide-mitigation measures on
1184 the natural environment. In *Proceedings of the International Conference on Slope Engineering,*
1185 *December 8–10, 2003, Hong Kong, China.*
- 1186 Schuster RL, Highland LM. 2007. Overview of the effects of mass wasting on the natural environment.
1187 *Environmental & Engineering Geoscience* 13:25-44
- 1188 Seddon N, Chausson A, Berry P, Girardin CA, Smith A, Turner B. 2020. Understanding the value and
1189 limits of nature-based solutions to climate change and other global challenges. *Philosophical*
1190 *Transactions of the Royal Society B* 375(1794):20190120
- 1191 Shanklin E. 1989. Exploding lakes and maleficent water in Grassfields legends and myth. *Journal of*
1192 *Volcanology and Geothermal Research* 39:233-246
- 1193 Shapley M, Finney B, Kruger C. 2019. Characteristics of landslide-formed lakes of central Idaho: high-
1194 resolution archives of watershed productivity and clastic sediment delivery. In: Starrett, S., Rosen, M.
1195 (Eds.) *From Saline to Freshwater: The Diversity of Western Lakes in Space and Time.* The Geological
1196 Society of America, NY, pp. 241–258.
- 1197 Shi JH, Zhang J, Gao HW, Tan SC, Yao XH, Ren JL. 2013. Concentration, solubility and deposition flux of
1198 atmospheric particulate nutrients over the Yellow Sea. *Deep Sea Research Part II: Topical Studies in*
1199 *Oceanography* 97:43-50
- 1200 Shoji J, Morimoto M. 2016. Changes in fish community in seagrass beds in Mangoku-ura Bay from 2009
1201 to 2014, the period before and after the tsunami following the 2011 off the Pacific coast of Tohoku
1202 earthquake. *Journal of Oceanography* 72(1):91-98
- 1203 Shoji J, Yoshikawa K, Tomiyama T, Kawamura T. 2021. Temporal changes of the fish community in
1204 seagrass beds in Funakoshi and Otsuchi bays after habitat destruction caused by a tsunami in 2011.
1205 *Fisheries Science* 87(6): 827-836
- 1206 Silliman BR, van de Koppel J, Bertness MD, Stanton LE, Mendelssohn IA. 2005. Drought, snails, and large-
1207 scale die-off of southern US salt marshes. *Science* 310:1803-1806

- 1208 Silva LG, Doyle KE, Duffy D, Humphries P, Horta A, Baumgartner LJ. 2020. Mortality events resulting from
1209 Australia's catastrophic fires threaten aquatic biota. *Global Change Biology* 26:5345-5350
- 1210 Smith EP. 2002. BACI design. *Encyclopedia of Environmetrics* 1:141-148
- 1211 Smith N. 2006. There's no such thing as a natural disaster. Understanding Katrina: Perspectives from the
1212 social sciences. Social Science Research Council. Retrieved on January 31, 2022 from
1213 [http://blogs.ubc.ca/naturalhazards/files/2016/03/Smith-There%E2%80%99s-No-Such-Thing-as-a-](http://blogs.ubc.ca/naturalhazards/files/2016/03/Smith-There%E2%80%99s-No-Such-Thing-as-a-Natural-Disaster.pdf)
1214 [Natural-Disaster.pdf](http://blogs.ubc.ca/naturalhazards/files/2016/03/Smith-There%E2%80%99s-No-Such-Thing-as-a-Natural-Disaster.pdf)
- 1215 Smith W. 2013. Amphibians and large, infrequent forest disturbances: an extreme wind event facilitates
1216 habitat creation and anuran breeding. *Herpetological Conservation and Biology* 8:732-740
- 1217 Somov BV. 1991. Physical processes in solar flares (Vol. 172). Springer Science & Business Media.
- 1218 Stelzer JAA, Mesman JP, Adrian R, Ibelings BW. 2021. Early warning signals of regime shifts for aquatic
1219 systems: Can experiments help to bridge the gap between theory and real-world application? *Ecological*
1220 *Complexity* 47:100944
- 1221 Stovall JP, Keeton WS, Kraft CE. 2009. Riparian forest-stream interactions: variability of forest structure,
1222 light, and periphyton along Adirondack streams. *Canadian Journal of Forestry Research* 39:2343–2354
- 1223 Swanson FJ, Major JJ. 2005. Physical events, environments, and geological–ecological interactions at
1224 Mount St. Helens—March 1980–2004. In: Dale VH, Swanson FJ, Crisafulli CM (Eds) Ecological responses
1225 to the 1980 eruption of Mount St. Helens. Springer, New York, pp 27–44
- 1226 Swanson FJ, Crisafulli CM. 2018. Volcano ecology: state of the field and contributions of Mount St.
1227 Helens research. In: Crisafulli CM, Dale VH (eds) Ecological responses at Mount St Helens: revisited 35
1228 years after the 1980 eruption. Springer, New York, pp 305–323
- 1229 Talbot CJ, Bennett EM, Cassell K, Hanes DM, Minor EC, Paerl H et al. 2018. The impact of flooding on
1230 aquatic ecosystem services. *Biogeochemistry* 141(3): 439-461
- 1231 Tapia J, Villagra F, Bertrán C, Espinoza J, Focardi S, Fierro P, et al. 2019. Effect of the earthquake-
1232 tsunami (Chile, 2010) on toxic metal content in the Chilean abalone mollusc *Concholepas concholepas*.
1233 *Ecotoxicology and Environmental Safety* 169:418-424
- 1234 Tassi F, Fazi S, Rossetti S, Pratesi P, Ceccotti M, Cabassi J, et al. 2018. The biogeochemical vertical
1235 structure renders a meromictic volcanic lake a trap for geogenic CO₂ (Lake Averno, Italy). *PLoS ONE*
1236 13:193914
- 1237 Thomsen MS, Metcalfe I, Siciliano A, South PM, Gerrity S, Alestra T, Schiel DR. 2020. Earthquake-driven
1238 destruction of an intertidal habitat cascade. *Aquatic Botany* 164:103217
- 1239 Tilloy A, Malamud BD, Winter H, Joly-Laugel A. 2019. A review of quantification methodologies for multi-
1240 hazard interrelationships. *Earth-Science Reviews* 196:102881
- 1241 Tittensor DP, Mora C, Jetz W, Lotze HK, Ricard D, Berghe EV, Worm B. 2010. Global patterns and
1242 predictors of marine biodiversity across taxa. *Nature* 466:1098-1101

- 1243 Toon OB, Zahnle K, Morrison D, Turco RP, Covey C. 1997. Environmental perturbations caused by the
1244 impacts of asteroids and comets. *Reviews of Geophysics* 35:1-78
- 1245 Tuckett QM, Koetsier P. 2018. Post-fire debris flows delay recovery and create novel headwater stream
1246 macroinvertebrate communities. *Hydrobiologia* 814:161-174
- 1247 Turner BA. 1976. The development of disasters—a sequence model for the analysis of the origins of
1248 disasters. *The Sociological Review* 24(4):753-774
- 1249 United Nations Environment Programme. 2022. Spreading like Wildfire – The Rising Threat of
1250 Extraordinary Landscape Fires. A UNEP Rapid Response Assessment. Nairobi.
- 1251 Unmack PJ. 2001. Biogeography of Australian freshwater fishes. *Journal of Biogeography* 28:1053-1089
- 1252 Urabe J, Nakashizuka T. 2016. Ecological impacts of tsunamis on coastal ecosystems. Springer, Japan.
- 1253
- 1254 Van Aalst MK. 2006. The impacts of climate change on the risk of natural disasters. *Disasters* 30:5-18
- 1255 Vannote RL, Minshall WG, Cummins KW, Sedell JR, Cushing CE. 1980. The river continuum concept.
1256 *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-37
- 1257 Vasskog K, Nesje A, Støren E, Waldmann N, Chapron E, Ariztegui D. 2011. A Holocene record of snow-
1258 avalanche and flood activity reconstructed from a lacustrine sedimentary sequence in Oldevatnet,
1259 western Norway. *The Holocene* 21:597-614
- 1260 Verkaik I, Rieradevall M, Cooper SD, Melack JM, Dudley TL, Prat N. 2013. Fire as a disturbance in
1261 Mediterranean climate streams. *Hydrobiologia* 719:353-382
- 1262 Waide RB. 1991. Summary of the response of animal populations to hurricanes in the Caribbean.
1263 *Biotropica* 23(4):508-512
- 1264 Walker LR, Shiels AB. 2012. Landslide ecology. Cambridge University Press
- 1265 Wang X, Wang W, Tong C. 2016. A review on impact of typhoons and hurricanes on coastal wetland
1266 ecosystems. *Acta Ecologica Sinica* 36(1): 23-29
- 1267 Wang CY, Manga M. 2010a. Lecture Notes in Earth Sciences, Earthquakes and Water. Springer-Verlag
1268 Berlin Heidelberg 2010
- 1269 Wang CY, Manga M. 2010b. Hydrologic responses to earthquakes and a general metric. *Geofluids*
1270 10:206-216
- 1271 Warren DR, Collins SM, Purvis EM, Kaylor MJ, Bechtold HA. 2017. Spatial variability in light yields
1272 colimitation of primary production by both light and nutrients in a forested stream ecosystem.
1273 *Ecosystems* 20:198-210
- 1274 Warren DR, Kraft CE. 2003. Brook trout (*Salvelinus fontinalis*) response to wood removal from high-
1275 gradient streams of the Adirondack Mountains (N.Y., U.S.A.). *Canadian Journal of Fisheries and Aquatic*
1276 *Sciences* 60:379-389

- 1277 West AJ, Lin CW, Lin TC, Hilton RG, Liu SH, Chang CT, Hovius N. 2011. Mobilization and transport of
1278 coarse woody debris to the oceans triggered by an extreme tropical storm. *Limnology and*
1279 *Oceanography* 56:77-85
- 1280 Wetz MS, Yoskowitz DW. 2013. An 'extreme' future for estuaries? Effects of extreme climatic events on
1281 estuarine water quality and ecology. *Marine Pollution Bulletin* 69:7-18
- 1282 Wilson SS, Dunton K. 2018. Hypersalinity during regional drought drives mass mortality of the seagrass
1283 *Syringodium filiforme* in a subtropical lagoon. *Estuaries and Coasts* 41:855-865
- 1284 Wiltschko W, Wiltschko R. 2005. Magnetic orientation and magnetoreception in birds and other animals.
1285 *Journal of Comparative Physiology A* 191:675-693
- 1286 Wingfield JC, Maney DL, Breuner CW, Jacobs JD, Lynn S, Ramenofsky M, Richardson RD. 1998. Ecological
1287 bases of hormone—behavior interactions: the “emergency life history stage”. *American Zoologist*
1288 38:191-206
- 1289 Woodward G, Perkins DM, Brown LE. 2010. Climate change and freshwater ecosystems: impacts across
1290 multiple levels of organization. *Philosophical Transactions of the Royal Society B: Biological Sciences*
1291 365:2093-2106
- 1292 Wydoski RS, Wick EJ. 2000. Flooding and aquatic ecosystems. In Wohl, E.E. *Inland flood hazards:*
1293 *human, riparian, and aquatic communities.* Cambridge University Press, UK. 238-268.
- 1294
- 1295 Zellar R, Pulkkinen A, Moore K, Rousseaux CS, Reeb D. 2021. Oceanic and Atmospheric Correlations to
1296 Cetacean Mass Stranding Events in Cape Cod, Massachusetts, USA. *Geophysical Research Letters* 20:48
- 1297 Zhang G, Cui P, Jin W, Zhang Z, Wang H, Bazai NA, Li Y, Liu D, Pasuto A. 2021. Changes in hydrological
1298 behaviours triggered by earthquake disturbance in a mountainous watershed. *Science of The Total*
1299 *Environment* 760:143349
- 1300 Zhang J, Connor T, Yang H, Ouyang Z, Li S, Liu J. 2018. Complex effects of natural disasters on protected
1301 areas through altering telecouplings. *Ecology and Society* 3:23
- 1302 Zink IC, Browder JA, Kelble CR, Stabenau E, Kavanagh C, Fratto ZW. 2020. Hurricane-mediated shifts in a
1303 subtropical seagrass associated fish and macroinvertebrate community. *Estuaries and Coasts* 43(5):
1304 1174-1193
- 1305
- 1306
- 1307
- 1308
- 1309
- 1310

1311 **Table 1.** Summary of how different types of disasters impact life in different aquatic realms, the relative
 1312 level of knowledge we have about those impacts, and key synthesis articles on the impacts of a given
 1313 type of disaster (where available – we only cite truly synthetic papers such as reviews).

1314

Type of Disaster	Aquatic Realm	Level of Knowledge	Key Syntheses
Floods	Inland waters (especially rivers), Estuaries	Very well studied, particularly in inland riverine systems	Wydoski and Wick 2000; Talbot et al. 2018; Merz et al. 2021
Droughts	Inland waters, Estuaries	Very well studied, particularly in inland rivers, lakes and wetlands; Much of the research based in Australia	Bond et al. 2008; Lake 2003, 2011
Wildfires	Inland waters, Estuaries, Coastal marine	Well studied with much recent research reflecting increasing intensity of such events	Gresswell 1999; Bixby et al. 2015; Gomez Isaza et al. 2022
Hurricanes (including cyclones and typhoons)	Inland waters, Estuaries, Coastal marine, Offshore marine	Well studied with much recent interest on impacts on vertebrate life	Waide 1991; Greening et al. 2006; Mallin and Corbett 2006; Wang et al. 2016
Tornadoes	Inland waters, Estuaries, Coastal marine	Poorly studied – a few empirical studies	NA
Dust storms	Inland waters, Estuaries, Coastal marine, Offshore marine	Some research although tends to focus on long-term changes in water chemistry rather than biological impacts; Most research from Middle East region	Griffin and Kellogg 2004
Ice storms	Inland waters (usually small rivers)	Poorly studied – a few empirical studies	NA

Avalanches	Inland waters, Estuaries, Coastal marine	Poorly studied – a few empirical studies	Muller and Straub 2016
Landslides	Inland waters, Estuaries, Coastal marine	Some research on aquatic impacts although mostly from the Pacific Northwest of North America	Geertsema and Pojar 2007; Geertsema et al. 2009
Volcanic eruptions	Inland waters, Estuaries, Coastal marine	Extensive research but typically focused around long-term study sites (e.g., Mount St. Helens in the USA)	Swanson and Crisafulli 2018; Carrillo and Díaz-Villanueva 2021
Earthquakes	Inland waters, Estuaries, Coastal marine, Offshore marine	Poorly studied	Freund and Stolc 2013
Limnic eruptions	Inland waters (usually volcanic lakes)	Some research with focus on lakes that are subject to such events – mostly in Africa and South America	Rouwet et al. 2014
Tsunamis	Inland waters (near coasts), Estuaries, Coastal marine, Offshore marine	Increasing body of research following relatively recent tsunamis over last decade, mostly in Asia	Urabe and Nakashizuka 2016
Cosmic events	Inland waters, Estuaries, Coastal marine, Offshore marine	Relatively little research on this topic, mostly lab-based	NA

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326 **Figure Captions**

1327 **Figure 1.** Natural disasters have diverse impacts on aquatic ecosystems as highlighted in the examples
1328 illustrated here.

1329

Draft

EFFECTS OF NATURAL DISASTERS ON AQUATIC LIFE

